

Decreasing Universe: The 'Dark Matter' Effect

João Carlos Holland de Barcellos, University of São Paulo

Abstract

After a brief review of the *Decreasing Universe* (D.U.) model, we will derive a formula and the graph for the *apparent velocity* of stars (of a Messier_33 galaxy) as a function of their radial distance from the galactic center. This formula reproduces the same velocity curve traditionally attributed to dark matter.

Keywords: Dark Matter, Hubble's Law, Dark Energy, Decreasing Universe, Universe Expansion, Messier 33.

1. Introduction

The *Decreasing Universe* (D.U.) theory posits that the gravitational field continuously contracts space and all objects within it. This effect offers an alternative explanation for two key cosmological phenomena:

1. The **apparent accelerated expansion** of galaxies (without invoking dark energy).
2. The **anomalous orbital velocities** of stars (without requiring dark matter).

While counterintuitive, this idea aligns with established relativistic effects:

- **Equivalence Principle:** In an accelerating elevator, space and objects inside contract in the direction of motion.
- **Schwarzschild Metric:** Near a massive body, radial distances are compressed.

Though D.U. does not directly derive from these effects, they illustrate that gravitational contraction is a plausible mechanism.

2. Dark Energy and Universe Expansion

The D.U. model explains the apparent cosmic acceleration as a consequence of **local** space contraction. Since Earth-bound measuring instruments also shrink, distant galaxies appear to recede faster than expected—even if their actual motion is unchanged. This contraction-based interpretation reproduces **Hubble's Law**, eliminating the need for dark energy.

For details, see:

- *Derivation of Hubble's Law and its relation to dark energy and dark matter* [01].
 - *Decreasing Universe: Distance as a function of redshift* [02].
-

3. Dark Matter and Galactic Rotation Curves

Dark matter is traditionally invoked to explain why stars in galactic outskirts orbit faster than predicted by Newtonian mechanics. The D.U. model offers an alternative: **differential gravitational contraction** within galaxies.

- **Near the galactic center:** Higher contraction rates shorten emitted photon wavelengths more severely.
- **Farther out:** Reduced contraction yields less wavelength compression.

Since stellar velocities are measured via redshift (z), this differential mimics the Doppler effect, making outer stars appear faster—**without dark matter**.

4. Testing D.U. Against Λ CDM

The standard Λ CDM model relies on dark energy and dark matter. D.U. makes a testable prediction:

- **D.U.:** More massive galaxies at the same distance should exhibit **lower redshifts** due to stronger contraction.
- **Λ CDM:** Predicts the opposite (gravitational redshift increases with mass).

Empirical data support D.U. and contradict Λ CDM ([\[031\]](#)).

5. The Dark Matter Effect

5.1 Core Hypothesis

The gravitational field (g) contracts space **continuously**, with the rate depending on g 's intensity. From [\[01\]](#), the contraction formula is:

$$L(t) = \frac{L_0}{2^{\Delta t/T_j}} \quad (F1)$$

(Contraction formula from the point of view of a non-shrinking observer)

Where:

$L(t)$ = Measure of L_0 (decreasing) by a "Sidereal Observer".

t_0 = Initial time (arbitrary)

L_0 = Length measured in $t=t_0$

T_j = Jocaxian Time (Time to shrink to half)

$\Delta t = t - t_0$

Rewritten using the *Jocaxian Factor* (F_j):

$$F_j(\Delta t) = 2^{\Delta t/T_j} \quad (F2)$$

Jocaxian's Factor

We can also rewrite the same Jocaxian Factor (F_j) in a more friendly way:

$$F_j(\Delta t) = e^{\left(\ln(2) \cdot \frac{\Delta t}{T_j}\right)} \quad (\text{F3})$$

We can rewrite (F1):

$$L = L_0 / F_j(\Delta t) \quad (\text{F2})$$

5.2 Gravitational Dependence

We know that the rate of contraction depends on the gravitational field (g). If the gravitational field increases, the contraction will be faster, that is, the time for space to fall by half decreases.

Therefore, we will include a new hypothesis:

The contraction time is inversely proportional to the intensity of the gravitational field.

So, we have:

$$T_j(g) = T_j \cdot \frac{g_0}{g} \quad (\text{F3})$$

Where

T_j = Jocaxian Time (Time to Length contract to half)

$g_0 = GM_t / R_t^2$ (gravitational field at Earth (Milk Way)).

$g = GM / R^2$ (gravitational field at distance R from galactic center).

This leads to the **generalized Jocaxian Factor**:

$$F_j(\Delta t, M, R) = e^{\frac{J_0 M \Delta t}{R^2}} \quad (\text{F4})$$

Where

$$J_0 = \frac{H_0 R_t^2}{M_t} \quad (\text{F5})$$

(Jocax Constant $\approx 1,37\text{E-}18 \text{ m}^2 \cdot \text{s}^{-1} \cdot \text{kg}^{-1}$).

Where

H_0 = Hubble Constant ($\approx 2,2\text{E-}18 \text{ s}^{-1}$)


R_t = Distance from Earth to center of Milk Way ($\approx 2,5\text{E}20 \text{ m}$)

M_t = Mass of Milk Way ($\approx 1\text{E}41 \text{ Kg}$)

6. Galactic Rotation Curves

Dark Matter is a hypothesis created to explain the anomalous speed of stars and gases orbiting galaxies. This speed appears to increase as the orbit gets further away from the center of the galaxy. This apparent increase in speed contradicts Newton's laws, which indicate lower speeds for stars orbiting further away from the center of the galaxy.

The following graph -Rotation Curve for the Messier Galaxy_33- (Taken by observational measures) illustrates this behavior:



Rotation curve of spiral galaxy Messier 33 plotted by the observational measures [4]

6.1 Photon Redshift Calculation

To understand what happens to the Photon until it reaches Earth, we must consider that:

a) Both the galaxy from which the photon originated and the Milky Way have been contracting since their formation. We will consider that both began the contraction at the same time T_0 . (We will take T_0 as the time of the formation of the galaxy until the present day: approximately **13 billion years** ($=4E16$ s).

b) The contraction rate, as we have seen, depends on the mass of the galaxy and therefore may have different contraction rates. The *Jocaxian Factor* will consider these differences.

c) It is important to note, as we saw previously, that the *contraction rate* of the star that emits the photon depends on its distance (R) from the center of the galaxy. Thus, the redshift of the outgoing photon (considering only this factor), will be greater the further away the star is from its origin.

d) Of course, part of the redshift will be due to the *rotation velocity* of the star, which is greater the closer the star is to the center.

e) This difference in the outgoing redshift as a function of the distance from the center of the galaxy, will define the "dark matter effect" and its peculiar curve.

f) Let us consider that T_1 is the instant at which the photon is emitted from the source galaxy towards Earth. When it leaves the source galaxy, it ceases to contract, since the gravitational field of the galaxy becomes negligible.

g) We will calculate the apparent velocity of the star (in Messier 33 galaxy) (as a function of its distance from the center) *considering the redshift up to the moment it leaves the galaxy*. (We will **not** consider the "dark energy" effect, which is the contraction of our galaxy **after** the photon leaves the emitting galaxy until the Earth).

6.2- Let's do the math

To simplify, let's assume that the galaxies formed at the *same time*. In this case, the photons began to contract differently for each galaxy with different masses.

Redshift is the variation in wavelength in relation to a standard on Earth.

Initially, shortly after their formation, both galaxies (the emitter (Messier_33, with a lower mass) and the receiver (the Milky Way, with a higher mass) emitted photons at the same wavelength λ_0 .

After about 13 billion years (T_0), the contraction rates were different and, therefore, the wavelength decreased differently for both galaxies.

In the Milky Way, the photon after T_0 will have a shorter wavelength:

$$\lambda_{0}^T = \lambda_0 / F_J^T(T_0) = \lambda_0 / \exp(H_0 T_0)$$

Where:

λ_0 = Wave length at beginning

λ_{0}^T = wave length of photon in Earth after T_0

F_J^T = Jocaxian Factor on Earth

T_0 = Time from beginning of Galaxy until Now (13G Years)

In Messier_33 the contraction rate is different, and we must use the general F_j :

$$\lambda_{0}^M = \lambda_0 / F_j(T_0, M, R)$$

Where:

λ_{0}^M = wave length of photon in Messier_33 Earth **after** T_0

$F_j(T_0, M, R)$ = generalized Jocaxian Factor

M = Messier_33 Mass (1,4E39 Kg)

R = Distance of star to center of Messier_33

However, the Messier_33 star that emits the photon, will present a redshift that depends on its *rotation speed*:

$$V = cz$$

Where c = Speed of light and z = redshift

However

$$V = \text{SQRT}(MG/R)$$

$$z = V/c = (\lambda^M_1 / \lambda^M_0 - 1)$$

Where

λ^M_1 = wave length of photon in Messier33 due rotation velocity

$$\lambda^M_1 = \lambda^M_0 (\text{SQRT}(MG/R)/c + 1)$$

The final redshift relative to the Milky Way will be:

$$Z_f = (\lambda^M_1 - \lambda^T_0) / \lambda^T_0$$

So,

$$Z_f = (\lambda^M_0 (\text{SQRT}(MG/R)/c + 1) / \lambda^T_0 - 1$$

Replacing we will have:

$$Z_f = (\exp(H_0 T_0) / F_j(T_0, M, R)) (\text{SQRT}(MG/R)/c + 1) - 1$$

But $v = cz$, then **the speed measured on Earth** (without considering the effect of "dark energy" = Our contraction during the photon's journey), will be:

$$V = c \{ (\exp(H_0 T_0) / F_j(T_0, M, R)) (\text{SQRT}(MG/R)/c + 1) - 1 \}$$

Substituting $F_j(T_0, M, R)$ we will finally have the galaxy's rotation curve:

$$V_t = c \left(\left(\frac{1}{c} \sqrt{\frac{MG}{R}} + 1 \right) e^{H_0 T_0 - \frac{J_0 M T_0}{R^2}} - 1 \right)$$

Curva de Rotação da Galáxia (Velocidade aparente)

Where:

C = Speed of Light ($3E8$ m/s)

M = Mass of the Emitting Galaxy ($1,4E40$ Kg)

G = Gravitational Constant ($6.7E-11$)

R = Distance from the Star to the center of the galaxy

H_0 = Hubble Constant ($2.2E-18$)

$T_0 = \text{Contraction Time (13 billion years} = 4E16\text{s)}$

$J_0 = \text{Jocax Constant (1,37E-18)}$


6.2-For the galaxy Messier 33

Substituting the constant values we will have the Speed of the Messier Galaxy Star Observed from Earth:

$$V_t = 3E5 \left(\left(\sqrt{\frac{2,4E6}{R}} + 1 \right) e^{(0,088 - \frac{7,7E38}{R^2})} - 1 \right)$$

Star Speed in km/s as a function of distance to the Messier Galaxy_33

We can finally plot the rotation curve for Messier_33 predicted by the "Decreasing Universe" theory:



7. Conclusion

The D.U. model provides a unified framework for:

- **Apparent cosmic acceleration** (replacing dark energy).
- **Anomalous rotation curves** (replacing dark matter).

By reinterpreting gravitational contraction, it challenges the need for undetected exotic components in cosmology.

References

- [01] [Derivation of Hubble's Law](#)
- [02] [Distance as a function of redshift](#)
- [03] [Redshifts Refute \$\Lambda\$ CDM](#)
- [04] [Messier 33 Rotation Curve](#)
- [05] [Galaxies Grow Hotter with Age](#)