

# First-order swap structures semantics for some Logics of Formal Inconsistency

Marcelo E. Coniglio<sup>1</sup>, Aldo Figallo-Orellano<sup>2</sup> and Ana C. Golzio<sup>3</sup>

<sup>1</sup>Institute of Philosophy and the Humanities (IFCH) and  
Centre for Logic, Epistemology and The History of Science (CLE),  
University of Campinas (UNICAMP), Campinas, SP, Brazil.  
E-mail: coniglio@unicamp.br

<sup>2</sup>Departamento de Matemática, Universidad Nacional del Sur (UNS),  
Bahia Blanca, Argentina and  
Centre for Logic, Epistemology and The History of Science (CLE),  
University of Campinas (UNICAMP), Campinas, SP, Brazil.  
E-mail: aldofigallo@gmail.com

<sup>3</sup>São Paulo State University (Unesp), Marilia Campus, Brazil.  
E-mail: anaclaudiagolzio@yahoo.com.br

## Abstract

The logics of formal inconsistency (**LFI**s, for short) are paraconsistent logics (that is, logics containing contradictory but non-trivial theories) having a consistency connective which allows to recover the *ex falso quodlibet* principle in a controlled way. The aim of this paper is considering a novel semantical approach to first-order **LFI**s based on Tarskian structures defined over swap structures, a special class of multialgebras. The proposed semantical framework generalizes previous approaches to quantified **LFI**s presented in the literature. The case of **QmbC**, the simpler quantified **LFI** expanding classical logic, will be analyzed in detail. An axiomatic extension of **QmbC** called **QLFI**<sub>o</sub> is also studied, which is equivalent to the quantified version of da Costa and D'Ottaviano 3-valued logic **J3**. The semantical structures for this logic turn out to be Tarskian structures based on twist structures. The expansion of **QmbC** and **QLFI**<sub>o</sub> with a standard equality predicate is also considered.

*Keywords:* First-order logics; Logics of formal inconsistency; Paraconsistent logics; Swap structures; Non-deterministic matrices; Twist structures

## 1 Introduction

A logic is said to be *paraconsistent* if it contains in its language a negation and it has a contradictory theory (with respect to such negation) which is non-trivial. Such negation is called a *paraconsistent* or *non-explosive* negation. This is why paraconsistent logics are said to be *tolerant to contradictions*. The class of paraconsistent logics known as

logics of formal inconsistency (**LFI**s, for short) was introduced by W. Carnielli and J. Marcos in [11]. In its simplest form, they have a non-explosive negation  $\neg$ , as well as a (primitive or derived) *consistency connective*  $\circ$  which allows to recover the explosion law in a controlled way.

Defining interesting and elucidative semantics for paraconsistent logics is a challenging task for paraconsistentists, taking into account that, in general, these logics are not algebraizable by means of the standard techniques. Several kinds of semantics of non-deterministic character were proposed for these logics, in particular for **LFI**s. Among them, the non-deterministic matrices (or Nmatrices), introduced by A. Avron and I. Lev in [4] (see also [5]) constitutes an interesting and useful semantical framework for dealing with such logics.

The situation is even more delicate in the case of first-order paraconsistent logics. The aim of this paper is considering a novel semantical approach for first-order **LFI**s based on Tarskian structures defined over a special class of multialgebras called *swap structures*, which were introduced by Carnielli and Coniglio in [7]. The proposed semantical framework for quantified **LFI**s generalizes previous approaches in the literature.

In order to understand what lies behind the non-deterministic semantical framework for first-order **LFI**s proposed here, let us recall briefly the standard lattice-theoretic algebraic approach to first-order classical logic (**CFOL**, in short). To simplify the exposition, let us consider only sentences instead of formulas with free variables. Let us begin by recalling that the standard semantics for **CFOL** is given by Tarskian first-order structures. Given such a structure  $\mathbf{A}$ , in order to evaluate sentences of the given language, it is used the standard two-valued logical matrix for propositional classical logic **CPL** based on the two-element Boolean algebra  $\mathcal{A}_2$  with domain  $A_2 = \{0, 1\}$ , expanded by quantification operators  $\tilde{Q} : (\mathcal{P}(A_2) - \{\emptyset\}) \rightarrow A_2$  for  $Q \in \{\forall, \exists\}$  given, respectively, by

$$\tilde{\forall}(X) = \bigwedge X \quad \text{and} \quad \tilde{\exists}(X) = \bigvee X.$$

It is known that this class of semantical structures can be enlarged by considering structures  $\mathfrak{A} = \langle U, I_{\mathfrak{A}} \rangle$  defined over a complete Boolean algebra  $\mathcal{A}$  instead of the two-element Boolean algebra  $\mathcal{A}_2$  (see, for instance, [28, Supplement–Section 8]). This means that  $I_{\mathfrak{A}}(P)$  is now a mapping  $I_{\mathfrak{A}}(P) : U^n \rightarrow A$  for any  $n$ -ary predicate symbol  $P$ , where  $A$  is the domain of  $\mathcal{A}$ . The interpretation of sentences in **CFOL** is then given in the context  $(\mathfrak{A}, \mathcal{M}_{\mathcal{A}})$ , where  $\mathcal{M}_{\mathcal{A}} = \langle \mathcal{A}, \{1\} \rangle$  constitutes a logical matrix for **CPL** in which 1 is the only designated value, and where the quantifiers are interpreted *mutatis mutandis* as in the case of  $\mathcal{A}_2$ . This kind of algebraic approach to quantified logics was firstly proposed by Mostowski in [27], and later on generalized by Henkin in [24] and Rasiowa and Sikorski (see [29, 28]). This approach was recently extended by Cintula and Noguera in [16] to first-order logics based on algebraizable logics in the sense of Blok and Pigozzi.

What is proposed here for **QmbC** is a generalization of the algebraic approach to **CFOL** mentioned above, in which the logical matrix  $\mathcal{M}_{\mathcal{A}}$  is replaced by a non-deterministic matrix  $\mathcal{M}(\mathcal{B})$  for **mbC**, where  $\mathcal{B}$  is a swap structure (a multialgebra of a special kind) for **mbC** which is induced by a complete Boolean algebra  $\mathcal{A}$ . That is, sentences of **QmbC** will be interpreted in contexts  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$  for any swap structure  $\mathcal{B}$  for **mbC**. The analogy with the semantics of **CFOL** is more accurate than it first appears: while **CFOL** is fully characterized by the class of contexts  $(\mathfrak{A}, \mathcal{M}_2)$  such that  $\mathcal{M}_2 = \langle \mathcal{A}_2, \{1\} \rangle$  (which is equivalent, up to presentation, to the class of standard Tarskian structures with the usual semantics), **QmbC** can be characterized by contexts  $(\mathfrak{A}, \mathcal{M}_5)$  defined over the 5-element Nmatrix  $\mathcal{M}_5$ , which is the one induced by the greatest swap structure defined over  $\mathcal{A}_2$ ,

as we shall see in Section 9. This is why semantical contexts for **QmbC** of the form  $(\mathfrak{A}, \mathcal{M}_5)$  are considered to be ‘classical’, in analogy to **CFOL**. It is worth noting that the semantics for **QmbC** given by the ‘classical’ contexts coincide, up to notational aspects, with the non-deterministic semantics proposed by Avron and Zamansky in [6].

An interesting feature of the mutialgebraic semantics studied here is that, by adding axioms to a given **LFI**, conditions on the multioperations, and even on the domain of the swap structures, naturally arise. In particular, consider the 3-valued **LFI**<sub>o</sub>, which is equivalent (up to language) to several well-known 3-valued logics such as da Costa-D’Ottaviano logic **J3**. As shown in [14], the swap structures for **LFI**<sub>o</sub>, which is an axiomatic extension of **mbC**, are deterministic, and as such they become twist structures, that is, ordinary agebras of a certain kind. From this, the first-order swap structures for **QLFI**<sub>o</sub>, the quantified extension of **LFI**<sub>o</sub>, become first-order twist structures. As we shall see, when the ‘classical’ structures (that is, the twist structures induced by the Boolean algebra  $\mathcal{A}_2$ ) are considered, the corresponding first-order structures are defined over the characteristic 3-valued logical matrix for **LFI**<sub>o</sub>, hence this semantics is equivalent (up to presentation) with the early 3-valued model theory for quantified **J3** introduced in [17], [18], [19] and [20].

This paper is organized as follows: in the first sections, the swap structures semantics for **mbC** will be recalled. In Section 6 a semantics based on swap structures for **QmbC** will be introduced, proving in the following sections the corresponding soundness and completeness theorems. As we shall see, the swap structures semantics generalizes the interpretation semantics for **QmbC** given in [7], as well as the Nmatrix semantics proposed in [6]. The extension of **QmbC** by adding a standard equality predicate will be analyzed in Sections 10 and 11. The analysis of **QLFI**<sub>o</sub>, whose semantics is based on twist structures, will be done in Sections 12 to 14. Some conclusions are given in the last section.

## 2 The logic mbC

In this section, the notion of logics of formal inconsistency will be recalled, and the basic **LFI** called **mbC** will be briefly described. Let  $\Sigma'$  be a propositional signature, and assume a denumerable set  $\mathcal{V} = \{p_1, p_2, \dots\}$  of propositional variables. The propositional language generated by  $\Sigma'$  from  $\mathcal{V}$  will be denoted by  $\mathcal{L}_{\Sigma'}$

**Definition 2.1.** Let  $\mathbf{L} = \langle \Sigma', \vdash \rangle$  be a Tarskian, finitary and structural logic defined over a propositional signature  $\Sigma'$ , which contains a negation  $\neg$ , and let  $\circ$  be a (primitive or defined) unary connective. Then,  $\mathbf{L}$  is said to be a *logic of formal inconsistency* with respect to  $\neg$  and  $\circ$  if the following holds:<sup>1</sup>

- (i)  $\varphi, \neg\varphi \not\vdash \psi$  for some  $\varphi$  and  $\psi$ ;
- (ii) there are two formulas  $\alpha$  and  $\beta$  such that
  - (ii.a)  $\circ\alpha, \alpha \not\vdash \beta$ ;
  - (ii.b)  $\circ\alpha, \neg\alpha \not\vdash \beta$ ;
- (iii)  $\circ\varphi, \varphi, \neg\varphi \vdash \psi$  for every  $\varphi$  and  $\psi$ .

Condition (iii) states that *ex falso quodlibet* is controllably recovered in **LFIs** by assuming that the contradictory formula  $\varphi$  is consistent, i.e.,  $\circ\varphi$ .

---

<sup>1</sup>The original definition of **LFIs** allows to consider a set  $\bigcirc(p)$  of formulas depending on a single propositional variable  $p$ , instead of a single formula  $\circ p$ . See [11, 9, 7].

**Definition 2.2.** ([7, Definition 2.1.12]) Let  $\Sigma = \{\wedge, \vee, \rightarrow, \neg, \circ\}$  be a propositional signature. The calculus **mbC** over the propositional language  $\mathcal{L}_\Sigma$  is defined by means of the following Hilbert calculus:

**Axiom schemas:**

- (A1)  $\alpha \rightarrow (\beta \rightarrow \alpha)$
- (A2)  $(\alpha \rightarrow (\beta \rightarrow \gamma)) \rightarrow ((\alpha \rightarrow \beta) \rightarrow (\alpha \rightarrow \gamma))$
- (A3)  $\alpha \rightarrow (\beta \rightarrow (\alpha \wedge \beta))$
- (A4)  $(\alpha \wedge \beta) \rightarrow \alpha$
- (A5)  $(\alpha \wedge \beta) \rightarrow \beta$
- (A6)  $\alpha \rightarrow (\alpha \vee \beta)$
- (A7)  $\beta \rightarrow (\alpha \vee \beta)$
- (A8)  $(\alpha \rightarrow \gamma) \rightarrow ((\beta \rightarrow \gamma) \rightarrow ((\alpha \vee \beta) \rightarrow \gamma))$
- (A9)  $\alpha \vee (\alpha \rightarrow \beta)$
- (A10)  $\alpha \vee \neg\alpha$
- (A11)  $\circ\alpha \rightarrow (\alpha \rightarrow (\neg\alpha \rightarrow \beta))$

**Inference rule:**

$$(MP) \frac{\alpha \quad \alpha \rightarrow \beta}{\beta}$$

Observe that **mbC** is obtained from positive classical logic by adding axioms (A10) and (A11) governing the new connectives  $\neg$  (paraconsistent negation) and  $\circ$  (consistency operator). It is easy to see that **mbC** is an **LFI**. Indeed, it is the least **LFI** which contains propositional classical logic **CPL** (see Remark 8.2).

### 3 Swap structures for mbC

It is well-known that **mbC**, as well as several axiomatic extensions of it, are neither algebraizable (see [11, Section 3.12]), nor characterizable by a single finite logical matrix (see for instance [9, Theorems 121 and 125]). In this section a non-deterministic semantics for **mbC** based on multialgebras called *swap structures*, introduced in [7, Chapter 6], will be briefly recalled.

**Definition 3.1.** Let  $\Omega$  be a propositional signature. A *multialgebra* (or *hyperalgebra*) over  $\Omega$  is a pair  $\mathcal{A} = \langle A, \sigma \rangle$  such that  $A$  is a nonempty set (the *universe* or *support* of  $\mathcal{A}$ ) and  $\sigma$  is a mapping assigning to each  $n$ -ary connective  $c$ , a function (called *multioperation* or *hyperoperation*)  $c^{\mathcal{A}} : A^n \rightarrow (\mathcal{P}(A) - \{\emptyset\})$ . In particular,  $\emptyset \neq c^{\mathcal{A}} \subseteq A$  if  $c$  is a constant symbol.

**Definition 3.2.** Let  $\Omega$  be a propositional signature. A *non-deterministic matrix* (or *Nmatrix*) is a pair  $\mathcal{M} = \langle \mathcal{A}, D \rangle$  such that  $\mathcal{A} = \langle A, \sigma \rangle$  is a multialgebra over  $\Omega$  with support  $A$ , and  $D$  is a subset of  $A$ . The elements in  $D$  are called *designated* elements.

**Notation 3.3.** Let  $\mathcal{A}$  be a Boolean algebra with domain  $A$ . If  $x \in A \times A \times A$  then  $(x)_i$  (or simply  $x_i$ ) will denote the  $i$ th-projection of  $x$ , that is,  $\pi_i(x)$ , where  $\pi_i$  is the  $i$ th-canonical projection for  $i = 1, 2, 3$ .

**Definition 3.4.** ([7, Definition 6.4.1]) Let  $\mathcal{A} = (A, \wedge, \vee, \rightarrow, 0, 1)$  be a Boolean algebra, and  $B_{\mathcal{A}} = \{x \in A \times A \times A : x_1 \vee x_2 = 1 \text{ and } x_1 \wedge x_2 \wedge x_3 = 0\}$ . A *swap structure for  $\mathbf{mbC}$  over  $\mathcal{A}$*  is any multialgebra  $\mathcal{B} = (B, \tilde{\wedge}, \tilde{\vee}, \tilde{\rightarrow}, \tilde{\sim}, \tilde{\circ})$  over  $\Sigma$  such that  $B \subseteq B_{\mathcal{A}}$  and where the multioperations satisfy the following, for every  $x$  and  $y$  in  $B$ :

- (i)  $\emptyset \neq x \# y \subseteq \{z \in B : z_1 = x_1 \# y_1\}$ , for each  $\# \in \{\wedge, \vee, \rightarrow\}$ ;
- (ii)  $\emptyset \neq \tilde{\sim}x \subseteq \{z \in B : z_1 = x_2\}$ ;
- (iii)  $\emptyset \neq \tilde{\circ}x \subseteq \{z \in B : z_1 = x_3\}$ .

**Definition 3.5.** The *full swap structure for  $\mathbf{mbC}$  over  $\mathcal{A}$* , denoted by  $\mathcal{B}_{\mathcal{A}}$ , is the unique swap structure for  $\mathbf{mbC}$  over  $\mathcal{A}$  with domain  $B_{\mathcal{A}}$ , in which ‘ $\subseteq$ ’ is replaced by ‘ $=$ ’ in items (i)-(iii) of Definition 3.4.

Observe that  $\mathcal{B}_{\mathcal{A}}$  is the greatest swap structure for  $\mathbf{mbC}$  over  $\mathcal{A}$  (see [14]). The elements of a given swap structure are called *snapshots*. This terminology is inspired by its use in computer science to refer to states. Accordingly, a triple  $(a, b, c)$  of a swap structure  $\mathcal{B}$  keeps track simultaneously of the value  $a$  of a given formula  $\varphi$ , a possible value  $b$  for  $\neg\varphi$ , and a possible value  $c$  for  $\circ\varphi$ .

Given that any swap structure is a multialgebra, the consequence relation over swap structures will be defined by means of non-deterministic matrices, in analogy with the corresponding notion for twist structures.

**Definition 3.6.** ([7, Definition 6.4.3]) Given a Boolean algebra  $\mathcal{A}$  and a swap structure  $\mathcal{B}$  for  $\mathbf{mbC}$  over  $\mathcal{A}$  with domain  $B$ , let  $D_B = \{x \in B : x_1 = 1\}$  be the set of *designated* elements. The non-deterministic matrix associated to  $\mathcal{B}$  is  $\mathcal{M}(\mathcal{B}) = (\mathcal{B}, D_B)$  (or simply  $\mathcal{M}(\mathcal{B}) = (\mathcal{B}, D)$ ). The Nmatrix associated to  $\mathcal{B}_{\mathcal{A}}$  will be denoted by  $\mathcal{M}_{\mathcal{A}}$ . The class of all the Nmatrices defined by swap structures for  $\mathbf{mbC}$  will be denoted by  $\mathbb{M}_{\mathbf{mbC}}$ , that is:

$$\mathbb{M}_{\mathbf{mbC}} = \{\mathcal{M}(\mathcal{B}) : \mathcal{B} \text{ is a swap structure for } \mathbf{mbC} \text{ over } \mathcal{A}, \text{ for some } \mathcal{A}\}.$$

**Definition 3.7.** Let  $\mathcal{M}(\mathcal{B}) \in \mathbb{M}_{\mathbf{mbC}}$ . A *valuation* over  $\mathcal{M}(\mathcal{B})$  is a function  $v : \mathcal{L}_{\Sigma} \rightarrow |\mathcal{B}|$  such that, for every  $\varphi, \psi \in \mathcal{L}_{\Sigma}$ :

- (i)  $v(\varphi \# \psi) \in v(\varphi) \# v(\psi)$ , for every  $\# \in \{\wedge, \vee, \rightarrow\}$ ;
- (ii)  $v(\neg\varphi) \in \tilde{\sim}v(\varphi)$ ;
- (iii)  $v(\circ\varphi) \in \tilde{\circ}v(\varphi)$ .

**Definition 3.8.** Let  $\mathcal{M}(\mathcal{B}) \in \mathbb{M}_{\mathbf{mbC}}$ , and let  $\Gamma \cup \{\varphi\} \subseteq \mathcal{L}_{\Sigma}$ . We say that  $\varphi$  is a *consequence* of  $\Gamma$  in  $\mathcal{M}(\mathcal{B})$ , denoted by  $\Gamma \models_{\mathcal{M}(\mathcal{B})} \varphi$ , if the following holds: for every valuation  $v$  over  $\mathcal{M}(\mathcal{B})$ , if  $v[\Gamma] \subseteq D$  then  $v(\varphi) \in D$ . In particular,  $\varphi$  is *valid* in  $\mathcal{M}(\mathcal{B})$ , denoted by  $\models_{\mathcal{M}(\mathcal{B})} \varphi$ , if  $v(\varphi) \in D$  for every valuation  $v$  over  $\mathcal{M}(\mathcal{B})$ . The *swap consequence relation*  $\models_{\mathbb{M}_{\mathbf{mbC}}}$  for  $\mathbf{mbC}$  is given by:  $\Gamma \models_{\mathbb{M}_{\mathbf{mbC}}} \varphi$  whenever  $\Gamma \models_{\mathcal{M}(\mathcal{B})} \varphi$  for every  $\mathcal{M}(\mathcal{B}) \in \mathbb{M}_{\mathbf{mbC}}$ .

**Theorem 3.9.** (Adequacy of  $\mathbf{mbC}$  w.r.t. swap structures, [7, Theorem 6.4.8] and [14, Theorem 7.1]) *For every  $\Gamma \cup \{\varphi\} \subseteq \mathcal{L}_{\Sigma}$ :  $\Gamma \vdash_{\mathbf{mbC}} \varphi$  iff  $\Gamma \models_{\mathbb{M}_{\mathbf{mbC}}} \varphi$ .*

## 4 The 5-valued characteristic Nmatrix $\mathcal{M}_5$ for mbC

It is illustrative to compare Theorem 3.9 with the adequacy of classical propositional logic **CPL** w.r.t. Boolean algebras semantics. As it is well known, it is enough to consider just one Boolean algebra to semantically characterize **CPL**, namely the two-element Boolean algebra  $\mathcal{A}_2$  with domain  $A_2 = \{0, 1\}$  and the associated logical matrix with 1 as designated value. In the case of swap structures semantics, it is enough to consider the Nmatrix  $\mathcal{M}_5 = \mathcal{M}(\mathcal{B}_{\mathcal{A}_2})$  induced by the full swap structure  $\mathcal{B}_{\mathcal{A}_2}$  defined over  $\mathcal{A}_2$ . The Nmatrix  $\mathcal{M}_5$  was originally introduced by A. Avron in [1] to semantically characterize **mbC**. Observe that  $B_{\mathcal{A}_2} = \{T, t, t_0, F, f_0\}$  where  $T = (1, 0, 1)$ ,  $t = (1, 1, 0)$ ,  $t_0 = (1, 0, 0)$ ,  $F = (0, 1, 1)$ , and  $f_0 = (0, 1, 0)$ . The set D of designated elements of  $\mathcal{M}_5$  is  $D = \{T, t, t_0\}$ , while  $ND = \{F, f_0\}$  is the set of non-designated truth-values. The multioperations proposed by Avron over the set  $B_{\mathcal{A}_2}$  coincide with the corresponding ones for  $\mathcal{B}_{\mathcal{A}_2}$ , and so his 5-valued Nmatrix coincides with  $\mathcal{M}(\mathcal{B}_{\mathcal{A}_2})$ . Observe that the swap structure of  $\mathcal{M}_5$  is defined as follows:

$\wedge^{\mathcal{M}_5}$	$T$	$t$	$t_0$	$F$	$f_0$
$T$	D	D	D	ND	ND
$t$	D	D	D	ND	ND
$t_0$	D	D	D	ND	ND
$F$	ND	ND	ND	ND	ND
$f_0$	ND	ND	ND	ND	ND

$\vee^{\mathcal{M}_5}$	$T$	$t$	$t_0$	$F$	$f_0$
$T$	D	D	D	D	D
$t$	D	D	D	D	D
$t_0$	D	D	D	D	D
$F$	D	D	D	ND	ND
$f_0$	D	D	D	ND	ND

$\rightarrow^{\mathcal{M}_5}$	$T$	$t$	$t_0$	$F$	$f_0$
$T$	D	D	D	ND	ND
$t$	D	D	D	ND	ND
$t_0$	D	D	D	ND	ND
$F$	D	D	D	D	D
$f_0$	D	D	D	D	D

	$\neg^{\mathcal{M}_5}$
$T$	ND
$t$	D
$t_0$	ND
$F$	D
$f_0$	D

	$\circ^{\mathcal{M}_5}$
$T$	D
$t$	ND
$t_0$	ND
$F$	D
$f_0$	ND

**Theorem 4.1.** (Adequacy of **mbC** w.r.t.  $\mathcal{M}_5$ , [1, Theorem 3.6]) *For every set of formulas  $\Gamma \cup \{\varphi\} \subseteq \mathcal{L}_\Sigma$ :  $\Gamma \vdash_{\mathbf{mbC}} \varphi$  iff  $\Gamma \models_{\mathcal{M}_5} \varphi$ .*

A new proof of this result was obtained in [7, Corollary 6.4.10], by using bivaluations for **mbC** in connection with the Nmatrix  $\mathcal{M}(\mathcal{B}_{\mathcal{A}_2})$ .

**Definition 4.2.** ([9, Definition 54]) A function  $\rho : \mathcal{L}_\Sigma \rightarrow \{0, 1\}$  is a *bivaluation for mbC* if it satisfies the following clauses:

- (**vAnd**)  $\rho(\alpha \wedge \beta) = 1$  iff  $\rho(\alpha) = \rho(\beta) = 1$
- (**vOr**)  $\rho(\alpha \vee \beta) = 1$  iff  $\rho(\alpha) = 1$  or  $\rho(\beta) = 1$
- (**vImp**)  $\rho(\alpha \rightarrow \beta) = 1$  iff  $\rho(\alpha) = 0$  or  $\rho(\beta) = 1$
- (**vNeg**)  $\rho(\neg\alpha) = 0$  implies  $\rho(\alpha) = 1$
- (**vCon**)  $\rho(\circ\alpha) = 1$  implies  $\rho(\alpha) = 0$  or  $\rho(\neg\alpha) = 0$ .

The consequence relation of **mbC** w.r.t. bivaluations, denoted by  $\models_{\mathbf{mbC}}^2$ , is given by:  $\Gamma \models_{\mathbf{mbC}}^2 \varphi$  iff  $\rho(\varphi) = 1$  for every bivaluation for **mbC** such that  $\rho[\Gamma] \subseteq \{1\}$ .

**Theorem 4.3.** (Adequacy of **mbC** w.r.t. bivaluations, [9, Theorems 56 and 61]) *For every set of formulas  $\Gamma \cup \{\varphi\} \subseteq \mathcal{L}_\Sigma$ :  $\Gamma \vdash_{\mathbf{mbC}} \varphi$  iff  $\Gamma \models_{\mathbf{mbC}}^2 \varphi$ .*

**Theorem 4.4.** ([7, Theorem 6.4.9]) *Any bivaluation  $\rho$  for **mbC** induces a valuation  $v^\rho$  over the Nmatrix  $\mathcal{M}_5$  given by  $v^\rho(\alpha) \stackrel{\text{def}}{=} (\rho(\alpha), \rho(\neg\alpha), \rho(\circ\alpha))$  such that, for every formula  $\alpha$ :  $\rho(\alpha) = 1$  iff  $v^\rho(\alpha) \in D$ .*

From the previous result, Theorem 4.1 follows easily (see [7, Corollary 6.4.10]). The characteristic Nmatrix  $\mathcal{M}_5$  of **mbC** can be considered as the ‘classical’ model of it, since it is based on the ‘classical’ Boolean algebra  $\mathcal{A}_2$ . In Section 9 it will be shown that **QmbC**, the first-order version of **mbC**, can be characterized by first-order structures defined over  $\mathcal{M}_5$ . These structures can be considered as ‘classical’ in this sense.

## 5 The logic QmbC

In this section the first-order logic **QmbC**, introduced in [10] (see also [7]) as an extension of **mbC** to first-order languages, will be briefly recalled. In Section 6 a new semantics of first-order swap structures for **QmbC** will be defined.

**Definition 5.1.** Assume the propositional signature  $\Sigma = \{\wedge, \vee, \rightarrow, \neg, \circ\}$  for **mbC**, as well as the symbols  $\forall$  (universal quantifier) and  $\exists$  (existential quantifier), with the punctuation marks (commas and parenthesis). Let  $Var = \{v_1, v_2, \dots\}$  be a denumerable set of individual variables. A first-order signature  $\Theta$  for **QmbC** is composed by the following elements:

- a set  $\mathcal{C}$  of individual constants;
- for each  $n \geq 1$ , a set  $\mathcal{F}_n$  of function symbols of arity  $n$ ,
- for each  $n \geq 1$ , a set  $\mathcal{P}_n$  of predicate symbols of arity  $n$ .<sup>2</sup>

**Notation 5.2.** *Let  $\Theta$  be a first-order signature for **QmbC**. The sets of terms and formulas generated by  $\Theta$  from  $Var$  we will denote by  $Ter(\Theta)$  and  $For(\Theta)$ , respectively. The set of sentences (formulas without free variables) and the set of closed terms (terms without variables) over  $\Theta$  are denoted by  $Sen(\Theta)$  and  $CTer(\Theta)$ , respectively. Given a formula  $\varphi$ , the formula obtained from  $\varphi$  by substituting every free occurrence of a variable  $x$  by a term  $t$  will be denoted by  $\varphi[x/t]$ .*

The notions of *subformula*, *scope* of an occurrence of a quantifier in a formula, *free* and *bound* occurrences of a variable in a formula, and of *term free for a variable in a formula*, are the usual ones (see, for instance, [26]).

**Definition 5.3.** ([7, Definition 7.1.4]) Let  $\varphi$  and  $\psi$  be formulas. If  $\varphi$  can be obtained from  $\psi$  by means of addition or deletion of void quantifiers, or by renaming bound variables (keeping the same free variables in the same places), we say that  $\varphi$  and  $\psi$  are *variant* of each other.

**Definition 5.4.** ([7, Definition 7.1.5]) Let  $\Theta$  be a first-order signature. The logic **QmbC** is obtained from the Hilbert calculus **mbC** extended by the following axioms and rules:

**Axiom schemas:**

---

<sup>2</sup>It will be assumed, as usual, that  $\Theta$  has at least one predicate symbol, in order to have a non-empty set of formulas. For instance, it could be assumed from the beginning an equality predicate (see Section 10).

- (Ax12)  $\varphi[x/t] \rightarrow \exists x\varphi$ , if  $t$  is a term free for  $x$  in  $\varphi$
- (Ax13)  $\forall x\varphi \rightarrow \varphi[x/t]$ , if  $t$  is a term free for  $x$  in  $\varphi$
- (Ax14)  $\alpha \rightarrow \beta$ , whenever  $\alpha$  is a variant of  $\beta$

**Inference rules:**

- ( $\exists$ -In)  $\frac{\varphi \rightarrow \psi}{\exists x\varphi \rightarrow \psi}$ , where  $x$  does not occur free in  $\psi$
- ( $\forall$ -In)  $\frac{\varphi \rightarrow \psi}{\varphi \rightarrow \forall x\psi}$ , where  $x$  does not occur free in  $\varphi$

**Definition 5.5.** If  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ , then  $\Gamma \vdash_{\mathbf{QmbC}} \varphi$  will denote that there exists a derivation in  $\mathbf{QmbC}$  of  $\varphi$  from  $\Gamma$ .

In [7] it was proved that the logic  $\mathbf{QmbC}$  enjoys the Deduction meta-theorem (DMT), as usually presented in first-order logics:

**Theorem 5.6** (Deduction Meta-Theorem (DMT) for  $\mathbf{QmbC}$ ). *Suppose that there exists in  $\mathbf{QmbC}$  a derivation of  $\psi$  from  $\Gamma \cup \{\varphi\}$ , such that no application of the rules ( $\exists$ -In) and ( $\forall$ -In) have, as their quantified variables, free variables of  $\varphi$  (in particular, this holds when  $\varphi$  is a sentence). Then  $\Gamma \vdash_{\mathbf{QmbC}} \varphi \rightarrow \psi$ .*

## 6 First-Order Swap Structures

The traditional approach to first-order structures based on algebraic structures (see for instance [27, 24, 29]) will be adapted to swap structures semantics. Thus, from now on the Boolean algebras to be considered are assumed to be complete.<sup>3</sup>

**Definition 6.1.** Let  $\mathcal{M}(\mathcal{B}) = (\mathcal{B}, D)$  be a non-deterministic matrix defined by a swap structure  $\mathcal{B}$  for  $\mathbf{mbC}$  over a complete Boolean algebra  $\mathcal{A}$ , and let  $\Theta$  be a first-order signature (see Definition 5.1). A (first-order) *structure* over  $\mathcal{M}(\mathcal{B})$  and  $\Theta$  is pair  $\mathfrak{A} = \langle U, I_{\mathfrak{A}} \rangle$  such that  $U$  is a nonempty set (the domain of the structure) and  $I_{\mathfrak{A}}$  is an interpretation function which assigns:

- to each individual constant  $c \in \mathcal{C}$ , an element  $I_{\mathfrak{A}}(c)$  of  $U$ ;
- to each function symbol  $f$  of arity  $n$ , a function  $I_{\mathfrak{A}}(f) : U^n \rightarrow U$ ;
- to each predicate symbol  $P$  of arity  $n$ , a function  $I_{\mathfrak{A}}(P) : U^n \rightarrow |\mathcal{B}|$ .

From now on, the expressions  $c^{\mathfrak{A}}$ ,  $f^{\mathfrak{A}}$  and  $P^{\mathfrak{A}}$  will be used instead of  $I_{\mathfrak{A}}(c)$ ,  $I_{\mathfrak{A}}(f)$  and  $I_{\mathfrak{A}}(P)$ , for an individual constant symbol  $c$ , a function symbol  $f$  and a predicate symbol  $P$ , respectively.

**Definition 6.2.** Let  $\mathfrak{A}$  be a structure over  $\mathcal{M}(\mathcal{B})$  and  $\Theta$ . A function  $\mu : \text{Var} \rightarrow U$  is called an *assignment* over  $\mathfrak{A}$ .

---

<sup>3</sup>Instead of this we could consider arbitrary Boolean algebras, thus obtaining partial models in which some quantified formulas has no denotation because of the lack of infima and/or suprema of some subsets. Since every Boolean algebra can be completed (see Section 8), we decide to restrict the semantic to complete Boolean algebras. An interesting discussion concerning this topic can be found in [16, footnote 3].



**Definition 6.3.** Let  $\mathfrak{A}$  be a structure over  $\mathcal{M}(\mathcal{B})$  and  $\Theta$ , and let  $\mu : Var \rightarrow U$  be an assignment. For each term  $t$ , we define  $\llbracket t \rrbracket_\mu^{\mathfrak{A}}$  in  $U$  such that:

- $\llbracket c \rrbracket_\mu^{\mathfrak{A}} = c^{\mathfrak{A}}$  if  $c$  is an individual constant;
- $\llbracket x \rrbracket_\mu^{\mathfrak{A}} = \mu(x)$  if  $x$  is a variable;
- $\llbracket f(t_1, \dots, t_n) \rrbracket_\mu^{\mathfrak{A}} = f^{\mathfrak{A}}(\llbracket t_1 \rrbracket_\mu^{\mathfrak{A}}, \dots, \llbracket t_n \rrbracket_\mu^{\mathfrak{A}})$  if  $f$  is a function symbol of arity  $n$  and  $t_1, \dots, t_n$  are terms.

**Definition 6.4.** Consider a structure  $\mathfrak{A}$  over  $\mathcal{M}(\mathcal{B})$  and  $\Theta$ . The *diagram language* of  $\mathfrak{A}$  is the set of formulas  $For(\Theta_U)$ , where  $\Theta_U$  is the signature obtained from  $\Theta$  by adding a new individual constant  $\bar{a}$  for each element  $a$  of the domain  $U$  of  $\mathfrak{A}$ .

**Definition 6.5.** The structure  $\widehat{\mathfrak{A}} = \langle U, I_{\widehat{\mathfrak{A}}} \rangle$  over  $\Theta_U$  is the structure  $\mathfrak{A}$  over  $\Theta$  extended by  $I_{\widehat{\mathfrak{A}}}(\bar{a}) = a$  for every  $a \in U$ .

Observe that  $s^{\widehat{\mathfrak{A}}} = s^{\mathfrak{A}}$  if  $s$  is a symbol (individual constant, function symbol or predicate symbol) of  $\Theta$ .

**Notation 6.6.** For any formula  $\varphi$ ,  $FV(\varphi)$  will denote the set of free variables of  $\varphi$ . The set of (closed) sentences (formulas without free variables) of the diagram language of  $\mathfrak{A}$  will be denoted by  $Sen(\Theta_U)$ , and the set of terms and of closed terms over  $\Theta_U$  will be denoted by  $Ter(\Theta_U)$  and  $CTer(\Theta_U)$ , respectively.

**Remark 6.7.** Clearly, if  $t$  is a closed term then the value of  $\llbracket t \rrbracket_\mu^{\mathfrak{A}}$  does not depend on the assignment  $\mu$ , that is:  $\llbracket t \rrbracket_\mu^{\mathfrak{A}} = \llbracket t \rrbracket_{\mu'}^{\mathfrak{A}}$ , for every assignments  $\mu$  and  $\mu'$ . Thus, if  $t$  is a closed term we can write  $\llbracket t \rrbracket^{\mathfrak{A}}$  instead of  $\llbracket t \rrbracket_\mu^{\mathfrak{A}}$ , for any assignment  $\mu$ .

**Definition 6.8.** Let  $\widehat{\mathfrak{A}} = \langle U, I_{\widehat{\mathfrak{A}}} \rangle$  be as above. Any assignment  $\mu$  over  $\widehat{\mathfrak{A}}$  induces a function  $\widehat{\mu} : (Ter(\Theta_U) \cup For(\Theta_U)) \rightarrow (Ter(\Theta_U) \cup For(\Theta_U))$  given by  $\widehat{\mu}(s) = s[x_1/\overline{\mu(x_1)}, \dots, x_n/\overline{\mu(x_n)}]$ , if either  $s \in Ter(\Theta_U)$  such that  $Var(s) \subseteq \{x_1, \dots, x_n\}$  or  $s \in For(\Theta_U)$  such that  $FV(s) \subseteq \{x_1, \dots, x_n\}$ .

It is worth observing that  $\widehat{\mu}(\varphi) \in Sen(\Theta_U)$  if  $\varphi \in For(\Theta_U)$ , and  $\widehat{\mu}(t) \in CTer(\Theta_U)$  if  $t \in Ter(\Theta_U)$ .

The next step is to define the notion of interpretation (or denotation) of a formula  $\varphi \in For(\Theta_U)$  in a given (extended) structure  $\widehat{\mathfrak{A}}$  and assignment  $\mu$ , which could be denoted by  $\llbracket \varphi \rrbracket_\mu^{\widehat{\mathfrak{A}}}$  (being coherent with the previous notation). Is exactly at this point when non-determinism enters. Observe that, in the traditional (truth-functional or algebraic) first-order semantical approach, any structure and assignment induce together a (unique) denotation for any formula. In the present framework, this is also true for atomic formulas, since predicates are interpreted by means of functions, and taking into account that the denotation of any term is uniquely determined given a structure and an assignment. However, the denotation of complex formulas is possibly non-deterministic (i.e., ambiguous), given that it involves logical symbols (connectives and quantifiers) to be evaluated over a non-deterministic matrix. As happens with the propositional case, we are not interested in assigning sets of truth-values to single formulas: instead of this, valuations (*legal valuations*, in Avron and Lev's terminology) are used in order to *choose*, in a coherent way, a single truth-value for any formula.<sup>4</sup> The definition of (legal) valuations over

<sup>4</sup>As we shall see in Section 12, in the case of the first-order logic  $\mathbf{LF11}_o$ , which is an axiomatic extension of  $\mathbf{QmbC}$  based on a truth-functional 3-valued logic, any structure and assignment will determine a unique truth-value for any single formula. In this case, it will not necessary to consider valuations.

first-order swap structures involves an additional technical complication with respect to the propositional case: the validity of the Substitution Lemma – a crucial result which allows to substitute a universally quantified variable by any term free for such variable in a given formula – is far from being true in our non-deterministic environment. Indeed, this technical result is trivially true for first-order logics in which the semantics is obtained by algebraic manipulations over the interpretation of the subformulas of the formula being interpreted. Since in **QmbC** it is necessary to introduce the valuations as intermediaries between the formulas and the multioperators of the swap structures, such valuations must satisfy additional requirements in order to guarantee the validity of the Substitution Lemma (namely, clause (vi) in Definition 6.9 below). To summarize, in order to interpret formulas in the present non-deterministic framework, it is necessary a structure, an assignment, and a (first-order) valuation over the underlying swap structure, which will be called a **QmbC-valuation**.

Given an assignment  $\mu$  over a structure  $\mathfrak{A}$ , a variable  $x$  and  $a \in U$ , the assignment  $\mu_a^x$  over  $\mathfrak{A}$  is given by  $\mu_a^x(y) = a$ , if  $y = x$ , and  $\mu_a^x(y) = \mu(y)$  otherwise. Thus, the previous considerations lead us to the following notion:

**Definition 6.9.** (**QmbC-valuations**) Let  $\mathcal{M}(\mathcal{B}) = (\mathcal{B}, D)$  be a non-deterministic matrix defined by a swap structure  $\mathcal{B}$  for **mbC**, and let  $\mathfrak{A}$  be a structure over  $\Theta$  and  $\mathcal{M}(\mathcal{B})$ . A function  $v : \text{Sen}(\Theta_U) \rightarrow |\mathcal{B}|$  is a *valuation* for **QmbC** (or a **QmbC-valuation**) over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$ , if it satisfies the following clauses:

- (i)  $v(P(t_1, \dots, t_n)) = P^{\mathfrak{A}}(\llbracket t_1 \rrbracket^{\mathfrak{A}}, \dots, \llbracket t_n \rrbracket^{\mathfrak{A}})$ , if  $P(t_1, \dots, t_n)$  is atomic;<sup>5</sup>
- (ii)  $v(\#\varphi) \in \#v(\varphi)$ , for every  $\# \in \{\neg, \circ\}$ ;
- (iii)  $v(\varphi\#\psi) \in v(\varphi)\#v(\psi)$ , for every  $\# \in \{\wedge, \vee, \rightarrow\}$ ;
- (iv)  $v(\forall x\varphi) \in \{z \in |\mathcal{B}| : z_1 = \bigwedge \{\pi_1(v(\varphi[x/\bar{a}])) : a \in U\}\}$ ;
- (v)  $v(\exists x\varphi) \in \{z \in |\mathcal{B}| : z_1 = \bigvee \{\pi_1(v(\varphi[x/\bar{a}])) : a \in U\}\}$ ;
- (vi) Let  $t$  be free for  $z$  in  $\varphi$  and  $\psi$ ,  $\mu$  an assignment and  $b = \llbracket t \rrbracket_{\mu}^{\mathfrak{A}}$ . Then:
  - (vi.1) If  $v(\widehat{\mu}(\varphi[z/t])) = v(\widehat{\mu}(\varphi[z/\bar{b}]))$ , then  $v(\widehat{\mu}(\#\varphi[z/t])) = v(\widehat{\mu}(\#\varphi[z/\bar{b}]))$ , for every  $\# \in \{\neg, \circ\}$ ;
  - (vi.2) If  $v(\widehat{\mu}(\varphi[z/t])) = v(\widehat{\mu}(\varphi[z/\bar{b}]))$  and  $v(\widehat{\mu}(\psi[z/t])) = v(\widehat{\mu}(\psi[z/\bar{b}]))$ , then  $v(\widehat{\mu}(\varphi\#\psi[z/t])) = v(\widehat{\mu}(\varphi\#\psi[z/\bar{b}]))$ , for every  $\# \in \{\wedge, \vee, \rightarrow\}$ ;
  - (vi.3) Let  $x$  be such that  $x \neq z$  and  $x$  does not occur in  $t$ . If  $v(\widehat{\mu}_a^x(\varphi[z/t])) = v(\widehat{\mu}_a^x(\varphi[z/\bar{b}]))$ , for every  $a \in U$ , then  $v(\widehat{\mu}((Qx\varphi)[z/t])) = v(\widehat{\mu}((Qx\varphi)[z/\bar{b}]))$ , for every  $Q \in \{\forall, \exists\}$ ;
- (vii) If  $\varphi$  and  $\varphi'$  are variant, then  $v(\varphi) = v(\varphi')$ .

Observe that clause (i) in the previous definition is the only one that uses the information of the structure  $\mathfrak{A}$ , and it allows to interpret the atomic formulas. In order to obtain a single denotation for a complex formula, the valuation is used to choose (coherently) a denotation for the formula from the denotation of its components. Clause (vi) guarantees the validity of the Substitution Lemma, a crucial step for obtaining the soundness of the proposed semantics.

**Definition 6.10.** Let  $v$  be a **QmbC-valuation** over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$ . Given an assignment  $\mu$  over  $\mathfrak{A}$ , we define  $v_{\mu} : \text{For}(\Theta_U) \rightarrow |\mathcal{B}|$  as  $v_{\mu}(\varphi) = v(\widehat{\mu}(\varphi))$ .

**Definition 6.11.** Let  $\mathfrak{A}$  be a structure over  $\Theta$  and  $\mathcal{M}(\mathcal{B})$ . If  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta_U)$ ,  $\varphi$  is said to be a *semantical consequence* of  $\Gamma$  over  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$ , denoted by  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}(\mathcal{B}))} \varphi$ , if the following holds: for every **QmbC-valuation**  $v$  over  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$ , if  $v_{\mu}(\gamma) \in D$ , for every formula  $\gamma \in \Gamma$  and every assignment  $\mu$ , then  $v_{\mu}(\varphi) \in D$ , for every assignment  $\mu$ .

<sup>5</sup>For the notation used here, recall Remark 6.7.

**Definition 6.12.** (Semantical consequence relation in **QmbC** w.r.t. swap structures) If  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ ,  $\varphi$  is said to be a *semantical consequence* of  $\Gamma$  in **QmbC** w.r.t. first-order swap structures, denoted by  $\Gamma \models_{\mathbf{QmbC}} \varphi$ , if  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}(\mathcal{B}))} \varphi$  for every  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$ .

As mentioned in the Introduction, the semantical contexts  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$  for **QmbC** generalize the semantical contexts  $(\mathfrak{A}, \mathcal{M}_{\mathcal{A}})$  for first-order classical logic **CFOL**, where  $\mathcal{M}_{\mathcal{A}} = \langle \mathcal{A}, \{1\} \rangle$ . The latter, by its turn, generalize the class of standard Tarskian structures for **CFOL** with the usual semantics, by taking the two-element Boolean algebra  $\mathcal{A}_2$ .

## 7 Soundness of **QmbC** w.r.t. swap structures

In this section the soundness of **QmbC** w.r.t. first-order swap structures semantics for **QmbC** will be proved. As mentioned in the previous section, a key result for proving soundness is the Substitution Lemma, which can be proved easily by induction on the complexity of  $\varphi$ .

**Theorem 7.1.** (Substitution Lemma) *Let  $v$  be a **QmbC**-valuation over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$  and let  $\mu$  be an assignment. If  $t$  is a term free for  $z$  in  $\varphi$  and  $b = \llbracket t \rrbracket_{\mu}^{\mathfrak{A}}$ , then  $v_{\mu}(\varphi[z/t]) = v_{\mu}(\varphi[z/\bar{b}])$ .*

A useful property of the semantics of the universal quantifier can be obtained now. The easy proof is omitted.

**Proposition 7.2.** *Let  $v$  be a **QmbC**-valuation over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$ , and let  $\varphi$  be a formula such that  $FV(\varphi) \subseteq \{x_1, \dots, x_n\}$ . Then,  $v(\forall x_1 \dots \forall x_n \varphi) \in D$  iff  $v_{\mu}(\varphi) \in D$ , for every  $\mu$ .*

If  $\alpha$  and  $\beta$  are formulas in  $\text{For}(\Theta)$  then  $\alpha \leftrightarrow \beta$  will denote the formula  $(\alpha \rightarrow \beta) \wedge (\beta \rightarrow \alpha)$  in  $\text{For}(\Theta)$ .

**Proposition 7.3.**

- (i)  $\alpha, \alpha \rightarrow \beta \models_{\mathbf{QmbC}} \beta$ ;
- (ii)  $\alpha \rightarrow \beta \models_{\mathbf{QmbC}} \exists x \alpha \rightarrow \beta$ , if  $x \notin FV(\beta)$ ;
- (iii)  $\alpha \rightarrow \beta \models_{\mathbf{QmbC}} \alpha \rightarrow \forall x \beta$ , if  $x \notin FV(\alpha)$ ;
- (iv)  $\models_{\mathbf{QmbC}} \forall x \alpha \rightarrow \alpha[x/t]$ , if  $t$  is a term free for  $x$  in  $\alpha$ ;
- (v)  $\models_{\mathbf{QmbC}} \alpha[x/t] \rightarrow \exists x \alpha$ , if  $t$  is a term free for  $x$  in  $\alpha$ ;
- (vi)  $\models_{\mathbf{QmbC}} \alpha \leftrightarrow \alpha'$ , if  $\alpha$  and  $\alpha'$  are variant.

*Proof.* (i): It is obvious from the definitions.

(ii): Let  $v$  be a **QmbC**-valuation over  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$  such that  $v_{\mu}(\alpha \rightarrow \beta) \in D$ , for every assignment  $\mu$ . Hence  $(v_{\mu}(\alpha))_1 \leq (v_{\mu}(\beta))_1$ , for every  $\mu$ . From this, for every  $a \in U$ :  $(v_{\mu_a^x}(\alpha))_1 \leq (v_{\mu_a^x}(\beta))_1 = (v_{\mu}(\beta))_1$ , since  $x \notin FV(\beta)$ . But then:  $(v_{\mu}(\exists x \alpha))_1 = \bigvee_{a \in U} (v_{\mu}(\alpha[x/a]))_1 = \bigvee_{a \in U} (v_{\mu_a^x}(\alpha))_1 \leq (v_{\mu}(\beta))_1$ . Hence,  $v_{\mu}(\exists x \alpha \rightarrow \beta) \in D$ , for every  $\mu$ . This shows that  $\alpha \rightarrow \beta \models_{\mathbf{QmbC}} \exists x \alpha \rightarrow \beta$ . Item (iii) is proved analogously.

(iv): Assume that  $t$  is a term free for  $x$  in  $\alpha$ . Let  $v$  be a **QmbC**-valuation over  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$  and let  $\mu$  be an assignment. If  $b = \llbracket t \rrbracket_{\mu}^{\mathfrak{A}}$ , then, by Theorem 7.1,  $v_{\mu}(\alpha[x/t]) = v_{\mu}(\alpha[x/\bar{b}])$ . Then,  $(v_{\mu}(\forall x \alpha))_1 = \bigwedge_{a \in U} (v_{\mu}(\alpha[x/\bar{a}]))_1 \leq (v_{\mu}(\alpha[x/\bar{b}]))_1 = (v_{\mu}(\alpha[x/t]))_1$ . Hence,  $v_{\mu}(\forall x \alpha \rightarrow \alpha[x/t]) \in D$ . Item (v) is proved analogously.

(vi): Let  $v$  be a **QmbC**-valuation and let  $\mu$  be an assignment. If  $\alpha$  and  $\alpha'$  are variant, so are  $\hat{\mu}(\alpha)$  and  $\hat{\mu}(\alpha')$ . By Definition 6.9(vii),  $v_{\mu}(\alpha \leftrightarrow \alpha') \in D$ .  $\square$

**Corollary 7.4.** *Let  $v$  be a **QmbC**-valuation over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$ . Then:*

- (1) *If  $\alpha$  is an instance of a **QmbC** axiom schema then  $v_\mu(\alpha) \in D$ , for every assignment  $\mu$ .*
- (2) *If  $\alpha$  and  $\beta$  are formulas such that  $v_\mu(\alpha) \in D$  and  $v_\mu(\alpha \rightarrow \beta) \in D$  for every assignment  $\mu$ , then  $v_\mu(\beta) \in D$  for every assignment  $\mu$ .*
- (3) *If  $\alpha$  and  $\beta$  are formulas such that  $v_\mu(\alpha \rightarrow \beta) \in D$  for every assignment  $\mu$ , and if  $x$  does not occur free in  $\beta$ , then  $v_\mu(\exists x\alpha \rightarrow \beta) \in D$  for every  $\mu$ .*
- (4) *If  $\alpha$  and  $\beta$  are formulas such that  $v_\mu(\alpha \rightarrow \beta) \in D$ , for every  $\mu$ , and if  $x$  does not occur free in  $\alpha$ , then  $v_\mu(\alpha \rightarrow \forall x\beta) \in D$  for every  $\mu$ .*

*Proof.* Item (1) follows by Theorem 2.2.2 in [7] and by Proposition 7.3(iv)-(vi). The rest of the proof follows by Proposition 7.3(i)-(iii).  $\square$

From this corollary it follows easily:

**Theorem 7.5.** (Soundness of **QmbC** w.r.t. first-order swap structures) *For every set  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ : if  $\Gamma \vdash_{\mathbf{QmbC}} \varphi$ , then  $\Gamma \models_{\mathbf{QmbC}} \varphi$ .*

## 8 Completeness of **QmbC** w.r.t. swap structures

In this section the completeness of **QmbC** w.r.t. first-order swap structures semantics for **QmbC** will be obtained. In order to do this, some definitions and results given in Sections 7.5.1 and 7.5.2 of [7] for proving the completeness theorem for **QmbC** w.r.t. interpretations will be adapted. In addition, the technique for proving the completeness of **mbC** w.r.t. swap structures presented in [14, Theorem 7.1] will be also used. The first step is considering a notion of  $C$ -Henkin theory a bit stronger than the one proposed in [7, Definition 7.5.1].

**Definition 8.1.** Consider a theory  $\Delta \subseteq \text{Sen}(\Theta)$  and a nonempty set  $C$  of constants of the signature  $\Theta$ . Then,  $\Delta$  is called a  $C$ -Henkin theory in **QmbC** if it satisfies the following: for every formula  $\varphi$  with (at most) a free variable  $x$ , there exists a constant  $c$  in  $C$  such that  $\Delta \vdash_{\mathbf{QmbC}} \exists x\varphi \rightarrow \varphi[x/c]$ .

**Remark 8.2.** Recall by [7, Section 2.4] that  $\perp_\beta \stackrel{\text{def}}{=} \beta \wedge (\neg\beta \wedge \circ\beta)$  is a bottom in **mbC**, hence  $\sim_\beta\alpha \stackrel{\text{def}}{=} \alpha \rightarrow \perp_\beta$  is a classical negation in **mbC**. This construction does not depend on  $\beta$  (up to logical equivalence), hence we will write  $\sim\alpha$  instead of  $\sim_\beta\alpha$ . This can be also done in **QmbC**. By [7, Proposition 7.2.2] it follows that  $\exists x\sim\varphi \rightarrow \sim\varphi[x/c] \vdash_{\mathbf{QmbC}} \varphi[x/c] \rightarrow \forall x\varphi$ . Thus, if  $\Delta$  is a  $C$ -Henkin theory in **QmbC** and  $\varphi$  is a formula with (at most) a free variable  $x$  then there is a constant  $c$  in  $C$  such that  $\Delta \vdash_{\mathbf{QmbC}} \varphi[x/c] \rightarrow \forall x\varphi$ .

**Definition 8.3.** Let  $\Theta_C$  be the signature obtained from  $\Theta$  by adding a set  $C$  of new individual constants. The consequence relation  $\vdash_{\mathbf{QmbC}}^C$  is the consequence relation of **QmbC** over the signature  $\Theta_C$ .

Recall that, given a Tarskian and finitary logic  $\mathbf{L} = \langle \text{For}, \vdash \rangle$  (where  $\text{For}$  is the set of formulas of  $\mathbf{L}$ ), and given a set  $\Gamma \cup \{\varphi\} \subseteq \text{For}$ , the set  $\Gamma$  is said to be *maximally non-trivial with respect to  $\varphi$  in  $\mathbf{L}$*  if the following holds: (i)  $\Gamma \not\vdash \varphi$ , and (ii)  $\Gamma, \psi \vdash \varphi$  for every  $\psi \notin \Gamma$ .

**Proposition 8.4.** ([7, Corollary 7.5.4]) *Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sen}(\Theta)$  such that  $\Gamma \not\vdash_{\mathbf{QmbC}} \varphi$ . Then, there exists a set of sentences  $\Delta \subseteq \text{Sen}(\Theta)$  which is maximally non-trivial with respect to  $\varphi$  in **QmbC** (by restricting  $\vdash_{\mathbf{QmbC}}$  to sentences) and such that  $\Gamma \subseteq \Delta$ .*

**Definition 8.5.** Let  $\Delta \subseteq \text{Sen}(\Theta)$  be non-trivial in  $\mathbf{QmbC}$ , that is: there is some sentence  $\varphi$  in  $\text{Sen}(\Theta)$  such that  $\Delta \not\vdash_{\mathbf{QmbC}} \varphi$ . Let  $\equiv_{\Delta} \subseteq \text{Sen}(\Theta)^2$  be the relation in  $\text{Sen}(\Theta)$  defined as follows:  $\alpha \equiv_{\Delta} \beta$  iff  $\Delta \vdash_{\mathbf{QmbC}} \alpha \leftrightarrow \beta$ .

By adapting the proof of [14, Theorem 7.1] it follows that  $\equiv_{\Delta}$  is an equivalence relation. Moreover, in the quotient set  $A_{\Delta} \stackrel{\text{def}}{=} \text{Sen}(\Theta)/\equiv_{\Delta}$  it is possible to define binary operators  $\bar{\wedge}, \bar{\vee}, \bar{\rightarrow}$  as follows:  $[\alpha]_{\Delta} \bar{\#} [\beta]_{\Delta} \stackrel{\text{def}}{=} [\alpha \# \beta]_{\Delta}$  for any  $\# \in \{\wedge, \vee, \rightarrow\}$ , where  $[\alpha]_{\Delta}$  denotes the equivalence class of formula  $\alpha$  w.r.t.  $\equiv_{\Delta}$ . Using the axioms of  $\mathbf{QmbC}$  coming from  $\mathbf{mbC}$  it follows:

**Proposition 8.6.** *The structure  $\mathcal{A}_{\Delta} \stackrel{\text{def}}{=} \langle A_{\Delta}, \bar{\wedge}, \bar{\vee}, \bar{\rightarrow}, 0_{\Delta}, 1_{\Delta} \rangle$  is a Boolean algebra with  $0_{\Delta} \stackrel{\text{def}}{=} [\varphi \wedge (\neg\varphi \wedge \circ\varphi)]_{\Delta}$  and  $1_{\Delta} \stackrel{\text{def}}{=} [\varphi \vee \neg\varphi]_{\Delta}$ , for any sentence  $\varphi$ .*

In order to construct the canonical model for  $\mathbf{QmbC}$  w.r.t.  $\Delta$ , the Boolean algebra  $\mathcal{A}_{\Delta}$  needs to be completed. Recall (see, for instance, [23, Chapter 25]) that a Boolean algebra  $\mathcal{B}$  is a *completion* of a Boolean algebra  $\mathcal{A}$  if: (1)  $\mathcal{B}$  is complete, and (2)  $\mathcal{B}$  includes  $\mathcal{A}$  as a dense subalgebra (that is: every element in  $\mathcal{B}$  is the supremum, in  $\mathcal{B}$ , of some subset of  $\mathcal{A}$ ). As a consequence of the definition, it follows that  $\mathcal{B}$  preserves all the existing infima and suprema in  $\mathcal{A}$ . In formal terms: there exists a monomorphism of Boolean algebras (therefore an injective mapping)  $*$  :  $\mathcal{A} \rightarrow \mathcal{B}$  such that  $*(\bigvee_{\mathcal{A}} X) = \bigvee_{\mathcal{B}} *[X]$  for every  $X \subseteq \mathcal{A}$  such that the supremum  $\bigvee_{\mathcal{A}} X$  exists, where  $*[X] = \{*(a) : a \in X\}$ . Analogously,  $*(\bigwedge_{\mathcal{A}} X) = \bigwedge_{\mathcal{B}} *[X]$  for every  $X \subseteq \mathcal{A}$  such that the infimum  $\bigwedge_{\mathcal{A}} X$  exists. By the (independent) results of MacNeille and Tarski, it is known that every Boolean algebra has a completion; moreover, the completion is unique up to isomorphisms. Thus, let  $C\mathcal{A}_{\Delta}$  be the completion of  $\mathcal{A}_{\Delta}$  and let  $*$  :  $\mathcal{A}_{\Delta} \rightarrow C\mathcal{A}_{\Delta}$  be the associated monomorphism.

**Definition 8.7.** Let  $C\mathcal{A}_{\Delta}$  be the complete Boolean algebra defined as above. The full swap structure for  $\mathbf{mbC}$  over  $C\mathcal{A}_{\Delta}$  (recall Definition 3.5) will be denoted by  $\mathcal{B}_{\Delta}$ . The associated Nmatrix (recall Definition 3.6) will be denoted by  $\mathcal{M}(\mathcal{B}_{\Delta}) \stackrel{\text{def}}{=} (\mathcal{B}_{\Delta}, D_{\Delta})$ .

Notice that  $(*([\alpha]_{\Delta}), *([\beta]_{\Delta}), *([\gamma]_{\Delta})) \in D_{\Delta}$  iff  $\Delta \vdash_{\mathbf{QmbC}} \alpha$ .

**Definition 8.8.** (Canonical Structure) Let  $\Theta$  be a signature with some individual constant. Let  $\Delta \subseteq \text{Sen}(\Theta)$  be non-trivial in  $\mathbf{QmbC}$ , let  $\mathcal{M}(\mathcal{B}_{\Delta})$  be as in Definition 8.7, and let  $U = C\text{Ter}(\Theta)$ . The *canonical structure induced by  $\Delta$*  is the structure  $\mathfrak{A}_{\Delta} = \langle U, I_{\mathfrak{A}_{\Delta}} \rangle$  over  $\mathcal{M}(\mathcal{B}_{\Delta})$  and  $\Theta$  such that:

- $c^{\mathfrak{A}_{\Delta}} = c$ , for each individual constant  $c$ ;
- $f^{\mathfrak{A}_{\Delta}} : U^n \rightarrow U$  is such that  $f^{\mathfrak{A}_{\Delta}}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ , for each function symbol  $f$  of arity  $n$ ;
- $P^{\mathfrak{A}_{\Delta}}(t_1, \dots, t_n) = (*([\varphi]_{\Delta}), *([\neg\varphi]_{\Delta}), *([\circ\varphi]_{\Delta}))$  with  $\varphi = P(t_1, \dots, t_n)$ , for each predicate symbol  $P$  of arity  $n$ .

Notice that  $[\varphi]_{\Delta} \bar{\vee} [\neg\varphi]_{\Delta} = [\varphi \vee \neg\varphi]_{\Delta} = 1$  and  $[\varphi]_{\Delta} \bar{\wedge} [\neg\varphi]_{\Delta} \bar{\wedge} [\circ\varphi]_{\Delta} = [\varphi \wedge \neg\varphi \wedge \circ\varphi]_{\Delta} = 0$ . Thus  $*([\varphi]_{\Delta}) \vee *([\neg\varphi]_{\Delta}) = *([\varphi]_{\Delta} \bar{\vee} [\neg\varphi]_{\Delta}) = 1$ , and  $*([\varphi]_{\Delta}) \wedge *([\neg\varphi]_{\Delta}) \wedge *([\circ\varphi]_{\Delta}) = *([\varphi]_{\Delta} \bar{\wedge} [\neg\varphi]_{\Delta} \bar{\wedge} [\circ\varphi]_{\Delta}) = 0$ . Hence  $P^{\mathfrak{A}_{\Delta}}(t_1, \dots, t_n) \in |\mathcal{B}_{\Delta}|$  and so  $\mathfrak{A}_{\Delta}$  is indeed a structure over  $\mathcal{M}(\mathcal{B}_{\Delta})$  and  $\Theta$ .

**Definition 8.9.** Let  $(\cdot)^{\triangleright} : (\text{Ter}(\Theta_U) \cup \text{For}(\Theta_U)) \rightarrow (\text{Ter}(\Theta) \cup \text{For}(\Theta))$  be the mapping such that  $(s)^{\triangleright}$  is the expression obtained from  $s$  by substituting every occurrence of a constant  $\bar{t}$  by the term  $t$  itself, for  $t \in C\text{Ter}(\Theta)$ .

For instance,  $\left( P(f(\bar{c}, x)) \wedge Q(\overline{f(c, x)}, z) \right)^\triangleright = P(f(c, x)) \wedge Q(f(c, x), z)$ .

**Definition 8.10.** (Canonical valuation) Let  $\Delta \subseteq \text{Sen}(\Theta)$  be a set of sentences over a signature  $\Theta$  such that  $\Delta$  is a  $C$ -Henkin theory in  $\mathbf{QmbC}$  for a nonempty set  $C$  of individual constants of  $\Theta$ , and  $\Delta$  is maximally non-trivial with respect to  $\varphi$  in  $\mathbf{QmbC}$ , for some sentence  $\varphi$ . The *canonical  $\mathbf{QmbC}$ -valuation induced by  $\Delta$  over  $\mathfrak{A}_\Delta$  and  $\mathcal{M}(\mathcal{B}_\Delta)$*  is the function  $v_\Delta : \text{Sen}(\Theta_U) \rightarrow |\mathcal{B}_\Delta|$  such that  $v_\Delta(\psi) = (*([\psi]^\triangleright)_\Delta, *([\neg(\psi)]^\triangleright)_\Delta, *([\circ(\psi)]^\triangleright)_\Delta)$ , for every sentence  $\psi$  over  $\Theta_U$ .

Notice that  $v_\Delta(\psi) \in D_\Delta$  iff  $\Delta \vdash_{\mathbf{QmbC}} (\psi)^\triangleright$ .

**Lemma 8.11.** Let  $\Delta \subseteq \text{Sen}(\Theta)$  be as in Definition 8.10. Then, for every formula  $\psi(x)$  in which  $x$  is the unique variable (possibly) occurring free, it holds:

- (1)  $[\forall x\psi]_\Delta = \bigwedge_{\mathcal{A}_\Delta} \{[\psi[x/t]]_\Delta : t \in C\text{Ter}(\Theta)\}$ , where  $\bigwedge_{\mathcal{A}_\Delta}$  denotes an existing infimum in the Boolean algebra  $\mathcal{A}_\Delta$ ;
- (2)  $[\exists x\psi]_\Delta = \bigvee_{\mathcal{A}_\Delta} \{[\psi[x/t]]_\Delta : t \in C\text{Ter}(\Theta)\}$ , where  $\bigvee_{\mathcal{A}_\Delta}$  denotes an existing supremum in the Boolean algebra  $\mathcal{A}_\Delta$ .

*Proof.*

(2) Observe that, in  $\mathcal{A}_\Delta$ ,  $[\alpha]_\Delta \leq [\beta]_\Delta$  iff  $\Delta \vdash_{\mathbf{QmbC}} \alpha \rightarrow \beta$ . Let  $\alpha$  be a formula in which  $x$  is the unique variable (possibly) occurring free. Then  $[\alpha[x/t]]_\Delta \leq [\exists x\alpha]_\Delta$  for every  $t \in C\text{Ter}(\Theta)$ , by **(Ax12)**. Let  $\beta$  be a sentence such that  $[\alpha[x/t]]_\Delta \leq [\beta]_\Delta$  for every  $t \in C\text{Ter}(\Theta)$ . That is,  $\Delta \vdash_{\mathbf{QmbC}} \alpha[x/t] \rightarrow \beta$  for every  $t \in C\text{Ter}(\Theta)$ . Since  $\Delta$  is a  $C$ -Henkin theory, there is a constant  $c$  of  $\Theta$  such that  $\Delta \vdash_{\mathbf{QmbC}} \exists x\alpha \rightarrow \alpha[x/c]$ . By hypothesis,  $\Delta \vdash_{\mathbf{QmbC}} \alpha[x/c] \rightarrow \beta$  and so  $\Delta \vdash_{\mathbf{QmbC}} \exists x\alpha \rightarrow \beta$ . That is,  $[\exists x\alpha]_\Delta \leq [\beta]_\Delta$ . This shows that  $[\exists x\alpha]_\Delta = \bigvee_{\mathcal{A}_\Delta} \{[\alpha[x/t]]_\Delta : t \in C\text{Ter}(\Theta)\}$ .

Item (1) is proved analogously, but now by using Remark 8.2. □

**Theorem 8.12.** The canonical  $\mathbf{QmbC}$ -valuation  $v_\Delta$  is a  $\mathbf{QmbC}$ -valuation over  $\mathfrak{A}_\Delta$  and  $\mathcal{M}(\mathcal{B}_\Delta)$ .

*Proof.* Let us see that  $v_\Delta$  satisfies all the requirements of Definition 6.9.

(i) If  $\varphi$  is an atomic formula  $P(t_1, \dots, t_n)$  then:

$$v_\Delta(\varphi) = (*([\varphi]^\triangleright)_\Delta, *([\neg(\varphi)]^\triangleright)_\Delta, *([\circ(\varphi)]^\triangleright)_\Delta) = P^{\mathfrak{A}_\Delta}((t_1)^\triangleright, \dots, (t_n)^\triangleright) = P^{\mathfrak{A}_\Delta}([\![t_1]\!]^{\mathfrak{A}_\Delta}, \dots, [\![t_n]\!]^{\mathfrak{A}_\Delta}).$$

(ii)  $v_\Delta(\neg\psi) = (*([\neg(\psi)]^\triangleright)_\Delta, *([\neg\neg(\psi)]^\triangleright)_\Delta, *([\circ\neg(\psi)]^\triangleright)_\Delta) \in \tilde{\sim}v_\Delta(\psi)$ . On the other hand,  $v_\Delta(\circ\psi) = (*([\circ(\psi)]^\triangleright)_\Delta, *([\neg\circ(\psi)]^\triangleright)_\Delta, *([\circ\circ(\psi)]^\triangleright)_\Delta) \in \tilde{\circ}v_\Delta(\psi)$ .

(iii) Since  $*([\delta\#\psi]_\Delta) = *([\delta]_\Delta) \# *([\psi]_\Delta)$ , then  $v_\Delta(\delta\#\psi) \in v(\delta)\tilde{\#}v_\Delta(\psi)$ , for every  $\# \in \{\wedge, \vee, \rightarrow\}$ .

(iv) By Lemma 8.11 (and recalling that  $U = C\text{Ter}(\Theta)$ ),  $[\forall x\psi]_\Delta = \bigwedge_{\mathcal{A}_\Delta} \{[\psi[x/t]]_\Delta : t \in U\}$  and so  $*([\forall x\psi]_\Delta) = \bigwedge_{C\mathcal{A}_\Delta} \{*([\psi[x/t]]_\Delta) : t \in U\}$ . Then,  $(v_\Delta(\forall x\psi))_1 = *([\forall x\psi]^\triangleright)_\Delta = \bigwedge_{t \in U} *([\psi[x/\bar{t}]]^\triangleright)_\Delta = \bigwedge_{t \in U} (v_\Delta(\psi[x/\bar{t}]))_1$ .

(v) The case  $\exists x\psi$  is treated analogously.

(vi) Let  $t$  be free for  $z$  in  $\varphi$ ,  $\mu$  an assignment and  $b = [\![t]\!]^{\mathfrak{A}_\Delta}$ . By induction on the complexity of  $\varphi$  it can be proved that  $(\widehat{\mu}(\varphi[z/t]))^\triangleright = (\widehat{\mu}(\varphi[z/\bar{b}]))^\triangleright$ . Hence,  $v_\Delta(\widehat{\mu}(\varphi[z/t])) = v_\Delta(\widehat{\mu}(\varphi[z/\bar{b}]))$ , by definition of  $v_\Delta$ . From this, it is obvious that  $v_\Delta$  satisfies conditions (vi.1)-(vi.3).

(vii) If  $\varphi$  and  $\varphi'$  are variant, so are  $(\varphi)^\triangleright$  and  $(\varphi')^\triangleright$ ;  $(\neg\varphi)^\triangleright$  and  $(\neg\varphi')^\triangleright$ ; and  $(\circ\varphi)^\triangleright$  and  $(\circ\varphi')^\triangleright$ . From this,  $v_\Delta(\varphi) = v_\Delta(\varphi')$ , by axiom **(Ax14)**. □

**Theorem 8.13.** (Completeness of **QmbC** restricted to sentences w.r.t. first-order swap structures) *Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sen}(\Theta)$ . If  $\Gamma \models_{\mathbf{QmbC}} \varphi$  then  $\Gamma \vdash_{\mathbf{QmbC}} \varphi$ .*

*Proof.* Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sen}(\Theta)$  such that  $\Gamma \not\vdash_{\mathbf{QmbC}} \varphi$ . Then, by recalling Definition 8.1 and by Theorem 7.5.3 in [7],<sup>6</sup> there exists a  $C$ -Henkin theory  $\Delta^H$  over  $\Theta_C$  in **QmbC** for a nonempty set  $C$  of new individual constants such that  $\Gamma \subseteq \Delta^H$  and, for every  $\alpha \in \text{Sen}(\Theta)$ :  $\Gamma \vdash_{\mathbf{QmbC}} \alpha$  iff  $\Delta^H \vdash_{\mathbf{QmbC}}^C \alpha$ . Hence,  $\Delta^H \not\vdash_{\mathbf{QmbC}}^C \varphi$ . Now, by Proposition 8.4, there exists a set of sentences  $\overline{\Delta^H}$  in  $\Theta_C$  extending  $\Delta^H$  which is maximally non-trivial with respect to  $\varphi$  in **QmbC** (defined over  $\text{Sen}(\Theta_C)$ ), such that  $\overline{\Delta^H}$  is a  $C$ -Henkin theory over  $\Theta_C$  in **QmbC**. Let  $\mathcal{M}(\mathcal{B}_{\overline{\Delta^H}})$ ,  $\mathfrak{A}_{\overline{\Delta^H}}$  and  $v_{\overline{\Delta^H}}$  be as in Definitions 8.7, 8.8 and 8.10, respectively. Then,  $v_{\overline{\Delta^H}}(\alpha) \in D_{\overline{\Delta^H}}$  iff  $\overline{\Delta^H} \vdash_{\mathbf{QmbC}}^C \alpha$ , for every  $\alpha$  in  $\text{Sen}(\Theta_C)$ . From this,  $v_{\overline{\Delta^H}}[\Gamma] \subseteq D_{\overline{\Delta^H}}$  and  $v_{\overline{\Delta^H}}(\varphi) \notin D_{\overline{\Delta^H}}$ . Finally, let  $\mathfrak{A}$  and  $v$  be the respective restrictions of  $\mathfrak{A}_{\overline{\Delta^H}}$  and  $v_{\overline{\Delta^H}}$  to  $\Theta$ . Then,  $\mathfrak{A}$  is a structure over  $\mathcal{M}(\mathcal{B}_{\overline{\Delta^H}})$ , and  $v$  is a valuation for **QmbC** over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B}_{\overline{\Delta^H}})$  such that  $v[\Gamma] \subseteq D_{\overline{\Delta^H}}$  but  $v(\varphi) \notin D_{\overline{\Delta^H}}$ . This shows that  $\Gamma \not\models_{\mathbf{QmbC}} \varphi$ .  $\square$

For any formula  $\psi$  in  $\text{For}(\Theta)$  let  $(\forall)\psi$  be the *universal closure* of  $\psi$ , defined as follows: if  $\psi$  is a sentence then  $(\forall)\psi \stackrel{\text{def}}{=} \psi$ ; and if  $\psi$  has exactly the variables  $x_1, \dots, x_n$  occurring free then  $(\forall)\psi \stackrel{\text{def}}{=} (\forall x_1) \cdots (\forall x_n)\psi$ . Note that  $(\forall)\psi \in \text{Sen}(\Theta)$ . If  $\Gamma$  is a set of formulas in  $\text{For}(\Theta)$  then  $(\forall)\Gamma \stackrel{\text{def}}{=} \{(\forall)\psi : \psi \in \Gamma\}$ . It is easy to show that, for every  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ : (i)  $\Gamma \vdash_{\mathbf{QmbC}} \varphi$  iff  $(\forall)\Gamma \vdash_{\mathbf{QmbC}} (\forall)\varphi$ ; and (ii)  $\Gamma \models_{\mathbf{QmbC}} \varphi$  iff  $(\forall)\Gamma \models_{\mathbf{QmbC}} (\forall)\varphi$ . From this, a general completeness result can be obtained:

**Corollary 8.14.** (Completeness of **QmbC** w.r.t. first-order swap structures) *Let  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ . If  $\Gamma \models_{\mathbf{QmbC}} \varphi$  then  $\Gamma \vdash_{\mathbf{QmbC}} \varphi$ .*

## 9 Completeness of **QmbC** w.r.t. structures over $\mathcal{M}_5$

Recall from Section 4 the 5-valued Nmatrix  $\mathcal{M}_5$  introduced by Avron in [1]. From the adequacy of **QmbC** w.r.t. first-order swap structures, and given that **mbC** can be characterized just with  $\mathcal{M}_5 = \mathcal{M}(\mathcal{B}_{A_2})$ , it is a natural question to determine if it is possible to extend the proof of [7, Theorem 6.4.9 and Corollary 6.4.10] (see Theorem 4.4 above) to **QmbC**. Namely, taking into account that **QmbC** can be characterized by standard Tarskian structures expanded with bivaluations which naturally extend the ones for **mbC** (see Theorem 9.3 below), it seems plausible to extend the technique of Theorem 4.4 to **QmbC**. In Theorem 9.6 below it will be shown that this is really the case, hence **QmbC** can be characterized by first-order structures over  $\mathcal{M}_5$ . Such structures, which were introduced by Avron and Zamansky in [6] (see Remark 9.4 below), can be considered as being ‘classical’, as discussed at the end of Section 4.

Consider a standard Tarskian first-order structure  $\mathbf{A} = \langle U, I_{\mathbf{A}} \rangle$  over a first-order signature  $\Theta$  (see, for instance, [26]). Observe that  $\mathbf{A}$  is defined as in Definition 6.1, but now any predicate symbol  $P$  of arity  $n$  is interpreted as a subset  $I_{\mathbf{A}}(P)$  of  $U^n$ . The notions of diagram language  $\text{For}(\Theta_U)$ , extended structure  $\widehat{\mathbf{A}}$  and  $\text{Sen}(\Theta_U)$  are defined as in Definitions 6.4 and 6.5, and Notation 6.6 above.

In [7] the notion of bivaluations for **mbC** was extended to *bivaluations for **QmbC*** as follows:

<sup>6</sup>The proof of Theorem 7.5.3 in [7] also holds for the notion of  $C$ -Henkin theory adopted here in Definition 8.1 (which is stronger than the one proposed in [7, Definition 7.5.1]), as it can be easily verified.

**Definition 9.1.** (Bivaluations for **QmbC**, [7, Definition 7.3.5]) Let  $\mathbf{A} = \langle U, I_{\mathbf{A}} \rangle$  be a standard Tarskian first-order structure over  $\Theta$ , and let  $\widehat{\mathbf{A}} = \langle U, I_{\widehat{\mathbf{A}}} \rangle$  be the expansion of  $\mathbf{A}$  to  $\Theta_U$  by setting  $I_{\widehat{\mathbf{A}}}(\bar{a}) = a$  for every  $a \in U$ . A *bivaluation*<sup>7</sup> for **QmbC** over  $\mathbf{A}$  is a function  $\rho : \text{Sen}(\Theta_U) \rightarrow \{0, 1\}$  satisfying the clauses of Definition 4.2 above plus the following:

(**vPred**)  $\rho(P(t_1, \dots, t_n)) = 1$  iff  $\langle \llbracket t_1 \rrbracket^{\widehat{\mathbf{A}}}, \dots, \llbracket t_n \rrbracket^{\widehat{\mathbf{A}}} \rangle \in I_{\mathbf{A}}(P)$ , for  $P(t_1, \dots, t_n)$  atomic

(**vVar**)  $\rho(\varphi) = \rho(\psi)$  whenever  $\varphi$  is a variant of  $\psi$

(**vUni**)  $\rho(\forall x\varphi) = 1$  iff  $\rho(\varphi[x/\bar{a}]) = 1$  for every  $a \in U$

(**vEx**)  $\rho(\exists x\varphi) = 1$  iff  $\rho(\varphi[x/\bar{a}]) = 1$  for some  $a \in U$

(**vSubs**) if  $\mu$  is an assignment,  $t$  is a term free for  $z$  in  $\varphi$  and  $b = \llbracket t \rrbracket^{\widehat{\mathbf{A}}}_{\mu}$ , then:  $\rho(\widehat{\mu}(\varphi[z/t])) = \rho(\widehat{\mu}(\varphi[z/\bar{b}]))$  implies  $\rho(\widehat{\mu}(\#\varphi[z/t])) = \rho(\widehat{\mu}(\#\varphi[z/\bar{b}]))$ , for  $\# \in \{\neg, \circ\}$ .

**Definition 9.2.** ([7, Definitions 7.3.6 and 7.3.12]) An *interpretation* for **QmbC** over a signature  $\Theta$  is a pair  $\langle \mathbf{A}, \rho \rangle$  such that  $\mathbf{A}$  is a Tarskian first-order structure over  $\Theta$  and  $\rho$  is a bivaluation for **QmbC** over  $\mathbf{A}$ . The *consequence relation*  $\models_{\mathbf{QmbC}}^2$  of **QmbC** w.r.t. interpretations is given by:  $\Gamma \models_{\mathbf{QmbC}}^2 \varphi$  if, for every interpretation  $\langle \mathbf{A}, \rho \rangle$ :  $\rho(\widehat{\mu}(\gamma)) = 1$  for every  $\gamma \in \Gamma$  and every assignment  $\mu$  implies that  $\rho(\widehat{\mu}(\varphi)) = 1$  for every assignment  $\mu$ .

**Theorem 9.3.** (Adequacy of **QmbC** w.r.t. interpretations, [7, Theorems 7.4.1. and 7.5.6]) If  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$  then:  $\Gamma \vdash_{\mathbf{QmbC}} \varphi$  iff  $\Gamma \models_{\mathbf{QmbC}}^2 \varphi$ .<sup>8</sup>

Now, Theorem 4.4 will be extended to **QmbC** (see Theorem 9.5 below). Previous to this, it is worth observing the following:

**Remark 9.4.** Consider once again the characteristic 5-valued Nmatrix  $\mathcal{M}_5 = \mathcal{M}(\mathcal{B}_{\mathcal{A}_2})$  for **mbC**, and let  $\mathfrak{A}$  be a structure over  $\Theta$  and  $\mathcal{M}_5$ . If  $v$  is a valuation for **QmbC** over  $\mathfrak{A}$  and  $\mathcal{M}_5$ , it is easy to see that clauses (iv) and (v) of Definition 6.9 are equivalent to the following: for every  $Q \in \{\forall, \exists\}$ ,  $v(Qx\varphi) \in \tilde{Q}(\{v(\varphi[x/\bar{a}]) : a \in U\})$  where  $\tilde{Q} : (\mathcal{P}(B_{\mathcal{A}_2}) - \{\emptyset\}) \rightarrow (\mathcal{P}(B_{\mathcal{A}_2}) - \{\emptyset\})$  is

$$\tilde{\forall}(X) = \begin{cases} \text{D}, & \text{if } X \subseteq \text{D} \\ \text{ND}, & \text{otherwise} \end{cases} \quad \text{and} \quad \tilde{\exists}(X) = \begin{cases} \text{D}, & \text{if } X \cap \text{D} \neq \emptyset \\ \text{ND}, & \text{otherwise} \end{cases}$$

and  $B_{\mathcal{A}_2} = |\mathcal{M}_5| = \{T, t, t_0, F, f_0\}$ ,  $\text{D} = \{T, t, t_0\}$  and  $\text{ND} = \{F, f_0\}$  (recall Section 4). It is not hard to prove that the notions of structures over  $\Theta$  and  $\mathcal{M}_5$ , and valuations over them, coincide with the corresponding notions introduced in [6]. Thus, the present framework generalizes, from  $\mathcal{A}_2$  to arbitrary complete Boolean algebras, the semantical framework proposed in [6].

**Theorem 9.5.** Let  $\mathcal{I} = \langle \mathbf{A}, \rho \rangle$  be an interpretation for **QmbC** over a signature  $\Theta$ . Then, it induces a first-order structure  $\mathfrak{A}_{\mathcal{I}}$  over  $\mathcal{M}_5$  and  $\Theta$ , and a **QmbC**-valuation  $v^\rho$  over  $\mathfrak{A}_{\mathcal{I}}$  and  $\mathcal{M}_5$  given by  $v^\rho(\alpha) \stackrel{\text{def}}{=} (\rho(\alpha), \rho(\neg\alpha), \rho(\circ\alpha))$  such that:  $\rho(\alpha) = 1$  iff  $v^\rho(\alpha) \in \text{D}$ , for every sentence  $\alpha \in \text{Sen}(\Theta_U)$ .

<sup>7</sup>It was called **QmbC**-valuation in [7, Definition 7.3.5].

<sup>8</sup>Rigourously speaking, in [7, Theorem 7.5.6] it was obtained completeness of **QmbC** w.r.t. interpretations, but only for sentences. However, completeness of **QmbC** (in the full language) w.r.t. interpretations follows easily, as we have done here in Corollary 8.14.



*Proof.* Given  $\mathcal{I} = \langle \mathbf{A}, \rho \rangle$  consider the first-order structure  $\mathfrak{A}_{\mathcal{I}}$  over  $\mathcal{M}_5$  and  $\Theta$  obtained from  $\mathbf{A}$  by taking the same domain  $U$ ;  $I_{\mathfrak{A}_{\mathcal{I}}}$  coincides with  $I_{\mathbf{A}}$  for every individual constant and function symbol; and  $I_{\mathfrak{A}_{\mathcal{I}}}(P) : U^n \rightarrow |\mathcal{M}_5|$  is given by  $I_{\mathfrak{A}_{\mathcal{I}}}(P)(a_1, \dots, a_n) = v^\rho(P(\bar{a}_1, \dots, \bar{a}_n))$  for every predicate symbol  $P$  of arity  $n$ , where  $v^\rho : \text{Sen}(\Theta_U) \rightarrow |\mathcal{M}_5|$  is defined by  $v^\rho(\alpha) = (\rho(\alpha), \rho(\neg\alpha), \rho(\circ\alpha))$ , for every  $\alpha \in \text{Sen}(\Theta_U)$ . Clearly  $\rho(\alpha) = 1$  iff  $v^\rho(\alpha) \in \text{D}$ , for every  $\alpha \in \text{Sen}(\Theta_U)$ . Thus, it remains to prove that  $v^\rho$  is indeed a **QmbC**-valuation over  $\mathfrak{A}_{\mathcal{I}}$  and  $\mathcal{M}_5$ . It is clear that clauses (i)-(iii) of Definition 6.9 are satisfied, since  $\llbracket t \rrbracket^{\mathfrak{A}_{\mathcal{I}}} = \llbracket t \rrbracket^{\hat{\mathbf{A}}}$  for every closed term  $t$ , and by Theorem 4.4. With respect to clause (iv), suppose that  $\{v^\rho(\varphi[x/\bar{a}]) : a \in U\} \subseteq \text{D}$ . Then  $\rho(\varphi[x/\bar{a}]) = 1$  for every  $a \in U$  and so  $\rho(\forall x\varphi) = 1$ , by (*vUni*). Hence  $v^\rho(\forall x\varphi) \in \text{D}$ . This means that  $v^\rho(\forall x\varphi) \in \tilde{\text{V}}(\{v^\rho(\varphi[x/\bar{a}]) : a \in U\})$ . If  $\{v^\rho(\varphi[x/\bar{a}]) : a \in U\} \not\subseteq \text{D}$  then, by a similar reasoning, it is shown that, once again,  $v^\rho(\forall x\varphi) \in \tilde{\text{V}}(\{v^\rho(\varphi[x/\bar{a}]) : a \in U\})$ . Analogously it can be proven that  $v^\rho$  satisfies clause (v). Clause (vi) is satisfied by  $v^\rho$ , since  $\llbracket t \rrbracket_\mu^{\mathfrak{A}_{\mathcal{I}}} = \llbracket t \rrbracket_\mu^{\hat{\mathbf{A}}}$  for every term  $t$ , and by the fact that  $\rho$  satisfies the Substitution Lemma:  $\rho(\hat{\mu}(\varphi[z/t])) = \rho(\hat{\mu}(\varphi[z/\bar{b}]))$  for  $b = \llbracket t \rrbracket_\mu^{\hat{\mathbf{A}}}$ . Clause (vii) is also satisfied, since  $\rho$  satisfies (*vVar*). This concludes the proof.  $\square$

**Theorem 9.6.** (Adequacy of **QmbC** w.r.t. first-order structures over  $\mathcal{M}_5$ ) *For every set  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ :  $\Gamma \vdash_{\text{QmbC}} \varphi$  iff  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}_5)} \varphi$  for every structure  $\mathfrak{A}$  over  $\Theta$  and  $\mathcal{M}_5$ .*

*Proof.*

‘Only if’ part (Soundness): It is a consequence of Theorem 7.5.

‘If’ part (Completeness): Suppose that  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}_5)} \varphi$  for every structure  $\mathfrak{A}$  over  $\Theta$  and  $\mathcal{M}_5$ . Let  $\mathcal{I} = \langle \mathbf{A}, \rho \rangle$  be an interpretation for **QmbC** over  $\Theta$  such that  $\rho(\hat{\mu}(\gamma)) = 1$  for every  $\gamma \in \Gamma$  and every assignment  $\mu$ . Let  $\mathfrak{A}_{\mathcal{I}}$  and  $v^\rho$  be as in Theorem 9.5. Then  $v^\rho_\mu(\gamma) \in \text{D}$ , for every formula  $\gamma \in \Gamma$  and every assignment  $\mu$ . By hypothesis,  $v^\rho_\mu(\varphi) \in \text{D}$ , for every assignment  $\mu$ . This implies that  $\rho(\hat{\mu}(\varphi)) = 1$ , for every assignment  $\mu$ . That is,  $\Gamma \models_{\text{QmbC}}^2 \varphi$ . Therefore  $\Gamma \vdash_{\text{QmbC}} \varphi$ , by Theorem 9.3.  $\square$

Taking into account Remark 9.4, the last result is a restatement of the adequacy for **QmbC** w.r.t. first-order structures over  $\mathcal{M}_5$  obtained in [6, Theorem 24].

## 10 Adding standard equality to **QmbC**

In this section a binary predicate  $\approx$  for dealing with equality will be considered. As expected, this predicate will be always interpreted as the standard identity. This means that the predicate  $\approx$  will be seen, from a semantical point of view, as a logical symbol. The resulting logic will be called **QmbC** $^\approx$ . The definition of **QmbC** $^\approx$  will follow closely [7, Section 7.7].

**Definition 10.1.** Let  $\Theta$  be a first-order signature. The induced signature with equality  $\Theta_\approx$  is obtained from  $\Theta$  by adding a new binary predicate symbol  $\approx$ .

The expression  $(t_1 \approx t_2)$  will stand for the atomic formula  $\approx(t_1, t_2)$ . If  $\varphi$  is a formula and  $y$  is a variable free for the variable  $x$  in  $\varphi$ ,  $\varphi[x \wr y]$  denotes any formula obtained from  $\varphi$  by replacing some, but not necessarily all (maybe none), free occurrences of  $x$  by  $y$ .

**Definition 10.2.** ([7, Definition 7.7.1]) Let  $\Theta_\approx$  be a first-order signature with equality. The logic **QmbC** $^\approx$  (over  $\Theta_\approx$ ) is the extension of **QmbC** over  $\text{For}(\Theta_\approx)$  obtained by

adding to  $\mathbf{QmbC}$ , besides all the new instances of axioms and inference rules involving the equality predicate  $\approx$ , the following axiom schemas:

$$(\mathbf{AxEq1}) \quad \forall x(x \approx x)$$

$$(\mathbf{AxEq2}) \quad (x \approx y) \rightarrow (\varphi \rightarrow \varphi[x \wr y]), \text{ if } y \text{ is a variable free for } x \text{ in } \varphi$$

Notice that the axioms for equality are the same considered for classical logic (see, for instance, [26]). Given that  $\mathbf{QmbC}^\approx$  is an axiomatic extension of  $\mathbf{QmbC}$ , it satisfies the deduction meta-theorem DMT (recall Theorem 5.6). Let  $\vdash_{\mathbf{QmbC}^\approx}$  be the consequence relation of the Hilbert calculus  $\mathbf{QmbC}^\approx$ . The semantics of first-order swap structures for  $\mathbf{QmbC}$  can be easily extended to the equality predicate.

**Definition 10.3.** Let  $\mathcal{M}(\mathcal{B}) = (\mathcal{B}, D)$  be a non-deterministic matrix defined by a swap structure  $\mathcal{B}$  for  $\mathbf{mbC}$ , and let  $\Theta_\approx$  be a first-order signature with equality. A (first-order) *structure with standard equality* over  $\mathcal{M}(\mathcal{B})$  and  $\Theta_\approx$  is a structure over  $\mathcal{M}(\mathcal{B})$  and  $\Theta_\approx$  such that  $I_{\mathfrak{A}}(\approx)(a, b) \in D$  iff  $a = b$ .

In what follows,  $(a \approx^{\mathfrak{A}} b)$  will stand for  $I_{\mathfrak{A}}(\approx)(a, b)$ , for every structure  $\mathfrak{A}$  and any  $a, b \in U$ . Given a structure  $\mathfrak{A}$ , the signature obtained from  $\Theta$  by adding a new individual constant for each element of  $U$  (recall Definition 6.4) will be denoted by  $\Theta_{\tilde{U}}$ . The set of formulas and sentences over  $\Theta_{\tilde{U}}$  will be denoted by  $For(\Theta_{\tilde{U}})$  and  $Sen(\Theta_{\tilde{U}})$ , respectively.

If  $\mathfrak{A}$  is a structure with standard equality over  $\mathcal{M}(\mathcal{B})$  and  $v$  is a  $\mathbf{QmbC}$ -valuation over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$  then, by Definition 6.9 (i) it follows that, for every closed terms  $t_1$  and  $t_2$ ,  $v(t_1 \approx t_2) \in D$  iff  $\llbracket t_1 \rrbracket^{\mathfrak{A}} = \llbracket t_2 \rrbracket^{\mathfrak{A}}$ . This guarantees the validity of axiom  $(\mathbf{AxEq1})$ . However, in order to validate axiom  $(\mathbf{AxEq2})$ , the valuations must be additionally restricted:

**Definition 10.4.** ( $\mathbf{QmbC}^\approx$ -valuations) Let  $\mathfrak{A}$  be a structure with standard equality over  $\Theta_\approx$  and  $\mathcal{M}(\mathcal{B})$ . A *valuation* for  $\mathbf{QmbC}^\approx$  (or a  $\mathbf{QmbC}^\approx$ -valuation) over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B})$  is a  $\mathbf{QmbC}$ -valuation  $v : Sen(\Theta_{\tilde{U}}) \rightarrow |\mathcal{B}|$  which satisfies, in addition, the following clause, for every  $\mu$ :

$$(viii) \quad v_\mu((x \approx y) \rightarrow (\varphi \rightarrow \varphi[x \wr y])) \in D, \text{ if } y \text{ is a variable free for } x \text{ in } \varphi.$$

The notion of  $\varphi$  being a  $\approx$ -semantical consequence of  $\Gamma$  over  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$ , denoted by  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}(\mathcal{B}))}^\approx \varphi$ , is as stated in Definition 6.11, but now restricted to structures with standard equality and  $\mathbf{QmbC}^\approx$ -valuations over them. Thus,  $\varphi$  is a *semantical consequence of  $\Gamma$  in  $\mathbf{QmbC}^\approx$  w.r.t. first-order swap structures*, denoted by  $\Gamma \models_{\mathbf{QmbC}^\approx} \varphi$ , if  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}(\mathcal{B}))}^\approx \varphi$  for every of such pairs  $(\mathfrak{A}, \mathcal{M}(\mathcal{B}))$ .

Observe that the Substitution Lemma still holds for  $\mathbf{QmbC}^\approx$ , since it holds for any structure and any  $\mathbf{QmbC}$ -valuation. From this, and from Definition 10.4, the following result can be easily derived by adapting the proof of Theorem 7.5:

**Theorem 10.5.** (Soundness of  $\mathbf{QmbC}^\approx$  w.r.t. first-order swap structures with standard equality) *For every set  $\Gamma \cup \{\varphi\} \subseteq For(\Theta_\approx)$ : if  $\Gamma \vdash_{\mathbf{QmbC}^\approx} \varphi$ , then  $\Gamma \models_{\mathbf{QmbC}^\approx} \varphi$ .*

In order to prove completeness of  $\mathbf{QmbC}^\approx$  w.r.t. swap structures semantics, the proof given in Section 8 will be adapted, in accordance with the argument given in [7, Section 7.7].

We begin by observing that the notion of  $C$ -Henkin theory in  $\mathbf{QmbC}^\approx$  can be defined by adapting Definition 8.1 in an obvious way. The signature obtained from  $\Theta_\approx$  by adding a set  $C$  of new individual constants will be denoted by  $\Theta_C^\approx$ , and the consequence relation in  $\mathbf{QmbC}^\approx$  over that signature will be denoted by  $\vdash_{\mathbf{QmbC}^\approx}^C$ . Clearly, Proposition 8.4 also

holds for  $\mathbf{QmbC}^\approx$ . This result, combined with [7, Theorem 7.5.3] (which can also be easily adapted to  $\mathbf{QmbC}^\approx$ ) produces the following:

**Proposition 10.6.** *Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sen}(\Theta_\approx)$  such that  $\Gamma \not\vdash_{\mathbf{QmbC}^\approx} \varphi$ . Then, there exists a set of sentences  $\Delta \subseteq \text{Sen}(\Theta_\approx^C)$ , for some set  $C$  of new individual constants, such that  $\Gamma \subseteq \Delta$ , it is a  $C$ -Henkin theory in  $\mathbf{QmbC}^\approx$ , and it is maximally non-trivial with respect to  $\varphi$  in  $\mathbf{QmbC}^\approx$  (by restricting  $\vdash_{\mathbf{QmbC}^\approx}^C$  to sentences in  $\text{Sen}(\Theta_\approx^C)$ ).*

**Definition 10.7.** Let  $\Delta \subseteq \text{Sen}(\Theta_\approx)$  be non-trivial in  $\mathbf{QmbC}^\approx$ , that is: there is some sentence  $\varphi$  in  $\text{Sen}(\Theta_\approx)$  such that  $\Delta \not\vdash_{\mathbf{QmbC}^\approx} \varphi$ . Let  $\equiv_\Delta^\approx \subseteq \text{Sen}(\Theta_\approx)^2$  be the relation in  $\text{Sen}(\Theta_\approx)$  defined as follows:  $\alpha \equiv_\Delta^\approx \beta$  iff  $\Delta \vdash_{\mathbf{QmbC}^\approx} \alpha \leftrightarrow \beta$ .

Then  $\equiv_\Delta^\approx$  is an equivalence relation which induces a Boolean algebra  $\mathcal{A}_\Delta^\approx$  whose domain is the quotient set  $A_\Delta^\approx \stackrel{\text{def}}{=} \text{Sen}(\Theta_\approx) / \equiv_\Delta^\approx$  such that the operations are defined as follows (here  $[\alpha]_\Delta^\approx$  denotes the equivalence class of  $\alpha$  w.r.t.  $\equiv_\Delta^\approx$ ):  $[\alpha]_\Delta^\approx \# [\beta]_\Delta^\approx \stackrel{\text{def}}{=} [\alpha \# \beta]_\Delta^\approx$  for any  $\# \in \{\wedge, \vee, \rightarrow\}$ ;  $0_\Delta^\approx \stackrel{\text{def}}{=} [\varphi \wedge \neg\varphi \wedge \circ\varphi]_\Delta^\approx$ , and  $1_\Delta^\approx \stackrel{\text{def}}{=} [\varphi \vee \neg\varphi]_\Delta^\approx$  (for any sentence  $\varphi$ ).

**Definition 10.8.** Let  $\mathcal{A}_\Delta^\approx$  be a Boolean algebra defined as above, and let  $C\mathcal{A}_\Delta^\approx$  be its completion with monomorphism  $*$  (recall Section 8). The full swap structure for  $\mathbf{mbC}$  over  $C\mathcal{A}_\Delta^\approx$  will be denoted by  $\mathcal{B}_\Delta^\approx$ , and the associated Nmatrix will be denoted by  $\mathcal{M}(\mathcal{B}_\Delta^\approx) \stackrel{\text{def}}{=} (\mathcal{B}_\Delta^\approx, D_\Delta^\approx)$ .

**Remark 10.9.** Note that  $(*([\alpha]_\Delta^\approx), *([\beta]_\Delta^\approx), *([\gamma]_\Delta^\approx)) \in D_\Delta^\approx$  iff  $\Delta \vdash_{\mathbf{QmbC}^\approx} \alpha$ .

**Definition 10.10.** (Canonical Structure in  $\mathbf{QmbC}^\approx$ ) Let  $\Delta \subseteq \text{Sen}(\Theta_\approx^C)$  be a non-trivial and  $C$ -Henkin theory in  $\mathbf{QmbC}^\approx$ . Define in the set  $C$  of constants the following relation:  $c \simeq d$  iff  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (c \approx d)$ . By the axioms of equality it follows that  $\simeq$  is an equivalence relation. For any  $c \in C$  let  $\tilde{c} = \{d \in C : c \simeq d\}$  be the equivalence class of  $c$  w.r.t.  $\simeq$ , and let  $U = \{\tilde{c} : c \in C\}$  be the corresponding quotient set. Let  $\mathcal{M}(\mathcal{B}_\Delta^\approx)$  be as in Definition 10.8. The canonical structure induced by  $\Delta$  in  $\mathbf{QmbC}^\approx$  is the structure  $\mathfrak{A}_\Delta^\approx = \langle U, I_{\mathfrak{A}_\Delta^\approx} \rangle$  over  $\Theta_\approx^C$  and  $\mathcal{M}(\mathcal{B}_\Delta^\approx)$  defined as follows:

- if  $c$  is an individual constant in  $\Theta_\approx^C$  then  $I_{\mathfrak{A}_\Delta^\approx}(c) = \tilde{d}$ , where  $d \in C$  is such that  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (c \approx d)$ ;
- if  $f$  is a function symbol,  $I_{\mathfrak{A}_\Delta^\approx}(f) : U^n \rightarrow U$  is given by  $I_{\mathfrak{A}_\Delta^\approx}(f)(\tilde{c}_1, \dots, \tilde{c}_n) = \tilde{c}$ , where  $c \in C$  is such that  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (f(c_1, \dots, c_n) \approx c)$ ;
- if  $P$  is a predicate symbol, then  $I_{\mathfrak{A}_\Delta^\approx}(P)$  is given by

$$I_{\mathfrak{A}_\Delta^\approx}(P)(\tilde{c}_1, \dots, \tilde{c}_n) = (*([P(c_1, \dots, c_n)]_\Delta^\approx), *([\neg P(c_1, \dots, c_n)]_\Delta^\approx), *([\circ P(c_1, \dots, c_n)]_\Delta^\approx)).$$

The proof that  $I_{\mathfrak{A}_\Delta^\approx}$  is well-defined for individual constants and function symbols is similar to that for classical logic (see [12]). Let  $P$  be predicate symbol of arity  $n$  and let  $(\tilde{c}_1, \dots, \tilde{c}_n) \in U^n$ . Let  $d_1, \dots, d_n \in C$  such that  $c_i \simeq d_i$  for every  $1 \leq i \leq n$ . Then  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (c_i \approx d_i)$  for  $1 \leq i \leq n$ . It is immediate that  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (\bigwedge_{i=1}^n (c_i \approx d_i)) \rightarrow (P(c_1, \dots, c_n) \leftrightarrow P(d_1, \dots, d_n))$ , hence  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (P(c_1, \dots, c_n) \leftrightarrow P(d_1, \dots, d_n))$ . Analogously it can be proven that  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (\neg P(c_1, \dots, c_n) \leftrightarrow \neg P(d_1, \dots, d_n))$  and  $\Delta \vdash_{\mathbf{QmbC}^\approx}^C (\circ P(c_1, \dots, c_n) \leftrightarrow \circ P(d_1, \dots, d_n))$ . This shows that  $I_{\mathfrak{A}_\Delta^\approx}(P)$  is well-defined. Moreover, by similar considerations to the ones given after Definition 8.8, it follows that  $I_{\mathfrak{A}_\Delta^\approx}(P)(\tilde{c}_1, \dots, \tilde{c}_n)$  belongs to  $|\mathcal{B}_\Delta^\approx|$  for every  $(\tilde{c}_1, \dots, \tilde{c}_n) \in U^n$ .

**Proposition 10.11.** *Let  $\Delta \subseteq \text{Sen}(\Theta_{\tilde{C}}^{\approx})$  be a non-trivial and  $C$ -Henkin theory in  $\mathbf{QmbC}^{\approx}$  and let  $\mathcal{M}(\mathcal{B}_{\tilde{\Delta}}^{\approx})$  be as in Definition 10.8. Then the canonical structure  $\mathfrak{A}_{\tilde{\Delta}}^{\approx}$  induced by  $\Delta$  in  $\mathbf{QmbC}^{\approx}$  is a structure with standard equality over  $\Theta_{\tilde{C}}^{\approx}$  and  $\mathcal{M}(\mathcal{B}_{\tilde{\Delta}}^{\approx})$ .*

*Proof.* As it was shown above, the mapping  $I_{\mathfrak{A}_{\tilde{\Delta}}^{\approx}}$  is well-defined, and  $I_{\mathfrak{A}_{\tilde{\Delta}}^{\approx}}(P)$  is a function from  $U^n$  to  $|\mathcal{B}_{\tilde{\Delta}}^{\approx}|$ . Let  $\tilde{c}_1, \tilde{c}_2 \in U$ . Then  $(\tilde{c}_1 \approx^{\mathfrak{A}_{\tilde{\Delta}}^{\approx}} \tilde{c}_2) \in D_{\tilde{\Delta}}^{\approx}$  iff  $\Delta \vdash_{\mathbf{QmbC}^{\approx}}^C (c_1 \approx c_2)$ , by Definition 10.10 and by Remark 10.9, iff  $c_1 \simeq c_2$  iff  $\tilde{c}_1 = \tilde{c}_2$ . This shows that  $\mathfrak{A}_{\tilde{\Delta}}^{\approx}$  is indeed a structure with standard equality.  $\square$

**Definition 10.12.** Let  $(\cdot)^{\triangleleft} : (\text{Ter}((\Theta_{\tilde{C}}^{\approx})_U) \cup \text{For}((\Theta_{\tilde{C}}^{\approx})_U)) \rightarrow (\text{Ter}(\Theta_{\tilde{C}}^{\approx}) \cup \text{For}(\Theta_{\tilde{C}}^{\approx}))$  be the mapping recursively defined as in Definition 8.9, but with the following difference:  $(\tilde{c})^{\triangleleft} = d$  for some  $d \in \tilde{c}$  previously chosen, for every  $\tilde{c} \in U$ .

Observe that, if  $s \in \text{Ter}((\Theta_{\tilde{C}}^{\approx})_U) \cup \text{For}((\Theta_{\tilde{C}}^{\approx})_U)$ , then  $(s)^{\triangleleft}$  is the expression in  $\text{Ter}(\Theta_{\tilde{C}}^{\approx}) \cup \text{For}(\Theta_{\tilde{C}}^{\approx})$  obtained from  $s$  by substituting every occurrence of a constant  $\tilde{c}$  by a constant  $d \in \tilde{c}$ . Suppose that  $(\cdot)^{\triangleleft'} : (\text{Ter}((\Theta_{\tilde{C}}^{\approx})_U) \cup \text{For}((\Theta_{\tilde{C}}^{\approx})_U)) \rightarrow (\text{Ter}(\Theta_{\tilde{C}}^{\approx}) \cup \text{For}(\Theta_{\tilde{C}}^{\approx}))$  is defined as  $(\cdot)^{\triangleleft}$ , but now  $(\tilde{c})^{\triangleleft'} = d'$  for another choice of  $d' \in \tilde{c}$  (possibly different to  $d$ ), for every  $\tilde{c} \in U$ . Then, it is easy to prove that  $\Delta \vdash_{\mathbf{QmbC}^{\approx}}^C (\psi)^{\triangleleft} \leftrightarrow (\psi)^{\triangleleft'}$  for every sentence  $\psi$ . This shows that the choice of each  $d \in \tilde{c}$  in order to define  $(\tilde{c})^{\triangleleft}$ , for every  $\tilde{c} \in U$ , is irrelevant.

**Proposition 10.13.** *Let  $\Delta \subseteq \text{Sen}(\Theta_{\tilde{C}}^{\approx})$  be a set of sentences over the signature  $\Theta_{\tilde{C}}^{\approx}$  such that  $\Delta$  is a  $C$ -Henkin theory in  $\mathbf{QmbC}^{\approx}$  which is also maximally non-trivial with respect to  $\varphi$  in  $\mathbf{QmbC}^{\approx}$ , for some sentence  $\varphi$ . Then, the canonical  $\mathbf{QmbC}$ -valuation induced by  $\Delta$  over  $\mathfrak{A}_{\tilde{\Delta}}^{\approx}$  and  $\mathcal{M}(\mathcal{B}_{\tilde{\Delta}}^{\approx})$  (see Definition 8.10) is a  $\mathbf{QmbC}^{\approx}$ -valuation, which will be denoted by  $v_{\tilde{\Delta}}^{\approx}$ , such that  $v_{\tilde{\Delta}}^{\approx}(\psi) \in D_{\tilde{\Delta}}^{\approx}$  iff  $\Delta \vdash_{\mathbf{QmbC}^{\approx}}^C (\psi)^{\triangleleft}$ .*

*Proof.* Observe that, by the very definitions,  $v_{\tilde{\Delta}}^{\approx}(\psi) \in D_{\tilde{\Delta}}^{\approx}$  iff  $\Delta \vdash_{\mathbf{QmbC}^{\approx}}^C (\psi)^{\triangleleft}$  (and, as observed above, ' $v_{\tilde{\Delta}}^{\approx}(\psi) \in D_{\tilde{\Delta}}^{\approx}$ ' does not depend on the choices made by  $(\cdot)^{\triangleleft}$ ). Hence, it suffices to prove that  $v_{\tilde{\Delta}}^{\approx}$  satisfies clause (viii) of Definition 10.4. Thus, let  $\alpha = (x \approx y) \rightarrow (\psi \rightarrow \psi[x\lambda y])$  (where  $y$  is a variable free for  $x$  in  $\varphi$ ) be an instance of axiom (**AxEq2**), and let  $\mu$  be an assignment. Given that  $\Delta$  is a closed theory in  $\mathbf{QmbC}^{\approx}$  over  $\Theta_{\tilde{C}}^{\approx}$ , it follows that  $\Delta \vdash_{\mathbf{QmbC}^{\approx}}^C (\hat{\mu}(\alpha))^{\triangleleft}$ . Then  $v_{\tilde{\Delta}}^{\approx}(\hat{\mu}(\alpha)) \in D_{\tilde{\Delta}}^{\approx}$ , showing that  $v_{\tilde{\Delta}}^{\approx}$  satisfies clause (viii).  $\square$

**Theorem 10.14.** (Completeness of  $\mathbf{QmbC}^{\approx}$  restricted to sentences w.r.t. first-order swap structures with standard equality) *Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sen}(\Theta_{\approx})$ . If  $\Gamma \models_{\mathbf{QmbC}^{\approx}} \varphi$  then  $\Gamma \vdash_{\mathbf{QmbC}^{\approx}} \varphi$ .*

*Proof.* Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sen}(\Theta_{\approx})$  such that  $\Gamma \not\vdash_{\mathbf{QmbC}^{\approx}} \varphi$ . By Proposition 10.6, there exists a set of sentences  $\Delta \subseteq \text{Sen}(\Theta_{\tilde{C}}^{\approx})$ , for some set  $C$  of new individual constants, such that  $\Gamma \subseteq \Delta$ ,  $\Delta$  is a  $C$ -Henkin theory in  $\mathbf{QmbC}^{\approx}$ , and it is maximally non-trivial with respect to  $\varphi$  in  $\mathbf{QmbC}^{\approx}$  (by restricting  $\vdash_{\mathbf{QmbC}^{\approx}}^C$  to sentences in  $\text{Sen}(\Theta_{\tilde{C}}^{\approx})$ ).

Now, let  $\mathcal{M}(\mathcal{B}_{\tilde{\Delta}}^{\approx})$ ,  $\mathfrak{A}_{\tilde{\Delta}}^{\approx}$  and  $v_{\tilde{\Delta}}^{\approx}$  be as in Definitions 10.8 and 10.10, and as in Proposition 10.13, respectively. Then,  $v_{\tilde{\Delta}}^{\approx}(\alpha) \in D_{\tilde{\Delta}}^{\approx}$  iff  $\Delta \vdash_{\mathbf{QmbC}^{\approx}}^C \alpha$ , for every  $\alpha$  in  $\text{Sen}(\Theta_{\tilde{C}}^{\approx})$  (by observing that  $(\alpha)^{\triangleleft} = \alpha$  if  $\alpha \in \text{Sen}(\Theta_{\tilde{C}}^{\approx})$ ). From this,  $v_{\tilde{\Delta}}^{\approx}[\Gamma] \subseteq D_{\tilde{\Delta}}^{\approx}$  and  $v_{\tilde{\Delta}}^{\approx}(\varphi) \notin D_{\tilde{\Delta}}^{\approx}$ . Finally, let  $\mathfrak{A}$  and  $v$  be the restriction to  $\Theta_{\approx}$  of  $\mathfrak{A}_{\tilde{\Delta}}^{\approx}$  and  $v_{\tilde{\Delta}}^{\approx}$ , respectively. Then,  $\mathfrak{A}$  is a structure with standard equality over  $\mathcal{M}(\mathcal{B}_{\tilde{\Delta}}^{\approx})$ , and  $v$  is a valuation for  $\mathbf{QmbC}^{\approx}$  over  $\mathfrak{A}$  and  $\mathcal{M}(\mathcal{B}_{\tilde{\Delta}}^{\approx})$  such that  $v[\Gamma] \subseteq D_{\tilde{\Delta}}^{\approx}$  but  $v(\varphi) \notin D_{\tilde{\Delta}}^{\approx}$ . This shows that  $\Gamma \not\vdash_{\mathbf{QmbC}^{\approx}} \varphi$ .  $\square$

By using universal closure, as it was done in Corollary 8.14, a general completeness result can be obtained for  $\mathbf{QmbC}^{\approx}$ :

**Corollary 10.15.** (Completeness of  $\mathbf{QmbC}^\approx$  w.r.t. first-order swap structures with standard equality) *Let  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta_\approx)$ . If  $\Gamma \models_{\mathbf{QmbC}^\approx} \varphi$  then  $\Gamma \vdash_{\mathbf{QmbC}^\approx} \varphi$ .*

## 11 Completeness of $\mathbf{QmbC}^\approx$ w.r.t. structures with standard equality over $\mathcal{M}_5$

In this section the adequacy of  $\mathbf{QmbC}$  w.r.t. first-order swap structures stated in Theorem 9.6 will be extended to  $\mathbf{QmbC}^\approx$ . In order to do this, some definitions taken from Section 9 will be adapted to  $\mathbf{QmbC}^\approx$ , by following the approach in [7, Section 7.7] with small modifications. In particular, [7, Definition 7.7.3] will be slightly adapted as follows:

**Definition 11.1.** An *interpretation* for  $\mathbf{QmbC}^\approx$  over a signature  $\Theta_\approx$  is a pair  $\langle \mathbf{A}, \rho \rangle$  such that  $\mathbf{A}$  is a standard Tarskian first-order structure with standard equality over  $\Theta_\approx$ <sup>9</sup> and  $\rho$  is a bivaluation for  $\mathbf{QmbC}$  over  $\mathbf{A}$ . The *consequence relation*  $\models_{\mathbf{QmbC}^\approx}^2$  of  $\mathbf{QmbC}^\approx$  w.r.t. interpretations is given by:  $\Gamma \models_{\mathbf{QmbC}^\approx}^2 \varphi$  if, for every interpretation  $\langle \mathbf{A}, \rho \rangle$  for  $\mathbf{QmbC}^\approx$ :  $\rho(\widehat{\mu}(\gamma)) = 1$  for every  $\gamma \in \Gamma$  and every  $\mu$  implies that  $\rho(\widehat{\mu}(\varphi)) = 1$  for every  $\mu$ .

**Remark 11.2.**

(1) In [7, Definition 7.7.3] it was introduced the notion of  $\mathbf{QmbC}^\approx$ -valuations, which are bivaluations for  $\mathbf{QmbC}$  over standard Tarskian structures  $\mathbf{A}$  over  $\Theta_\approx$  satisfying for  $\approx$ , instead of (*vPred*), the following clauses:

(**vEq1**)  $\rho(t_1 \approx t_2) = 1$  iff  $\llbracket t_1 \rrbracket^{\widehat{\mathbf{A}}} = \llbracket t_2 \rrbracket^{\widehat{\mathbf{A}}}$  for every  $t_1, t_2 \in \text{CTer}(\Theta_\approx)$ ;

(**vEq2**)  $\rho(\bar{a} \approx \bar{b}) = 1$  implies  $\rho(\alpha[x, y/\bar{a}, \bar{b}]) = \rho(\alpha[x \lambda y][x, y/\bar{a}, \bar{b}])$  for every  $a, b \in U$ , if  $y$  is a variable free for  $x$  in  $\alpha$ .

It is easy to see that (*vEq2*) is derivable from (*vEq1*). Indeed, suppose that  $\rho(\bar{a} \approx \bar{b}) = 1$ . Hence  $a = \llbracket \bar{a} \rrbracket^{\widehat{\mathbf{A}}} = \llbracket \bar{b} \rrbracket^{\widehat{\mathbf{A}}} = b$ , by clause (*vEq1*). But then  $\bar{a} = \bar{b}$ , which implies that  $\alpha[x, y/\bar{a}, \bar{b}] = \alpha[x \lambda y][x, y/\bar{a}, \bar{b}]$ . Thus,  $\rho(\alpha[x, y/\bar{a}, \bar{b}]) = \rho(\alpha[x \lambda y][x, y/\bar{a}, \bar{b}])$ , showing that  $\rho$  also satisfies (*vEq2*).

(2) Let  $\langle \mathbf{A}, \rho \rangle$  be an interpretation for  $\mathbf{QmbC}^\approx$  as in Definition 11.1. Then, by (*vPred*) applied to  $\approx$  (recall Definition 9.1), and by the fact that  $\approx^{\mathbf{A}} = \{(a, a) : a \in U\}$ , it follows that  $\rho$  satisfies clause (*vEq1*), hence it also satisfies (*vEq2*), by item (1) above. This means that  $\langle \mathbf{A}, \rho \rangle$  is an interpretation for  $\mathbf{QmbC}^\approx$  in the sense of [7, Definition 7.7.3]. Conversely, if  $\langle \mathbf{A}, \rho \rangle$  is an interpretation for  $\mathbf{QmbC}^\approx$  in the sense of [7, Definition 7.7.3] let  $\mathbf{A}'$  be the standard Tarskian structure over  $\Theta_\approx$  obtained from  $\mathbf{A}$  by setting  $\approx^{\mathbf{A}'} \stackrel{\text{def}}{=} \{(a, a) : a \in U\}$ . Hence  $\langle \mathbf{A}', \rho \rangle$  is an interpretation for  $\mathbf{QmbC}$  as in Definition 11.1, since  $\rho$  satisfies (*vEq1*) and so it satisfies (*vPred*) applied to  $\approx$ . This shows that our presentation is equivalent to that of [7].

**Theorem 11.3.** (Adequacy of  $\mathbf{QmbC}^\approx$  w.r.t. interpretations, [7, Theorem 7.7.5]) *If  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta_\approx)$  then:  $\Gamma \vdash_{\mathbf{QmbC}^\approx} \varphi$  iff  $\Gamma \models_{\mathbf{QmbC}^\approx}^2 \varphi$ .*<sup>10</sup>

Theorem 9.5 can be easily extended to  $\mathbf{QmbC}^\approx$ :

<sup>9</sup>That is,  $\mathbf{A}$  is a standard Tarskian first-order structure over signature  $\Theta_\approx$ , as considered in Section 9, in which the equality predicate  $\approx$  is interpreted as the identity:  $\approx^{\mathbf{A}} \stackrel{\text{def}}{=} \{(a, a) : a \in U\}$ .

<sup>10</sup>As in the case of  $\mathbf{QmbC}$ , in [7, Theorem 7.7.5] it was obtained adequacy of  $\mathbf{QmbC}^\approx$  w.r.t. interpretations, but only for sentences. Once again, adequacy for general formulas follows from that result by using universal closure.

**Theorem 11.4.** *Let  $\mathcal{I} = \langle \mathbf{A}, \rho \rangle$  be an interpretation for  $\mathbf{QmbC}^\approx$  over a signature  $\Theta_\approx$ . Then, it induces a first-order structure with standard equality  $\mathfrak{A}_\mathcal{I}$  over  $\mathcal{M}_5$  and  $\Theta_\approx$ , and a  $\mathbf{QmbC}^\approx$ -valuation  $v_\approx^\rho$  over  $\mathfrak{A}_\mathcal{I}$  and  $\mathcal{M}_5$  given by  $v_\approx^\rho(\alpha) \stackrel{\text{def}}{=} (\rho(\alpha), \rho(\neg\alpha), \rho(\circ\alpha))$  such that:  $\rho(\alpha) = 1$  iff  $v_\approx^\rho(\alpha) \in \mathbf{D}$ , for every sentence  $\alpha \in \text{Sen}(\Theta_\approx)$ .*

*Proof.* Let  $\mathcal{I} = \langle \mathbf{A}, \rho \rangle$  be an interpretation for  $\mathbf{QmbC}^\approx$  over signature  $\Theta_\approx$ . Consider the first-order structure  $\mathfrak{A}_\mathcal{I}$  over  $\mathcal{M}_5$  and  $\Theta_\approx$  obtained from  $\mathbf{A}$  as in the proof of Theorem 9.5. In particular,  $I_{\mathfrak{A}_\mathcal{I}}(\approx)(a_1, a_2) = v_\approx^\rho(\bar{a}_1 \approx \bar{a}_2)$ , where  $v_\approx^\rho : \text{Sen}(\Theta_\approx) \rightarrow |\mathcal{M}_5|$  is given by  $v_\approx^\rho(\alpha) = (\rho(\alpha), \rho(\neg\alpha), \rho(\circ\alpha))$ , for every  $\alpha \in \text{Sen}(\Theta_\approx)$ . It is immediate to see that  $\rho(\alpha) = 1$  iff  $v_\approx^\rho(\alpha) \in \mathbf{D}$ , for every  $\alpha \in \text{Sen}(\Theta_\approx)$ . From this,  $I_{\mathfrak{A}_\mathcal{I}}(\approx)(a_1, a_2) = v_\approx^\rho(\bar{a}_1 \approx \bar{a}_2) \in \mathbf{D}$  iff  $\rho(\bar{a}_1 \approx \bar{a}_2) = 1$  iff  $a_1 = a_2$ , since  $\rho$  satisfies clause (vEq1), by Remark 11.2. This shows that  $\mathfrak{A}_\mathcal{I}$  is a first-order structure with standard equality over  $\mathcal{M}_5$  and  $\Theta_\approx$ .

In order to see that  $v_\approx^\rho$  is a  $\mathbf{QmbC}^\approx$ -valuation  $v_\approx^\rho$  over  $\mathfrak{A}_\mathcal{I}$  and  $\mathcal{M}_5$  observe that  $v_\approx^\rho$  satisfies clause (i) of Definition 6.9 for every predicate symbol in  $\Theta$ . Concerning the equality predicate  $\approx$  it is easy to prove that that, by the axioms of equality, the properties of bivaluations for  $\mathbf{QmbC}$ , and the fact that  $a = \llbracket \bar{a} \rrbracket^{\widehat{\mathfrak{A}}} = \llbracket \bar{a} \rrbracket^{\widehat{\mathbf{A}}}$  for every  $a \in U$  (in particular, for  $a = \llbracket t \rrbracket^{\widehat{\mathfrak{A}}}$ , for  $t \in \text{CTer}(\Theta_\approx)$ ),  $v_\approx^\rho(t_1 \approx t_2) = v_\approx^\rho(\llbracket t_1 \rrbracket^{\widehat{\mathfrak{A}}} \approx \llbracket t_2 \rrbracket^{\widehat{\mathfrak{A}}}) = I_{\mathfrak{A}_\mathcal{I}}(\approx)(\llbracket t_1 \rrbracket^{\widehat{\mathfrak{A}}}, \llbracket t_2 \rrbracket^{\widehat{\mathfrak{A}}})$ . This shows that  $v_\approx^\rho$  also satisfies clause (i) of Definition 6.9 for the equality predicate  $\approx$ . By the proof of Theorem 9.5, it follows that  $v_\approx^\rho$  satisfies the other clauses of Definition 6.9 for  $\mathbf{QmbC}$ -valuation over  $\mathfrak{A}_\mathcal{I}$  and  $\mathcal{M}_5$ . It remains to prove that  $v_\approx^\rho$  is a  $\mathbf{QmbC}^\approx$ -valuation over  $\mathfrak{A}_\mathcal{I}$  and  $\mathcal{M}_5$ , that is,  $v_\approx^\rho(\widehat{\mu}(\alpha)) \in \mathbf{D}$  for every instance  $\alpha$  of axiom (AxEq2) and every assignment  $\mu$ . But this is easy to prove, by the properties of  $\rho$  and by an argument similar to that presented in Remark 11.2(1). This concludes the proof.  $\square$

As an immediate consequence of Theorems 10.5, 11.3 and 11.4:

**Theorem 11.5.** (Adequacy of  $\mathbf{QmbC}^\approx$  w.r.t. first-order structures with standard equality over  $\mathcal{M}_5$ ) *For every set  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta_\approx)$ :  $\Gamma \vdash_{\mathbf{QmbC}^\approx} \varphi$  iff  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}_5)}^\approx \varphi$  for every structure  $\mathfrak{A}$  with standard equality over  $\Theta_\approx$  and  $\mathcal{M}_5$ .*

## 12 First order twist structures based on the logic LFI<sub>0</sub>

The generalization of swap structures semantics to other quantified **LFI**s, defined as axiomatic extensions of **QmbC**, can be easily obtained. Indeed, by analyzing the swap structures semantics for axiomatic extensions of **mbC** given in [7, Section 6.5] (see also [14]), as well as the first-order version of such extensions proposed in [7, Section 7.8], it is immediate how to obtain first-order swap structures for all these logics. Thus, it is immediate to define **QmbCciw**, **QmbCci**, **QbC** and **QCi**, the quantified version of **mbCciw**, **mbCci**, **bC** and **Ci**, respectively, as well as the corresponding extensions of them by adding the standard equality. All these logics are characterized by means of first-order structures defined over 3-valued swap structures.<sup>11</sup>

<sup>11</sup>As proved by Avron in [2], the extension **mbCcl** of **mbC** by adding da Costa's axiom (cl):  $\neg(\varphi \wedge \neg\varphi) \rightarrow \circ\varphi$  cannot be characterized by a single finite Nmatrix. This negative result also holds for **Cila**, the presentation of a Costa's system  $C_1$  in the language of **LFI**s, hence it holds for  $C_1$  itself. Thus, these systems lies outside the scope of the present framework. A discussion about this question can be found

Instead of analyzing in this section these axiomatic extensions of **QmbC**, together with the corresponding swap structures semantics (a straightforward exercise), the first-order version of a quite interesting axiomatic extension of **mbC**, the logic **LFII**<sub>o</sub>, will be analyzed with full detail. This logic can be semantically characterized by a 3-valued logical matrix called **LFII**' , which is equivalent (up to language) to several 3-valued paraconsistent logics such as the well-known da Costa-D'Ottaviano logic **J3** and Carnielli-Marcos-de Amo logic **LFII**. From this, it follows that **LFII**<sub>o</sub> is algebraizable in the sense of Blok and Pigozzi (see [7, Chapter 4] for a discussion about this logic). The interesting point is that, as proved in [14], the swap structures for **LFII**<sub>o</sub> turn out to be deterministic, thus becoming twist structures, which represent the algebraic semantics for **LFII**<sub>o</sub>.<sup>12</sup> Because of this, the first-order structures for the first-order extension of **LFII**<sub>o</sub> presented here are based on twist structures.

**Definition 12.1.** ([7, Definition 4.4.39]) Let  $\mathcal{M}_{LFII} = \langle \mathcal{A}_{LFII}, \{1, \frac{1}{2}\} \rangle$  be the logical matrix such that  $\mathcal{A}_{LFII}$  is the 3-valued algebra over  $\Sigma$  with domain  $M = \{1, \frac{1}{2}, 0\}$ , where the operators are defined as follows:

$\wedge$	1	$\frac{1}{2}$	0	$\vee$	1	$\frac{1}{2}$	0	$\rightarrow$	1	$\frac{1}{2}$	0	$\neg$		$\circ$	
1	1	$\frac{1}{2}$	0	1	1	1	1	1	1	$\frac{1}{2}$	0	1	0	1	1
$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0	$\frac{1}{2}$	1	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	1	$\frac{1}{2}$	0	$\frac{1}{2}$	$\frac{1}{2}$	$\frac{1}{2}$	0
0	0	0	0	0	1	$\frac{1}{2}$	0	0	1	1	1	0	1	0	1

The logic associated to the logical matrix  $\mathcal{M}_{LFII}$  is called **LFII**'.

Recall that, by definition, the consequence relation  $\models_{\mathbf{LFII}'}$  of **LFII**' is given as follows: for every  $\Gamma \cup \{\alpha\} \subseteq \mathcal{L}_\Sigma$ ,  $\Gamma \models_{\mathbf{LFII}'} \alpha$  iff, for every homomorphism  $v : \mathcal{L}_\Sigma \rightarrow \mathcal{A}_{LFII}$  of algebras over  $\Sigma$ , if  $v[\Gamma] \subseteq \{1, \frac{1}{2}\}$  then  $v(\alpha) \in \{1, \frac{1}{2}\}$ .

A sound and complete Hilbert calculus for **LFII**' , called **LFII**<sub>o</sub>, was introduced in [7]. This calculus is an axiomatic extension of **mbC**.

**Definition 12.2.** ([7, Definition 4.4.41]) The Hilbert calculus **LFII**<sub>o</sub> over  $\Sigma$  is obtained from **mbC** by adding the following axioms:

- (ci)  $\neg \circ \alpha \rightarrow (\alpha \wedge \neg \alpha)$
- (dneg)  $\neg \neg \alpha \leftrightarrow \alpha$
- (neg $\vee$ )  $\neg(\alpha \vee \beta) \leftrightarrow (\neg \alpha \wedge \neg \beta)$
- (neg $\wedge$ )  $\neg(\alpha \wedge \beta) \leftrightarrow (\neg \alpha \vee \neg \beta)$
- (neg  $\rightarrow$ )  $\neg(\alpha \rightarrow \beta) \leftrightarrow (\alpha \wedge \neg \beta)$

**Theorem 12.3.** ([7, Theorem 4.4.45]) *The logic **LFII**<sub>o</sub> is sound and complete w.r.t. the matrix semantics of **LFII**' :  $\Gamma \vdash_{\mathbf{LFII}_o} \alpha$  iff  $\Gamma \models_{\mathbf{LFII}'} \alpha$*

**Remark 12.4.** By Propositions 3.1.10 and 3.2.3 in [7], and given that **LFII**<sub>o</sub> contains axiom (ci), it follows that  $\vdash_{\mathbf{LFII}_o} \circ \alpha \leftrightarrow \sim(\alpha \wedge \neg \alpha)$  (here,  $\sim$  is the classical negation defined as in Remark 8.2). On the other hand, taking into account that  $\vdash_{\mathbf{mbC}} (\alpha \wedge \neg \alpha) \rightarrow \neg \circ \alpha$ , it follows from (ci) that  $\vdash_{\mathbf{LFII}_o} \neg \circ \alpha \leftrightarrow (\alpha \wedge \neg \alpha)$ .

in [7, Section 6.5]. On the other hand, **Cila** is not algebraizable in the sense of Blok and Pigozzi (consult, for instance, [11, Section 3.12]). As a consequence of this, none of the logics **mbC**, **mbCciw**, **mbCci**, **bC** and **Ci** is algebraizable.

<sup>12</sup>Twist structures were independently introduced by Fidel [22] and Vakarelov [30]. However, as observed by Cignoli in [13], the basic algebraic ideas underlying twist structures were firstly introduced by Kalman in [25].

Since  $\mathbf{LFI1}_\circ$  is an axiomatic extension of  $\mathbf{mbC}$ , its first-order extension  $\mathbf{QLFI1}_\circ$  can be defined as an axiomatic extension of  $\mathbf{QmbC}$ ,<sup>13</sup> hence the semantics of first-order swap structures for  $\mathbf{QmbC}$  given in the previous sections can be adapted to  $\mathbf{QLFI1}_\circ$ , obtaining so a semantics based on first-order twist structures. Indeed, as shown in [14], each multioperation in the corresponding swap structures for  $\mathbf{LFI1}_\circ$  is deterministic, and so these swap structures are twist structures (which are ordinary algebras presented in a particular form).

**Definition 12.5.** Let  $\mathcal{A} = \langle A, \wedge, \vee, \rightarrow, 0, 1 \rangle$  be a Boolean algebra. The *twist domain* generated by  $\mathcal{A}$  is the set  $T_{\mathcal{A}} = \{(z_1, z_2) \in A \times A : z_1 \vee z_2 = 1\}$ .

**Definition 12.6.** ([14, Definition 9.2]) Let  $\mathcal{A}$  be a Boolean algebra. The *twist structure* for  $\mathbf{LFI1}_\circ$  over  $\mathcal{A}$  is the algebra  $\mathcal{T}_{\mathcal{A}} = \langle T_{\mathcal{A}}, \tilde{\wedge}, \tilde{\vee}, \tilde{\rightarrow}, \tilde{\sim}, \tilde{\circ} \rangle$  over  $\Sigma$  such that the operations are defined as follows, for every  $(z_1, z_2), (w_1, w_2) \in T_{\mathcal{A}}$ :

- (i)  $(z_1, z_2) \tilde{\wedge} (w_1, w_2) = (z_1 \wedge w_1, z_2 \vee w_2)$ ;
- (ii)  $(z_1, z_2) \tilde{\vee} (w_1, w_2) = (z_1 \vee w_1, z_2 \wedge w_2)$ ;
- (iii)  $(z_1, z_2) \tilde{\rightarrow} (w_1, w_2) = (z_1 \rightarrow w_1, z_1 \wedge w_2)$ ;
- (iv)  $\tilde{\sim}(z_1, z_2) = (z_2, z_1)$ ;
- (v)  $\tilde{\circ}(z_1, z_2) = (\sim(z_1 \wedge z_2), z_1 \wedge z_2)$ .<sup>14</sup>

The intuitive meaning of a snapshot  $(z_1, z_2)$  in  $T_{\mathcal{A}}$  is that  $z_1$  represents a value, in a given Boolean algebra, for the evidence for  $\varphi$ , while  $z_2$  represents a value for the evidence against  $\varphi$  (or a value for the evidence for  $\neg\varphi$ ).

**Definition 12.7.** The logical matrix associated to the twist structure  $\mathcal{T}_{\mathcal{A}}$  is  $\mathcal{MT}_{\mathcal{A}} = \langle \mathcal{T}_{\mathcal{A}}, D_{\mathcal{A}} \rangle$  where  $D_{\mathcal{A}} = \{(z_1, z_2) \in T_{\mathcal{A}} : z_1 = 1\} = \{(1, a) : a \in A\}$ . The consequence relation associated to  $\mathcal{MT}_{\mathcal{A}}$  will be denoted by  $\models_{\mathcal{T}_{\mathcal{A}}}$ , namely:  $\Gamma \models_{\mathcal{T}_{\mathcal{A}}} \alpha$  iff, for every homomorphism  $h : \mathcal{L}_{\Sigma} \rightarrow T_{\mathcal{A}}$  of algebras over  $\Sigma$ , if  $h(\gamma) \in D_{\mathcal{A}}$  for every  $\gamma \in \Gamma$  then  $h(\alpha) \in D_{\mathcal{A}}$ . Let  $\mathbb{M}_{\mathbf{LFI1}}$  be the class of logical matrices  $\mathcal{MT}_{\mathcal{A}}$ , for any Boolean algebra  $\mathcal{A}$ . The *twist consequence relation* for  $\mathbf{LFI1}_\circ$  is the consequence relation  $\models_{\mathbb{M}_{\mathbf{LFI1}}}$  associated to  $\mathbb{M}_{\mathbf{LFI1}}$ , namely:  $\Gamma \models_{\mathbb{M}_{\mathbf{LFI1}}} \alpha$  iff  $\Gamma \models_{\mathcal{T}_{\mathcal{A}}} \alpha$  for every Boolean algebra  $\mathcal{A}$ .

**Remark 12.8.** In [14, Theorem 9.6] it was shown that  $\mathbf{LFI1}_\circ$  is sound and complete w.r.t. twist structures semantics, namely:  $\Gamma \vdash_{\mathbf{LFI1}_\circ} \alpha$  iff  $\Gamma \models_{\mathbb{M}_{\mathbf{LFI1}}} \alpha$ , for every set of formulas  $\Gamma \cup \{\alpha\}$ . On the other hand, if  $\mathcal{A}_2$  is the two-element Boolean algebra with domain  $\{0, 1\}$  then  $\mathcal{T}_{\mathcal{A}_2}$  consists of three elements:  $(1, 0)$ ,  $(1, 1)$  and  $(0, 1)$ . By identifying these elements with  $1, \frac{1}{2}$  and  $0$ , respectively, then  $\mathcal{T}_{\mathcal{A}_2}$  coincides with the 3-valued algebra  $\mathcal{A}_{\mathbf{LFI1}}$  underlying the matrix  $\mathcal{M}_{\mathbf{LFI1}}$  (recall Definition 12.1). Moreover,  $\mathcal{MT}_{\mathcal{A}_2}$  coincides with  $\mathcal{M}_{\mathbf{LFI1}}$ . Taking into consideration Theorem 12.3, this situation is analogous to the semantical characterization of  $\mathbf{mbC}$  w.r.t. the 5-element swap structure over  $\mathcal{A}_2$ : it is enough to consider the structure induced by  $\mathcal{A}_2$  in order to characterize the logic.

A first-order version of  $\mathbf{LFI1}_\circ$ , which will be called  $\mathbf{QLFI1}_\circ$ , can be easily defined from  $\mathbf{QmbC}$ .

**Definition 12.9.** Let  $\Theta$  be a first-order signature. The logic  $\mathbf{QLFI1}_\circ$  is obtained from  $\mathbf{QmbC}$  by deleting axiom **(Ax14)** and by adding axioms **(ci)**, **(dneg)**, **(neg $\vee$ )**, **(neg $\wedge$ )** and **(neg $\rightarrow$ )** from  $\mathbf{LFI1}_\circ$ , plus the following:

<sup>13</sup>As we shall see in Remark 12.10, axiom **(Ax14)** will be redundant.

<sup>14</sup>Here,  $\sim$  denotes the Boolean complement  $\sim x = x \rightarrow 0$ .



$$(\mathbf{Ax}\neg\exists) \quad \neg\exists x\varphi \leftrightarrow \forall x\neg\varphi$$

$$(\mathbf{Ax}\neg\forall) \quad \neg\forall x\varphi \leftrightarrow \exists x\neg\varphi$$

**Remark 12.10.** Observe that  $\mathbf{QLFI1}_\circ$  can be alternatively defined as the Hilbert calculus obtained from  $\mathbf{LFI1}_\circ$  by adding axioms  $(\mathbf{Ax12})$  and  $(\mathbf{Ax13})$  from Definition 5.4,  $(\mathbf{Ax}\neg\exists)$  and  $(\mathbf{Ax}\neg\forall)$  above, and the inference rules  $(\exists\text{-In})$  and  $(\forall\text{-In})$  from Definition 5.4. The fact that axiom  $(\mathbf{Ax14})$  is no longer required is justified by the fact that it can now be derived from the other axioms. This can be proved easily after obtaining the completeness of  $\mathbf{QLFI1}_\circ$  w.r.t. twist structures semantics, since axiom  $(\mathbf{Ax14})$  is valid w.r.t. that semantics.

The consequence relation of  $\mathbf{QLFI1}_\circ$  will be denoted by  $\vdash_{\mathbf{QLFI1}_\circ}$ . Since  $\mathbf{QLFI1}_\circ$  does not add inference rules to  $\mathbf{QmbC}$ , it satisfies a deduction meta-theorem (DMT) analogous to  $\mathbf{QmbC}$  (see Theorem 5.6).

Now, the swap structures semantics for  $\mathbf{QmbC}$  can be adapted to  $\mathbf{QLFI1}_\circ$ , taking into account that the swap structures for  $\mathbf{LFI1}_\circ$  are exactly the twist structures introduced in Definition 12.6. This leads us to the following definition:

**Definition 12.11.** let  $\mathcal{A}$  be a complete Boolean algebra. Let  $\mathcal{MT}_{\mathcal{A}}$  be the logical matrix associated to the twist structure  $\mathcal{T}_{\mathcal{A}}$  for  $\mathbf{LFI1}_\circ$ , and let  $\Theta$  be a first-order signature. A (first-order) *structure* over  $\mathcal{MT}_{\mathcal{A}}$  and  $\Theta$ , or a  $\mathbf{QLFI1}_\circ$ -*structure* over  $\Theta$ , is a pair  $\mathfrak{A} = \langle U, I_{\mathfrak{A}} \rangle$  as in Definition 6.1, but now  $I_{\mathfrak{A}}(P)$  is a function from  $U^n$  to  $T_{\mathcal{A}}$ , for each predicate symbol  $P$  of arity  $n$ .

**Notation 12.12.** As it was done with  $\mathbf{QmbC}$ ,  $c^{\mathfrak{A}}$ ,  $f^{\mathfrak{A}}$  and  $P^{\mathfrak{A}}$  will denote the interpretation of an individual constant symbol  $c$ , a function symbol  $f$  and a predicate symbol  $P$ , respectively.

The notion of assignment over a  $\mathbf{QLFI1}_\circ$ -structure is as in Definition 6.2. The notion of interpretation  $\llbracket t \rrbracket_{\mu}^{\mathfrak{A}}$  of a term  $t$  given a structure  $\mathfrak{A}$  and an assignment  $\mu$  is identical to the one described in Definition 6.3. Given  $\mathfrak{A}$ , the structure  $\widehat{\mathfrak{A}} = \langle U, I_{\widehat{\mathfrak{A}}} \rangle$  over  $\Theta_U$  is defined analogously to the case of  $\mathbf{QmbC}$  (recall Definition 6.5).

**Notation 12.13.** By adapting Notation 3.3, if  $z \in T_{\mathcal{A}}$  then  $(z)_1$  and  $(z)_2$ , or simply  $z_1$  and  $z_2$ , will denote the first and second coordinates of  $z$ , respectively.

As it was discussed after Definition 6.8, in order to obtain a single denotation (truth-value) for a formula in  $\mathbf{QmbC}$ , a given interpretation and an assignment are not enough: valuations are necessary in order to choose a unique denotation, in case the formula is complex (that is, if it contains connectives or quantifiers). The case of  $\mathbf{QLFI1}_\circ$  is different, since twist structures are deterministic (that is, they are ordinary algebras). This being so, from a given denotation for the atomic formulas, the denotation for complex formulas is uniquely determined from the denotation of its components, which is in line with the traditional approach to first-order algebraic logic originated by Mostowski. Because of this, valuations over structures are no longer necessary for  $\mathbf{QLFI1}_\circ$ , and a structure  $\mathfrak{A}$  will assign a single denotation (truth-value), denoted by  $\llbracket \varphi \rrbracket^{\mathfrak{A}}$ , to each sentence  $\varphi$ . This lead us to the following definition:

**Definition 12.14.** ( $\mathbf{QLFI1}_\circ$  interpretation maps) Let  $\mathcal{A}$  be a complete Boolean algebra, and let  $\mathfrak{A}$  be a structure over  $\mathcal{MT}_{\mathcal{A}}$  and  $\Theta$ . The *interpretation map* for  $\mathbf{QLFI1}_\circ$  over  $\mathfrak{A}$  and  $\mathcal{MT}_{\mathcal{A}}$  is a function  $\llbracket \cdot \rrbracket^{\mathfrak{A}} : \text{Sen}(\Theta_U) \rightarrow T_{\mathcal{A}}$  satisfying the following clauses (using

Notation 12.13 in clauses (iv) and (v):

- (i)  $\llbracket P(t_1, \dots, t_n) \rrbracket^{\mathfrak{A}} = P^{\mathfrak{A}}(\llbracket t_1 \rrbracket^{\mathfrak{A}}, \dots, \llbracket t_n \rrbracket^{\mathfrak{A}})$ , if  $P(t_1, \dots, t_n)$  is atomic;
- (ii)  $\llbracket \# \varphi \rrbracket^{\mathfrak{A}} = \# \llbracket \varphi \rrbracket^{\mathfrak{A}}$ , for every  $\# \in \{\neg, \circ\}$ ;
- (iii)  $\llbracket \varphi \# \psi \rrbracket^{\mathfrak{A}} = \llbracket \varphi \rrbracket^{\mathfrak{A}} \# \llbracket \psi \rrbracket^{\mathfrak{A}}$ , for every  $\# \in \{\wedge, \vee, \rightarrow\}$ ;
- (iv)  $\llbracket \forall x \varphi \rrbracket^{\mathfrak{A}} = (\bigwedge_{a \in U} (\llbracket \varphi[x/\bar{a}] \rrbracket^{\mathfrak{A}})_1, \bigvee_{a \in U} (\llbracket \varphi[x/\bar{a}] \rrbracket^{\mathfrak{A}})_2)$ ;
- (v)  $\llbracket \exists x \varphi \rrbracket^{\mathfrak{A}} = (\bigvee_{a \in U} (\llbracket \varphi[x/\bar{a}] \rrbracket^{\mathfrak{A}})_1, \bigwedge_{a \in U} (\llbracket \varphi[x/\bar{a}] \rrbracket^{\mathfrak{A}})_2)$ .

The definition of the interpretation of the quantifiers in  $\mathbf{QLFI1}_\circ$  is coherent with the fact that  $\mathcal{T}_{\mathcal{A}}$  (ordered by:  $z \leq w$  iff  $z_1 \leq w_1$  and  $z_2 \geq w_2$ ) is a complete lattice (since  $\mathcal{A}$  is a complete Boolean algebra), in which

$$\bigwedge_{i \in I} z_i = (\bigwedge_{i \in I} (z_i)_1, \bigvee_{i \in I} (z_i)_2), \text{ and}$$

$$\bigvee_{i \in I} z_i = (\bigvee_{i \in I} (z_i)_1, \bigwedge_{i \in I} (z_i)_2)$$

for every family  $(z_i)_{i \in I}$  in  $T_{\mathcal{A}}$ . Note that  $1_{\mathcal{T}_{\mathcal{A}}} =_{\text{def}} (1, 0)$  and  $0_{\mathcal{T}_{\mathcal{A}}} =_{\text{def}} (0, 1)$  are the top and bottom elements of  $\mathcal{T}_{\mathcal{A}}$ , respectively.

**Definition 12.15.** Let  $\mathcal{A}$  be a complete Boolean algebra,  $\mathfrak{A}$  a structure over  $\mathcal{MT}_{\mathcal{A}}$  and  $\Theta$ , and let  $\mu$  be an assignment over  $\mathfrak{A}$ . The *extended interpretation map*  $\llbracket \cdot \rrbracket_{\mu}^{\mathfrak{A}} : \text{For}(\Theta_U) \rightarrow T_{\mathcal{A}}$  is given by  $\llbracket \varphi \rrbracket_{\mu}^{\mathfrak{A}} = \llbracket \hat{\mu}(\varphi) \rrbracket^{\mathfrak{A}}$ .

**Definition 12.16.** Let  $\mathcal{A}$  be a complete Boolean algebra, and let  $\mathfrak{A}$  be a structure over  $\mathcal{MT}_{\mathcal{A}}$  and  $\Theta$ . Given a set of formulas  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta_U)$ ,  $\varphi$  is said to be a *semantical consequence of  $\Gamma$  w.r.t.  $(\mathfrak{A}, \mathcal{MT}_{\mathcal{A}})$* , denoted by  $\Gamma \models_{(\mathfrak{A}, \mathcal{MT}_{\mathcal{A}})} \varphi$ , if the following holds: if  $\llbracket \gamma \rrbracket_{\mu}^{\mathfrak{A}} \in D$ , for every formula  $\gamma \in \Gamma$  and every assignment  $\mu$ , then  $\llbracket \varphi \rrbracket_{\mu}^{\mathfrak{A}} \in D$ , for every assignment  $\mu$ .

**Definition 12.17.** (Semantical consequence relation in  $\mathbf{QLFI1}_\circ$  w.r.t. twist structures) Let  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$  be a set of formulas. Then  $\varphi$  is said to be a *semantical consequence of  $\Gamma$  in  $\mathbf{QLFI1}_\circ$  w.r.t. first-order twist structures*, denoted by  $\Gamma \models_{\mathbf{QLFI1}_\circ} \varphi$ , if  $\Gamma \models_{(\mathfrak{A}, \mathcal{MT}_{\mathcal{A}})} \varphi$  for every  $(\mathfrak{A}, \mathcal{MT}_{\mathcal{A}})$ .

The soundness of  $\mathbf{QLFI1}_\circ$  w.r.t. first-order twist structures semantics can be easily obtained. The proof is analogous but much easier than the proof for  $\mathbf{QmbC}$  given in Theorem 7.5, given that valuations are no longer necessary.

**Proposition 12.18.**

- (i)  $\alpha, \alpha \rightarrow \beta \models_{\mathbf{QLFI1}_\circ} \beta$ ;
- (ii)  $\alpha \rightarrow \beta \models_{\mathbf{QLFI1}_\circ} \exists x \alpha \rightarrow \beta$ , if  $x$  is not free in  $\beta$ ;
- (iii)  $\alpha \rightarrow \beta \models_{\mathbf{QLFI1}_\circ} \alpha \rightarrow \forall x \beta$ , if  $x$  is not free in  $\alpha$ ;
- (iv)  $\models_{\mathbf{QLFI1}_\circ} \forall x \alpha \rightarrow \alpha[x/t]$ , if  $t$  is a term free for  $x$  in  $\alpha$ ;
- (v)  $\models_{\mathbf{QLFI1}_\circ} \alpha[x/t] \rightarrow \exists x \alpha$ , if  $t$  is a term free for  $x$  in  $\alpha$ ;
- (vi)  $\models_{\mathbf{QLFI1}_\circ} \neg \exists x \varphi \leftrightarrow \forall x \neg \varphi$ ;
- (vii)  $\models_{\mathbf{QLFI1}_\circ} \neg \forall x \varphi \leftrightarrow \exists x \neg \varphi$ .

*Proof.* The proofs for items (i)-(v) are easily obtained by adapting the corresponding proofs for  $\mathbf{QmbC}$  given in Proposition 7.3. Items (vi) and (vii) follow easily from the definitions.  $\square$

**Corollary 12.19.** *Let  $\mathcal{A}$  be a complete Boolean algebra, and let  $\mathfrak{A}$  be a structure over  $\mathcal{MT}_{\mathcal{A}}$  and  $\Theta$ . Then:*

- (1) *If  $\alpha$  is an instance of an axiom schema of  $\mathbf{QLFI1}_{\circ}$ , then  $\llbracket \alpha \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$ , for every assignment  $\mu$ .*
- (2) *If  $\alpha$  and  $\beta$  are formulas such that  $\llbracket \alpha \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$  and  $\llbracket \alpha \rightarrow \beta \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$  for every assignment  $\mu$ , then  $\llbracket \beta \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$  for every  $\mu$ .*
- (3) *If  $\alpha$  and  $\beta$  are formulas such that  $\llbracket \alpha \rightarrow \beta \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$  for every assignment  $\mu$ , and if  $x$  does not occur free in  $\beta$ , then  $\llbracket \exists x \alpha \rightarrow \beta \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$  for every  $\mu$ .*
- (4) *If  $\alpha$  and  $\beta$  are formulas such that  $\llbracket \alpha \rightarrow \beta \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$ , for every assignment  $\mu$ , and if  $x$  does not occur free in  $\alpha$ , then  $\llbracket \alpha \rightarrow \forall x \beta \rrbracket_{\mu}^{\mathfrak{A}} \in D_{\mathcal{A}}$  for every  $\mu$ .*

**Theorem 12.20.** (Soundness of  $\mathbf{QLFI1}_{\circ}$  w.r.t. first-order twist structures) *For every set  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ : if  $\Gamma \vdash_{\mathbf{QLFI1}_{\circ}} \varphi$ , then  $\Gamma \models_{\mathbf{QLFI1}_{\circ}} \varphi$ .*

*Proof.* By induction on the length  $n$  of a derivation of  $\varphi$  from  $\Gamma$  in  $\mathbf{QLFI1}_{\circ}$ , taking into account Corollary 12.19.  $\square$

## 13 Completeness of $\mathbf{QLFI1}_{\circ}$ w.r.t. twist structures

Now, the completeness of  $\mathbf{QLFI1}_{\circ}$  w.r.t. first-order twist structures semantics will be proved, by adapting the completeness proof for  $\mathbf{QmbC}$  given in Section 8.

The notion of *C-Henkin theory* in  $\mathbf{QLFI1}_{\circ}$  is analogous to the one for  $\mathbf{QmbC}$  given in Definition 8.1. The consequence relation  $\vdash_{\mathbf{QLFI1}_{\circ}}^C$  is the consequence relation of  $\mathbf{QLFI1}_{\circ}$  over the signature  $\Theta_C$ . The following result can be proved by adapting the proof for  $\mathbf{QmbC}$  (see Proposition 8.4):

**Proposition 13.1.** *Let  $\Gamma \cup \{\varphi\} \subseteq \text{Sent}(\Theta)$  such that  $\Gamma \not\vdash_{\mathbf{QLFI1}_{\circ}} \varphi$ . Then, there exists a set of sentences  $\Delta \subseteq \text{Sent}(\Theta)$  which is maximally non-trivial with respect to  $\varphi$  in  $\mathbf{QLFI1}_{\circ}$  (assuming that  $\vdash_{\mathbf{QLFI1}_{\circ}}$  is restricted to sentences) and such that  $\Gamma \subseteq \Delta$ .*

**Definition 13.2.** Let  $\Delta \subseteq \text{Sen}(\Theta)$  be a non-trivial theory in  $\mathbf{QLFI1}_{\circ}$ . Let  $\equiv_{\Delta}^1 \subseteq \text{Sen}(\Theta)^2$  be the relation in  $\text{Sen}(\Theta)$  given by:  $\alpha \equiv_{\Delta}^1 \beta$  iff  $\Delta \vdash_{\mathbf{QLFI1}_{\circ}} \alpha \leftrightarrow \beta$ .

As in the case of  $\mathbf{QmbC}$ ,  $\equiv_{\Delta}^1$  is an equivalence relation. The equivalence class of a sentence  $\alpha$  w.r.t.  $\equiv_{\Delta}^1$  will be denoted by  $|\alpha|_{\Delta}$ . By adapting the proof of Proposition 8.6 it is easy to prove the following:

**Proposition 13.3.** *The structure  $\mathcal{A}_{\Delta} \stackrel{\text{def}}{=} \langle A_{\Delta}, \bar{\wedge}, \bar{\vee}, \bar{\rightarrow}, 0_{\Delta}, 1_{\Delta} \rangle$  is a Boolean algebra with the following operations:  $|\alpha|_{\Delta} \bar{\#} |\beta|_{\Delta} \stackrel{\text{def}}{=} |\alpha \# \beta|_{\Delta}$  for any  $\# \in \{\wedge, \vee, \rightarrow\}$ ,  $0_{\Delta} \stackrel{\text{def}}{=} |\varphi \wedge (\neg\varphi \wedge \circ\varphi)|_{\Delta}$  and  $1_{\Delta} \stackrel{\text{def}}{=} |\varphi \vee \neg\varphi|_{\Delta}$ , for any sentence  $\varphi$ .*

It is worth noting that  $\sim|\alpha|_{\Delta} \stackrel{\text{def}}{=} |\sim\alpha|_{\Delta}$  is the Boolean complement of  $|\alpha|_{\Delta}$  in  $\mathcal{A}_{\Delta}$ .<sup>15</sup> As it was done with  $\mathbf{QmbC}$ , the construction of the canonical model for  $\mathbf{QLFI1}_{\circ}$  w.r.t.  $\Delta$  requires the use of the MacNeille-Tarski completion of the Boolean algebra  $\mathcal{A}_{\Delta}$ . Then, consider the following:

**Definition 13.4.** Let  $(C\mathcal{A}_{\Delta}, *)$  be the MacNeille-Tarski completion of  $\mathcal{A}_{\Delta}$ . The twist structure for  $\mathbf{LFI1}_{\circ}$  over  $C\mathcal{A}_{\Delta}$  will be denoted by  $\mathcal{T}(\Delta)$ , and its domain will be denoted by  $T(\Delta)$ . The associated logical matrix will be denoted by  $\mathcal{MT}(\Delta) \stackrel{\text{def}}{=} (\mathcal{T}(\Delta), D_{\Delta})$ .

<sup>15</sup>Observe that the occurrence of  $\sim$  on the right-hand side of the definition denotes a classical negation definable in  $\mathbf{LFI1}_{\circ}$ , as explained in Remark 8.2.

**Remark 13.5.** It is worth noting that  $(*(|\alpha|_\Delta), *(|\beta|_\Delta)) \in D_\Delta$  iff  $\Delta \vdash_{\mathbf{QLFI1}_\circ} \alpha$ .

**Definition 13.6.** (Canonical Structure) Let  $\Theta$  be a signature with some individual constant. Let  $\Delta \subseteq \text{Sen}(\Theta)$  be non-trivial in  $\mathbf{QLFI1}_\circ$ , let  $\mathcal{MT}(\Delta)$  be the matrix as in Definition 13.4, and let  $U \stackrel{\text{def}}{=} \text{CTer}(\Theta)$ . The *canonical  $\mathbf{QLFI1}_\circ$ -structure induced by  $\Delta$*  is the structure  $\mathfrak{A}_\Delta = \langle U, I_{\mathfrak{A}_\Delta} \rangle$  over  $\mathcal{MT}(\Delta)$  and  $\Theta$  such that:

- $c^{\mathfrak{A}_\Delta} = c$ , for every individual constant  $c \in \mathcal{C}$ ;
- $f^{\mathfrak{A}_\Delta} : U^n \rightarrow U$  is such that  $f^{\mathfrak{A}_\Delta}(t_1, \dots, t_n) = f(t_1, \dots, t_n)$ , for every function symbol  $f$  of arity  $n$ ;
- $P^{\mathfrak{A}_\Delta}(t_1, \dots, t_n) = (*(|\varphi|_\Delta), *(|\neg\varphi|_\Delta))$  with  $\varphi = P(t_1, \dots, t_n)$ , for every predicate symbol  $P$  of arity  $n$ .

Note that  $|\varphi|_\Delta \bar{\vee} |\neg\varphi|_\Delta = |\varphi \vee \neg\varphi|_\Delta = 1$  and so  $*(|\varphi|_\Delta) \vee *(|\neg\varphi|_\Delta) = *(|\varphi \vee \neg\varphi|_\Delta) = 1$ . Hence,  $P^{\mathfrak{A}_\Delta}(t_1, \dots, t_n) \in T(\Delta)$  and so  $\mathfrak{A}_\Delta$  is indeed a structure over  $\mathcal{MT}(\Delta)$  and  $\Theta$ .

Let  $(\cdot)^\triangleright : (\text{Ter}(\Theta_U) \cup \text{For}(\Theta_U)) \rightarrow (\text{Ter}(\Theta) \cup \text{For}(\Theta))$  be the function introduced in Definition 8.9 such that  $(s)^\triangleright$  is obtained from  $s$  by substituting every occurrence of a constant  $\bar{t}$  by the term  $t$  itself. Clearly  $(t)^\triangleright = \llbracket t \rrbracket^{\widehat{\mathfrak{A}_\Delta}}$  for every  $t \in \text{CTer}(\Theta_U)$ . By adapting the proof of Lemma 8.11 it follows:

**Lemma 13.7.** *Let  $\Delta \subseteq \text{Sen}(\Theta)$  be a set of sentences over a signature  $\Theta$  such that  $\Delta$  is a C-Henkin theory in  $\mathbf{QLFI1}_\circ$  for a nonempty set  $C$  of individual constants of  $\Theta$ , and  $\Delta$  is maximally non-trivial with respect to  $\varphi$  in  $\mathbf{QLFI1}_\circ$ , for some sentence  $\varphi$ . Then, for every formula  $\psi(x)$  in which  $x$  is the unique variable (possibly) occurring free, it holds:*

- (1)  $|\forall x\psi|_\Delta = \bigwedge_{\mathcal{A}_\Delta} \{|\psi[x/t]|_\Delta : t \in \text{CTer}(\Theta)\}$ , where  $\bigwedge_{\mathcal{A}_\Delta}$  denotes an existing infimum in the Boolean algebra  $\mathcal{A}_\Delta$ ;
- (2)  $|\exists x\psi|_\Delta = \bigvee_{\mathcal{A}_\Delta} \{|\psi[x/t]|_\Delta : t \in \text{CTer}(\Theta)\}$ , where  $\bigvee_{\mathcal{A}_\Delta}$  denotes an existing supremum in the Boolean algebra  $\mathcal{A}_\Delta$ .

**Proposition 13.8.** *Let  $\Delta \subseteq \text{Sen}(\Theta)$  be as in Lemma 13.7. Then, the interpretation map  $\llbracket \cdot \rrbracket^{\mathfrak{A}_\Delta} : \text{Sen}(\Theta_U) \rightarrow T(\Delta)$  is such that  $\llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} = (*(|\psi|_\Delta), *(|\neg\psi|_\Delta))$  for every sentence  $\psi$ . Moreover,  $\llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} \in D_\Delta$  iff  $\Delta \vdash_{\mathbf{QLFI1}_\circ} (\psi)^\triangleright$ . In particular,  $\llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} \in D_\Delta$  iff  $\Delta \vdash_{\mathbf{QLFI1}_\circ} \psi$  for every  $\psi \in \text{Sen}(\Theta)$ .*

*Proof.* The proof is done by induction on the complexity of the sentence  $\psi$  in  $\text{Sen}(\Theta_U)$ .

If  $\psi = P(t_1, \dots, t_n)$  is atomic then, by using Definition 12.14, the fact that  $\llbracket t \rrbracket^{\widehat{\mathfrak{A}_\Delta}} = (t)^\triangleright$  for every  $t \in \text{CTer}(\Theta_U)$ , and Definition 13.6, we have:

$$\begin{aligned} \llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} &= P^{\mathfrak{A}_\Delta}(\llbracket t_1 \rrbracket^{\widehat{\mathfrak{A}_\Delta}}, \dots, \llbracket t_n \rrbracket^{\widehat{\mathfrak{A}_\Delta}}) = P^{\mathfrak{A}_\Delta}((t_1)^\triangleright, \dots, (t_n)^\triangleright) \\ &= (*(|\psi|_\Delta), *(|\neg\psi|_\Delta)). \end{aligned}$$

If  $\psi = \neg\beta$  then, by Definition 12.14 and by induction hypothesis,

$$\llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} = \bar{\sim} \llbracket \beta \rrbracket^{\mathfrak{A}_\Delta} = \bar{\sim} (*(|(\beta)^\triangleright|_\Delta), *(|(\neg\beta)^\triangleright|_\Delta)) = (*(|(\neg\beta)^\triangleright|_\Delta), *(|(\beta)^\triangleright|_\Delta)).$$

But  $|\beta|_\Delta^\triangleright = |(\neg\neg\beta)^\triangleright|_\Delta$ , by (**dneg**). Hence,  $\llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} = (*(|\psi|_\Delta), *(|\neg\psi|_\Delta))$ .

If  $\psi = \circ\beta$  then, by Definition 12.14, by induction hypothesis, the definition of the operations in  $\mathcal{A}_\Delta$  and the fact that  $*$  is a homomorphism of Boolean algebras,

$$\begin{aligned} \llbracket \psi \rrbracket^{\mathfrak{A}_\Delta} &= \bar{\circ} \llbracket \beta \rrbracket^{\mathfrak{A}_\Delta} = \bar{\circ} (*(|(\beta)^\triangleright|_\Delta), *(|(\neg\beta)^\triangleright|_\Delta)) \\ &= (*(|(\sim(\beta \wedge \neg\beta))^\triangleright|_\Delta), *(|(\beta \wedge \neg\beta)^\triangleright|_\Delta)). \end{aligned}$$

But  $|(\sim(\beta \wedge \neg\beta))^\triangleright|_\Delta = |(\circ\beta)^\triangleright|_\Delta$  and  $|(\beta \wedge \neg\beta)^\triangleright|_\Delta = |(\neg\circ\beta)^\triangleright|_\Delta$ , by Remark 12.4. Hence,  $\llbracket\psi\rrbracket^{\mathfrak{A}_\Delta} = (*(|(\psi)^\triangleright|_\Delta), *(|(\neg\psi)^\triangleright|_\Delta))$ .

If  $\psi = \alpha\#\beta$  for  $\# \in \{\wedge, \vee, \rightarrow\}$ , the proof is analogous, but now axioms **(neg $\vee$ )**, **(neg $\wedge$ )** and **(neg $\rightarrow$ )** are required.

If  $\psi = \forall x\beta$  then, by Lemma 13.7 and using that  $U = CTer(\Theta)$ ,  $|\forall x\beta|_\Delta = \bigwedge_{\mathcal{A}_\Delta} \{\beta[x/t]|_\Delta : t \in U\}$  and so  $*(|\forall x\beta|_\Delta) = \bigwedge_{C\mathcal{A}_\Delta} \{*(|\beta[x/t]|_\Delta) : t \in U\}$ . Analogously,  $*(|\exists x\beta|_\Delta) = \bigvee_{C\mathcal{A}_\Delta} \{*(|\beta[x/t]|_\Delta) : t \in U\}$ . Then, by Definition 12.14, by induction hypothesis and by axiom **(Ax $\neg$ )**:

$$\begin{aligned} \llbracket\forall x\beta\rrbracket^{\mathfrak{A}_\Delta} &= (\bigwedge_{t \in U} (\llbracket\beta[x/t]\rrbracket^{\mathfrak{A}_\Delta})_1, \bigvee_{t \in U} (\llbracket\beta[x/t]\rrbracket^{\mathfrak{A}_\Delta})_2) \\ &= (\bigwedge_{t \in U} *(|\beta[x/t]|_\Delta), \bigvee_{t \in U} *(|\neg\beta[x/t]|_\Delta)) \\ &= (*(|\forall x\beta|_\Delta), *(|\exists x\neg\beta|_\Delta)) = (*(|\forall x\beta|_\Delta), *(|\neg\forall x\beta|_\Delta)). \end{aligned}$$

Hence,  $\llbracket\psi\rrbracket^{\mathfrak{A}_\Delta} = (*(|(\psi)^\triangleright|_\Delta), *(|(\neg\psi)^\triangleright|_\Delta))$ .

If  $\psi = \exists x\beta$ , the proof is analogous to the previous case.

This shows that  $\llbracket\psi\rrbracket^{\mathfrak{A}_\Delta} = (*(|(\psi)^\triangleright|_\Delta), *(|(\neg\psi)^\triangleright|_\Delta))$  for every sentence  $\psi$ . The rest of the proof follows by Remark 13.5.  $\square$

**Theorem 13.9.** (Completeness of **QLFI $_o$**  restricted to sentences w.r.t. first-order twist structures) *If  $\Gamma \models_{\mathbf{QLFI}_o} \varphi$  then  $\Gamma \vdash_{\mathbf{QLFI}_o} \varphi$ , for every  $\Gamma \cup \{\varphi\} \subseteq Sen(\Theta)$ .*

*Proof.* Suppose that  $\Gamma \cup \{\varphi\} \subseteq Sen(\Theta)$  is such that  $\Gamma \not\vdash_{\mathbf{QLFI}_o} \varphi$ . By adapting Theorem 7.5.3 in [7] to **QLFI $_o$** ,<sup>16</sup> there exists a  $C$ -Henkin theory  $\Delta^H$  over  $\Theta_C$  in **QLFI $_o$**  for some nonempty set  $C$  of new individual constants such that  $\Gamma \subseteq \Delta^H$  and, in addition:  $\Gamma \vdash_{\mathbf{QLFI}_o} \alpha$  iff  $\Delta^H \vdash_{\mathbf{QLFI}_o}^C \alpha$ , for every  $\alpha \in Sen(\Theta)$ . In consequence,  $\Delta^H \not\vdash_{\mathbf{QLFI}_o}^C \varphi$ . Because of Proposition 13.1, there is a set of sentences  $\overline{\Delta^H}$  in  $\Theta_C$  containing  $\Delta^H$  which is maximally non-trivial with respect to  $\varphi$  in **QLFI $_o$**  (restricted to  $Sen(\Theta_C)$ ), and such that  $\overline{\Delta^H}$  is also a  $C$ -Henkin theory over  $\Theta_C$  in **QLFI $_o$** . Consider now  $\mathcal{MT}(\overline{\Delta^H})$  and  $\mathfrak{A}_{\overline{\Delta^H}}$  as in Definitions 13.4 and 13.6, respectively. By Proposition 13.8,  $\llbracket\alpha\rrbracket^{\mathfrak{A}_{\overline{\Delta^H}}} \in D_{\overline{\Delta^H}}$  iff  $\overline{\Delta^H} \vdash_{\mathbf{QLFI}_o}^C \alpha$ , for every  $\alpha$  in  $Sen(\Theta_C)$ . But then  $\llbracket\gamma\rrbracket^{\mathfrak{A}_{\overline{\Delta^H}}} \in D_{\overline{\Delta^H}}$  for every  $\gamma \in \Gamma$  and  $\llbracket\varphi\rrbracket^{\mathfrak{A}_{\overline{\Delta^H}}} \notin D_{\overline{\Delta^H}}$ . Now, let  $\mathfrak{A}$  the reduct of  $\mathfrak{A}_{\overline{\Delta^H}}$  to  $\Theta$ . Hence,  $\mathfrak{A}$  is a structure over  $\mathcal{MT}(\overline{\Delta^H})$  and  $\Theta$  such that  $\llbracket\gamma\rrbracket^{\mathfrak{A}} \in D_{\overline{\Delta^H}}$  for every  $\gamma \in \Gamma$  but  $\llbracket\varphi\rrbracket^{\mathfrak{A}} \notin D_{\overline{\Delta^H}}$ . This means that  $\Gamma \not\vdash_{\mathbf{QLFI}_o} \varphi$ .  $\square$

**Corollary 13.10.** (Completeness of **QmbC** w.r.t. first-order twist structures) *Let  $\Gamma \cup \{\varphi\} \subseteq For(\Theta)$ . If  $\Gamma \models_{\mathbf{QLFI}_o} \varphi$  then  $\Gamma \vdash_{\mathbf{QLFI}_o} \varphi$ .*

## 14 Completeness of **QLFI $_o$** w.r.t. structures over $\mathcal{M}_{LFI1}$

In Remark 12.8 it was observed that  $\mathcal{T}_{\mathcal{A}_2}$ , the twist structure for **LFI $_o$**  defined over the two-element Boolean algebra  $\mathcal{A}_2$ , coincides (up to names) with the 3-valued algebra  $\mathcal{A}_{LFI1}$  underlying the matrix  $\mathcal{M}_{LFI1}$  and, moreover,  $\mathcal{MT}_{\mathcal{A}_2}$  coincides with the 3-valued characteristic matrix  $\mathcal{M}_{LFI1}$  of **LFI $_o$** . Recall that 1,  $\frac{1}{2}$  and 0 are identified with (1, 0), (1, 1) and (0, 1), respectively. Let  $\mathfrak{A}$  be a **QLFI $_o$** -structure over  $\mathcal{A}_2$ . If  $\varphi$  is a formula

<sup>16</sup>As observed in the proof of Theorem 8.13 above, Theorem 7.5.3 in [7] also holds for the definition of  $C$ -Henkin theory adopted in this paper.

in which  $x$  is the unique variable (possibly) occurring free, let  $X = \{\llbracket \varphi[x/\bar{a}] \rrbracket^{\mathfrak{A}} : a \in U\}$ . Then:

$$\llbracket \forall x \varphi \rrbracket^{\mathfrak{A}} = \begin{cases} 1 & \text{if } X = \{1\} \\ \frac{1}{2} & \text{if } X \subseteq \{1, \frac{1}{2}\} \text{ and } \frac{1}{2} \in X \\ 0 & \text{if } 0 \in X \end{cases} \quad \llbracket \exists x \varphi \rrbracket^{\mathfrak{A}} = \begin{cases} 1 & \text{if } 1 \in X \\ \frac{1}{2} & \text{if } X \cap \{1, \frac{1}{2}\} = \{\frac{1}{2}\} \\ 0 & \text{if } X = \{0\} \end{cases}$$

In Section 9 it was obtained a characterization of **QmbC** in terms of swap structures over the 5-element characteristic Nmatrix of **mbC**, which coincides with the one given in [6]. That result can be easily adapted to **QLFI1<sub>o</sub>**, by proving that **QLFI1<sub>o</sub>** can be characterized by first-order structures defined over  $\mathcal{M}_{LFI1}$ . Indeed, it is possible to adapt Theorem 9.5 to **QLFI1<sub>o</sub>**, taking into account that the bivaluations for **QLFI1<sub>o</sub>** satisfy additional clauses, see [7, Definition 7.9.16]. This lead us to the following result, in view of Remark 12.8 (details of the proof will be omitted):

**Theorem 14.1.** (Adequacy of **QLFI1<sub>o</sub>** w.r.t. first-order structures over  $\mathcal{M}_{LFI1}$ ) *For every set  $\Gamma \cup \{\varphi\} \subseteq \text{For}(\Theta)$ :  $\Gamma \vdash_{\mathbf{QLFI1}_o} \varphi$  iff  $\Gamma \models_{(\mathfrak{A}, \mathcal{M}_{LFI1})} \varphi$  for every structure  $\mathfrak{A}$  over  $\Theta$  and  $\mathcal{M}_{LFI1}$ .*

The latter result is a variant (up to language) of the adequacy theorem of first-order **J3** w.r.t. first-order structures given in [18] (see also [17, 19, 20]). Indeed, the semantics in terms of first-order structures over  $\mathcal{M}_{LFI1}$  is equivalent to the 3-valued first-order structures proposed by D'Ottaviano in [18] for a quantified version of **J3**, given that **LF1<sub>o</sub>** is equivalent, up to language, to **J3**. This shows that the twist-structures semantics for **QLFI1<sub>o</sub>** constitutes a generalization, to any complete Boolean algebra, of the above mentioned semantics for first-order **J3**.

The extension of **QLFI1<sub>o</sub>** with standard equality is straightforward, taking into account the construction for **QmbC<sup>≈</sup>** presented in Section 10. It is worth noting that, when restricted to structures over  $\mathcal{M}_{LFI1}$ , there are differences with D'Ottaviano's approach to first-order **J3** with equality. Indeed, she assumes that the equality must be classical, that is, every formula  $\circ(t_1 \approx t_2)$  is valid in her system, contrary to what happens in **QLFI1<sub>o</sub>** with equality.

## 15 Final remarks

In this paper, the semantical frameworks for **QmbC** already proposed in the literature were extended to a vast class of models based on the non-deterministic algebras known as swap structures. Indeed, the Nmatrix semantics proposed in [6] and the semantics given by interpretations (i.e., standard Tarskian structures plus bivaluations) considered in [10] and [7] coincide, and are particular cases of the swap structures semantics introduced here, as it was shown along this paper.

The advantage of considering models based on a class of swap structures instead of 'classical' models based on a finite Nmatrix (as done in [6]) is that this enlarged class of models allows us to consider applications to another fields such, for instance, algebraic logic (as done in [14]) or paraconsistent set theory. Concerning the latter, the Boolean valued models for set theory could be generalized to this setting, obtaining so swap structures models for several paraconsistent set theories based on **LFIs**, along the lines of the twist-valued models introduced in [8].

Two important model-theoretic results for **QmbC** (and some of its axiomatic extensions) were obtained by Ferguson in [21]: Łoś’ ultraproducts theorem, and a suitable version of the Keisler-Shelah isomorphism theorem, which states that two **QmbC**-models are strongly elementarily equivalent iff there exists an ultrafilter  $\mathcal{U}$  such that the corresponding ultrapowers over  $\mathcal{U}$  are strongly isomorphic. The notions of strong elementary equivalence and strong isomorphism were introduced in [21], as well as an adaptation of the method of atomization introduced by Skolem, which was used in order to prove the Keisler-Shelah theorem for quantified **LFI**s. It would be interesting to adapt Ferguson’s notions and constructions to the present semantical framework for quantified **LFI**s.

In other line of research, it would be interesting to extend the techniques developed in [3] for generating cut-free Gentzen-type calculi for propositional **LFI**s from Nmatrix semantics to the first-order framework described here.

**Acknowledgements:** This paper is the full version of the extended abstract [15]. Coniglio was financially supported by an individual research grant from CNPq, Brazil (308524/2014-4). Figallo-Orellano acknowledges financial support from a post-doctoral grant from FAPESP, Brazil (2016/21928-0). Golzio was financially supported by a post-doctoral grant from CNPq, Brazil (150064/2018-7) and by a post-doctoral grant from FAPESP, Brazil (2019/08442-9).

## References

- [1] Avron, A. (2005). Non-deterministic matrices and modular semantics of rules. In J.-Y. Béziau (Ed.), *Logica Universalis* (pp. 149–167). Birkhäuser Verlag.
- [2] Avron, A. (2007). Non-deterministic Semantics for Logics with a Consistency Operator. *International Journal of Approximate Reasoning*, 45(2), 271–287.
- [3] Avron, A., Konikowska, B. & Zamansky, A. (2013). Cut-free sequent calculi for C-systems with generalized finite-valued semantics. *Journal of Logic and Computation*, 23(3), 517–540.
- [4] Avron, A. & Lev, I. (2001). Canonical propositional Gentzen-type systems. In R. Gore, A. Leitsch, & T. Nipkow (Eds.) *Proceedings of the 1st International Joint Conference on Automated Reasoning (IJCAR 2001)* (pp. 529–544). LNAI 2083, Springer Verlag.
- [5] Avron, A. & Lev, I. (2005). Non-deterministic multi-valued structures. *Journal of Logic and Computation*, 15(3), 241–261.
- [6] Avron, A., & Zamansky, A. (2007). Many-valued non-deterministic semantics for first-order Logics of Formal (In)consistency. In S. Aguzzoli, A. Ciabattoni, B. Gerla, C. Manara, & V. Marra (Eds.) *Algebraic and Proof-theoretic Aspects of Non-classical Logics* (pp. 1–24). LNAI 4460, Springer.
- [7] Carnielli, W. A., & Coniglio, M. E. (2016) *Paraconsistent Logic: Consistency, Contradiction and Negation*, vol. 40, Logic, Epistemology, and the Unity of Science, Basel, Switzerland: Springer International Publishing.
- [8] Carnielli, W. A., & Coniglio, M. E. (2019) Twist-Valued Models for Three-valued Paraconsistent Set Theory. *arXiv:1911.11833* [math.LO].

- [9] Carnielli, W. A., Coniglio, M. E., & Marcos, J. (2007) Logics of Formal Inconsistency. In D. M. Gabbay, & F. Guenther (Eds.) *Handbook of Philosophical Logic (2nd. edition)* (vol. 14, pp. 1–93). Springer.
- [10] Carnielli, W. A., Coniglio, M. E., Podiacki, R., & Rodrigues, T. (2014) On the way to a wider model theory: Completeness theorems for first-order logics of formal inconsistency. *The Review of Symbolic Logic*, 7(3), 548–578.
- [11] Carnielli, W. A., & Marcos, J. (2002) A taxonomy of C-systems. In W. A. Carnielli, M. E. Coniglio, & I. M. L. D’Ottaviano (Eds.) *Paraconsistency: The Logical Way to the Inconsistent: Lecture Notes in Pure and Applied Mathematics* (vol. 228, pp. 1–94). New York: Marcel Dekker.
- [12] Chang, C. C., & Keisler, H. J. (1990) *Model Theory*. Elsevier, Third edition.
- [13] Cignoli, R. (1986) The class of Kleene algebras satisfying an interpolation property and Nelson algebras. *Algebra Universalis*, 23, 262–292.
- [14] Coniglio, M. E., Figallo-Orellano, A., & Golzio, A. C. (2018) Non-deterministic algebraization of logics by swap structures, *Logic Journal of the IGPL*. <https://doi.org/10.1093/jigpal/jzy072>
- [15] Coniglio, M. E., Figallo-Orellano, A., & Golzio, A. C. (2019) First-order swap structures semantics for QmbC (extended abstract). In N. Bezhanishvili & Y. Venema, (Eds.) *SYSMICS 2019: Syntax Meets Semantics - Book of Abstracts* (pp. 62–65). Institute for Logic, Language and Computation, University of Amsterdam.
- [16] Cintula, P., & Noguera, C. (2015) A Henkin-style proof of completeness for first-order algebraizable logics. *The Journal of Symbolic Logic*, 80(1), 341–358.
- [17] D’Ottaviano, I. M. L. (1982) *Sobre uma Teoria de Modelos Trivalente (On a three-valued model theory, in Portuguese)*. PhD thesis, IMECC, University of Campinas.
- [18] D’Ottaviano, I. M. L. (1985) The completeness and compactness of a three-valued first-order logic. *Revista Colombiana de Matemáticas*, XIX(1-2), 77–94.
- [19] D’Ottaviano, I. M. L. (1985) The model extension theorems for J3-theories. In C. A. Di Prisco (Ed.) *Methods in Mathematical Logic. Proceedings of the 6th Latin American Symposium on Mathematical Logic: Lecture Notes in Mathematics* (vol. 1130, pages 157–173), Berlin: Springer-Verlag.
- [20] D’Ottaviano, I. M. L. (1987) Definability and quantifier elimination for J3-theories. *Studia Logica*, 46(1), 37–54.
- [21] Ferguson, T. M. (2018) The Keisler-Shelah theorem for QmbC through semantical atomization. *Logic Journal of the IGPL*. <https://doi.org/10.1093/jigpal/jzy067>
- [22] Fidel, M. M. (1978) An algebraic study of a propositional system of Nelson. In A. I. Arruda, N. C. A. da Costa, & R. Chuaqui (Eds.) *Mathematical Logic. Proceedings of the First Brazilian Conference on Mathematical Logic, Campinas 1977: Lecture Notes in Pure and Applied Mathematics* (vol. 39, pp. 99–117). Marcel Dekker.
- [23] Givant, S., & Halmos, P. (2009) *Introduction to Boolean Algebras*. Springer.



- [24] Henkin, L. (1950) An algebraic characterization of quantifiers. *Fundamenta Mathematicae*, 37(1), 63–74.
- [25] Kalman, J. A. (1958) Lattices with involution. *Transactions of the American Mathematical Society*, 87(2), 485–491.
- [26] Mendelson, E. (1997) *Introduction to Mathematical Logic*, Fourth Edition. Chapman and Hall/CRC.
- [27] Mostowski, A. (1948) Proofs of non-deducibility in intuitionistic functional calculus. *The Journal of Symbolic Logic*, 13(4), 204–207.
- [28] Rasiowa, H. (1974) *An Algebraic Approach to Non-Classical Logics*. North Holland.
- [29] Rasiowa, H. & Sikorski, R. (1963) *The Mathematics of Metamathematics*. Państwowe Wydawnictwo Naukowe, Warszaw.
- [30] Vakarelov, D. (1977) Notes on N-lattices and constructive logic with strong negation. *Studia Logica*, 36(1-2), 109–125.