

Generalized Löb's Theorem. Strong Reflection Principles and Large Cardinal Axioms

J. Foukzon¹, E. R. Men'kova²

¹Israel Institute of Technology, Haifa, Israel

²Lomonosov Moscow State University, Moscow, Russia

Email: jaykovfoukzon@list.ru, E_Menkova@mail.ru

Received February 9, 2013; revised March 13, 2013; accepted April 14, 2013

Copyright © 2013 J. Foukzon, E. R. Men'kova. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

ABSTRACT

In this article, a possible generalization of the Löb's theorem is considered. Main result is: let κ be an inaccessible cardinal, then $\neg \text{Con}(ZFC + \exists \kappa)$.

Keywords: Löb's Theorem; Second Gödel Theorem; Consistency; Formal System; Uniform Reflection Principles; ω -Model of ZFC; Standard Model of ZFC; Inaccessible Cardinal

1. Introduction

Let Th be some fixed, but unspecified, consistent formal theory.

Theorem 1 [1]. (Löb's Theorem).

If $Th \vdash \exists x \text{Prov}_{Th}(x, \bar{n}) \rightarrow \phi_n$ where x is the Gödel number of the proof of the formula with Gödel number n , and \bar{n} is the numeral of the Gödel number of the formula ϕ_n , then $Th \vdash \phi_n$. Taking into account the second Gödel theorem it is easy to be able to prove

$\exists x \text{Prov}_{Th}(x, \bar{n}) \rightarrow \phi_n$, for disprovable (refutable) and undecidable formulas ϕ_n . Thus summarized, Löb's theorem says that for refutable or undecidable formula ϕ , the intuition "if exists proof of ϕ then ϕ " is fails.

Definition 1. Let M_ω^{Th} be an ω -model of the Th . We said that, $Th^\#$ is a nice theory over Th or a nice extension of the Th iff:

- 1) $Th^\#$ contains Th ;
- 2) Let Φ be any closed formula, then

$$\left[Th \vdash \text{Pr}_{Th}([\Phi]^c) \right] \& \left[M_\omega^{Th} \models \Phi \right]$$

implies $Th^\# \vdash \Phi$.

Definition 2. We said that, $Th^\#$ is a maximally nice theory over Th or a maximally nice extension of the Th iff $Th^\#$ is consistent and for any consistent nice extension Th' of the Th : $\text{Ded}(Th^\#) \subseteq \text{Ded}(Th')$ implies

$$\text{Ded}(Th^\#) = \text{Ded}(Th').$$

Theorem 2. (Generalized Löb's Theorem). Assume that 1) $\text{Con}(Th)$ and 2) Th has an ω -model M_ω^{Th} . Then

theory Th can be extended to a maximally consistent nice theory $Th^\#$.

2. Preliminaries

Let Th be some fixed, but unspecified, consistent formal theory. For later convenience, we assume that the encoding is done in some fixed formal theory S and that Th contains S . We do not specify S —it is usually taken to be a formal system of arithmetic, although a weak set theory is often more convenient. The sense in which S is contained in Th is better exemplified than explained: If S is a formal system of arithmetic and Th is, say, ZFC , then Th contains S in the sense that there is a well-known embedding, or interpretation, of S in Th . Since encoding is to take place in S , it will have to have a large supply of constants and closed terms to be used as codes. (e.g. in formal arithmetic, one has $\bar{0}, \bar{1}, \dots$) S will also have certain function symbols to be described shortly. To each formula, Φ , of the language of Th is assigned a closed term, $[\Phi]^c$, called the code of Φ . [N. B. If $\Phi(x)$ is a formula with a free variable x , then $[\Phi(x)]^c$ is a closed term encoding the formula $\Phi(x)$ with x viewed as a syntactic object and not as a parameter.] Corresponding to the logical connectives and quantifiers are function symbols, $\text{neg}(\cdot), \text{imp}(\cdot)$, etc., such that, for all formulae $\Phi, \Psi : S \vdash \neg([\Phi]^c)$

$$= [\neg\Phi]^c, S \vdash \text{imp}([\Phi]^c, [\Psi]^c) = [\Phi \rightarrow \Psi]^c \quad \text{etc.}$$

Of particular importance is the substitution operator, represented by the function symbol $\text{sub}(\cdot, \cdot)$. For formulae $\Phi(x)$, terms t with codes $[t]^c$:

$$S|- \text{sub}\left(\left[\Phi(x)\right]^c, [t]^c\right) = \left[\Phi(t)\right]^c. \quad (2.1)$$

Iteration of the substitution operator sub allows one to define function symbols $\text{sub}_3, \text{sub}_4, \dots, \text{sub}_n$ such that

$$S|- \text{sub}_n\left(\left[\Phi(x_1, x_2, \dots, x_n)\right]^c, [t_1]^c, [t_2]^c, \dots, [t_n]^c\right) = \left[\Phi(t_1, t_2, \dots, t_n)\right]^c \quad (2.2)$$

It well known [2,3] that one can also encode derivations and have a binary relation $\text{Prov}_{Th}(x, y)$ (read “ x proves y ” or “ x is a proof of y ”) such that for closed t_1, t_2 : $S|- \text{Prov}_{Th}(t_1, t_2)$ iff t_1 is the code of a derivation in Th of the formula with code t_2 . It follows that

$$Th \vdash \Phi \leftrightarrow S \vdash \text{Prov}_{Th}(t, [\Phi]^c) \quad (2.3)$$

for some closed term t . Thus one can define predicate $\text{Pr}_{Th}(y)$:

$$\text{Pr}_{Th}(y) \leftrightarrow \exists x \text{Prov}_{Th}(x, y), \quad (2.4)$$

and therefore one obtain a predicate asserting provability.

Remark 2.1. We note that is not always the case that [2,3]:

$$Th \vdash \Phi \leftrightarrow S \vdash \text{Pr}_{Th}([\Phi]^c). \quad (2.5)$$

It well known [3] that the above encoding can be carried out in such a way that the following important conditions $D1, D2$ and $D3$ are met for all sentences [2,3]:

$$\begin{aligned} D1. & Th \vdash \Phi \text{ implies } S \vdash \text{Pr}_{Th}([\Phi]^c), \\ D2. & S \vdash \text{Pr}_{Th}([\Phi]^c) \rightarrow \text{Pr}_{Th}\left(\left[\text{Pr}_{Th}([\Phi]^c)\right]^c\right), \\ D3. & S \vdash \text{Pr}_{Th}([\Phi]^c) \wedge \text{Pr}_{Th}([\Phi \rightarrow \Psi]^c) \\ & \rightarrow \text{Pr}_{Th}([\Psi]^c). \end{aligned} \quad (2.6)$$

Conditions $D1, D2$ and $D3$ are called the Derivability Conditions.

Assumption 2.1. We assume now that:

1) the language of Th consists of:

numerals $\bar{0}, \bar{1}, \dots$

countable set of the numerical variables: $\{v_0, v_1, \dots\}$

countable set F of the set variables:

$F = \{x, y, z, X, Y, Z, \mathfrak{R}, \dots\}$

countable set of the n -ary function symbols: f_0^n, f_1^n, \dots

countable set of the n -ary relation symbols: R_0^n, R_1^n, \dots

connectives: \neg, \rightarrow

quantifier: \forall .

2) Th contains ZFC

3) Th has an ω -model M_ω^{Th} .

Theorem 2.1. (Löb’s Theorem). Let be 1) $\text{Con}(Th)$ and 2) ϕ be closed. Then

$$Th \vdash \text{Pr}_{Th}([\phi]^c) \rightarrow \phi \text{ iff } Th \vdash \phi. \quad (2.7)$$

It well known that replacing the induction scheme in Peano arithmetic PA by the ω -rule with the meaning “if the formula $A(n)$ is provable for all n , then the formula $A(x)$ is provable”:

$$\frac{A(0), A(1), \dots, A(n), \dots}{\forall x A(x)}, \quad (2.8)$$

leads to complete and sound system PA_∞ where each true arithmetical statement is provable. S. Feferman showed that an equivalent formal system $Th^\#$ can be obtained by erecting on $Th = PA$ a transfinite progression of formal systems PA_α according to the following scheme

$$\begin{aligned} PA_0 &= PA \\ PA_{\alpha+1} &= PA_\alpha + \left\{ \forall x \text{Pr}_{PA_\alpha}([\hat{A}(x)]^c) \rightarrow \forall x A(x) \right\}, \\ PA_\lambda &= \bigcup_{\alpha < \lambda} PA_\alpha \end{aligned} \quad (2.9)$$

where $A(x)$ is a formula with one free variable and λ is a limit ordinal. Then $Th = \bigcup_{\alpha \in O} PA_\alpha$, O being Kleene’s system of ordinal notations, is equivalent to $Th^\# = PA_\infty$. It is easy to see that $Th^\# = PA^\#$, i.e. $Th^\#$ is a maximally nice extension of the PA .

3. Generalized Löb’s Theorem

Definition 3.1. An Th -wff Φ (well-formed formula Φ) is closed i.e., Φ is a Th -sentence iff it has no free variables; a wff Ψ is open if it has free variables. We’ll use the slang “ k -place open wff” to mean a wff with k distinct free variables. Given a model M^{Th} of the Th and a Th -sentence Φ , we assume known the meaning of $M \models \Phi$ —i.e. Φ is true in M^{Th} , (see for example [4-6]).

Definition 3.2. Let M_ω^{Th} be an ω -model of the Th . We shall say that, $Th^\#$ is a nice theory over Th or a nice extension of the Th iff:

1) $Th^\#$ contains Th ;

2) Let Φ be any closed formula, then

$$\left[Th \vdash \text{Pr}_{Th}([\Phi]^c) \right] \& \left[M_\omega^{Th} \models \Phi \right]$$

implies $Th^\# \vdash \Phi$.

Definition 3.3. We shall say that $Th^\#$ is a maximally nice theory over Th or a maximally nice extension of the Th iff $Th^\#$ is consistent and for any consistent nice extension Th' of the Th : $\text{Ded}(Th^\#) \subseteq \text{Ded}(Th')$ implies $\text{Ded}(Th^\#) = \text{Ded}(Th')$.

Lemma 3.1. Assume that: 1) $\text{Con}(Th)$; and 2) $Th \vdash \text{Pr}_{Th}([\Phi]^c)$, where Φ is a closed formula. Then $Th \not\vdash \text{Pr}_{Th}([\neg\Phi]^c)$.

Proof. Let $\text{Con}_{Th}(\Phi)$ be the formula

$$\begin{aligned} & \text{Con}_{Th}(\Phi) \\ & \triangleq \forall t_1 \forall t_2 \neg \left[\text{Prov}_{Th}(t_1, [\Phi]^c) \wedge \text{Prov}_{Th}(t_2, \text{neg}([\Phi]^c)) \right] \\ & \leftrightarrow \neg \exists t_1 \neg \exists t_2 \left[\text{Prov}_{Th}(t_1, [\Phi]^c) \wedge \text{Prov}_{Th}(t_2, \text{neg}([\Phi]^c)) \right] \end{aligned} \quad (3.1)$$

where t_1, t_2 is a closed term. We note that under canonical observation, one obtains

$Th + \text{Con}(Th) \vdash \text{Con}_{Th}(\Phi)$ for any closed wff Φ .

Suppose that $Th \vdash \text{Pr}_{Th}([\neg\Phi]^c)$, then assumption (ii) gives

$$Th \vdash \text{Pr}_{Th}([\Phi]^c) \wedge \text{Pr}_{Th}([\neg\Phi]^c). \quad (3.2)$$

From (3.1) and (3.2) one obtain

$$\exists t_1 \exists t_2 \left[\text{Prov}_{Th}(t_1, [\Phi]^c) \wedge \text{Prov}_{Th}(t_2, \text{neg}([\Phi]^c)) \right]. \quad (3.3)$$

But the Formula (3.3) contradicts the Formula (3.1).

Therefore: $Th \not\vdash \text{Pr}_{Th}([\neg\Phi]^c)$.

Lemma 3.2. Assume that: 1) $\text{Con}(Th)$; and 2)

$Th \vdash \text{Pr}_{Th}([\neg\Phi]^c)$, where Φ is a closed formula. Then $Th \not\vdash \text{Pr}_{Th}([\Phi]^c)$.

Theorem 3.1. [7,8]. (Generalized Löb's Theorem). Assume that: $\text{Con}(Th)$. Then theory Th can be extended to a maximally consistent nice theory $Th^\#$ over Th .

Proof. Let $\Phi_1 \dots \Phi_i \dots$ be an enumeration of all wff's of the theory Th (this can be achieved if the set of propositional variables can be enumerated). Define a chain $\mathcal{S} = \{Th_i \mid i \in \mathbb{N}\}$, $Th_1 = Th$ of consistent theories inductively as follows: assume that theory Th_i is defined.

1) Suppose that a statement (3.4) is satisfied

$$\begin{aligned} & Th \vdash \text{Pr}_{Th}([\Phi_i]^c) \text{ and} \\ & [Th_i \not\vdash \Phi_i] \& [M_\omega^{Th} \models \Phi_i] \end{aligned} \quad (3.4)$$

Then we define theory Th_{i+1} as follows

$$Th_{i+1} \triangleq Th_i \cup \{\Phi_i\}.$$

2) Suppose that a statement (3.5) is satisfied

$$\begin{aligned} & Th \vdash \text{Pr}_{Th}([\neg\Phi_i]^c) \text{ and} \\ & [Th_i \not\vdash \neg\Phi_i] \& [M_\omega^{Th} \models \neg\Phi_i] \end{aligned} \quad (3.5)$$

Then we define theory Th_{i+1} as follows:

$$Th_{i+1} \triangleq Th_i \cup \{\neg\Phi_i\}.$$

3) Suppose that a statement (3.6) is satisfied

$$Th \vdash \text{Pr}_{Th}([\Phi_i]^c) \text{ and } Th_i \vdash \Phi_i. \quad (3.6)$$

Then we define theory Th_{i+1} as follows:

$$Th_{i+1} \triangleq Th_i \cup \{\Phi_i\}.$$

4) Suppose that a statement (3.7) is satisfied

$$Th \vdash \text{Pr}_{Th}([\neg\Phi_i]^c) \text{ and } Th \vdash \neg\Phi_i. \quad (3.7)$$

Then we define theory Th_{i+1} as follows:

$$Th_{i+1} \triangleq Th_i.$$

We define now theory $Th^\#$ as follows:

$$Th^\# \triangleq \bigcup_{i \in \mathbb{N}} Th_i. \quad (3.8)$$

First, notice that each Th_i is consistent. This is done by induction on i and by Lemmas 3.1-3.2. By assumption, the case is true when $i=1$. Now, suppose Th_i is consistent. Then its deductive closure $\text{Ded}(Th_i)$ is also consistent. If a statement (3.6) is satisfied *i.e.*,

$Th \vdash \text{Pr}_{Th}([\Phi_i]^c)$ and $Th \vdash \Phi_i$, then clearly

$Th_{i+1} \triangleq Th_i \cup \{\Phi_i\}$ is consistent since it is a subset of closure $\text{Ded}(Th_i)$. If a statement (3.7) is satisfied, *i.e.*,

$Th \vdash \text{Pr}_{Th}([\neg\Phi_i]^c)$ and $Th \vdash \neg\Phi_i$, then clearly

$Th_{i+1} \triangleq Th_i \cup \{\neg\Phi_i\}$ is consistent since it is a subset of closure $\text{Ded}(Th_i)$.

Otherwise:

1) if a statement (3.4) is satisfied, *i.e.*

$Th_i \vdash \text{Pr}_{Th_i}([\Phi_i]^c)$ and $Th_i \not\vdash \Phi_i$, then clearly

$Th_{i+1} \triangleq Th_i \cup \{\Phi_i\}$ is consistent by Lemma 3.1 and by one of the standard properties of consistency: $\Delta \cup \{A\}$ is consistent iff $\Delta \not\vdash \neg A$;

2) if a statement (3.5) is satisfied, *i.e.*

$Th \vdash \text{Pr}_{Th}([\neg\Phi_i]^c)$ and $Th_i \not\vdash \neg\Phi_i$, then clearly

$Th_{i+1} \triangleq Th_i \cup \{\neg\Phi_i\}$ is consistent by Lemma 3.2 and by one of the standard properties of consistency: $\Delta \cup \{\neg A\}$ is consistent iff $\Delta \not\vdash A$.

Next, notice $\text{Ded}(Th^\#)$ is a maximally consistent nice extension of the set $\text{Ded}(Th)$. A set $\text{Ded}(Th^\#)$ is consistent because, by the standard Lemma 3.3 below, it is the union of a chain of consistent sets. To see that $\text{Ded}(Th^\#)$ is maximal, pick any wff Φ . Then Φ is some Φ_i in the enumerated list of all wff's. Therefore

for any Φ such that $Th \vdash \text{Pr}_{Th}([\Phi]^c)$ or $Th \vdash \text{Pr}_{Th}([\neg\Phi]^c)$, either $\Phi \in Th^\#$ or $\neg\Phi \in Th^\#$.

Since $\text{Ded}(Th_{i+1}) \subseteq \text{Ded}(Th^\#)$, we have $\Phi \in \text{Ded}(Th^\#)$ or $\neg\Phi \in \text{Ded}(Th^\#)$, which implies that $\text{Ded}(Th^\#)$ is maximally consistent nice extension of the $\text{Ded}(Th)$.

Lemma 3.3. The union of a chain $\wp = \{\Gamma_i \mid i \in \mathbb{N}\}$ of the consistent sets Γ_i , ordered by \subseteq , is consistent.

Definition 3.4. (a) Assume that a theory Th has an ω -model M_ω^{Th} and Φ is a Th -sentence. Let Φ_ω be a Th -sentence Φ with all quantifiers relativised to ω -model M_ω^{Th} ;

(b) Assume that a theory Th has a standard model SM^{Th} and Φ is a Th -sentence. Let Φ_{SM} be a Th -sentence Φ with all quantifiers relativized to a model SM^{Th} [9].

Remark 3.1. In some special cases we denote a sentence Φ_ω by a symbol $\Phi[M_\omega^{Th}]$ and we denote a sentence Φ_{SM} by symbol $\Phi[SM^{Th}]$ correspondingly.

Definition 3.5. (a) Assume that Th has an ω -model M_ω^{Th} . Let Th_ω be a theory Th relativized to a model M_ω^{Th} , that is, any Th_ω -sentence has a form Φ_ω for some Th -sentence Φ [9];

(b) Assume that Th has a standard model SM^{Th} . Let Th_{SM} be a theory Th relativized to a model SM^{Th} , that is, any Th_{SM} -sentence has a form Φ_{SM} for some Th -sentence Φ [9].

Remark 3.2. In some special cases we denote a theory Th_ω by symbol $Th[M_\omega^{Th}]$ and we denote a theory Th_{SM} by symbol $Th[SM^{Th}]$ correspondingly.

Theorem 3.2. (Strong Reflection Principle).

(i) Assume that: Th has an ω -model M_ω^{Th} . Then for any Th_ω -sentence Φ_ω

$$Th_\omega \vdash \text{Pr}_{Th_\omega}([\Phi_\omega]^c) \text{ iff } Th_\omega \vdash \Phi_\omega. \quad (3.9)$$

(ii) Assume that: Th has model M_{SM}^{Th} . Then for any Th_{SM} -sentence Φ_{SM}

$$Th_{SM} \vdash \text{Pr}_{Th_{SM}}([\Phi_{SM}]^c) \text{ iff } Th_{SM} \vdash \Phi_{SM}. \quad (3.10)$$

Proof. (i) The one direction is obvious. For the other, assume that

$$Th_\omega \vdash \text{Pr}_{Th_\omega}([\Phi_\omega]^c), Th_\omega \not\vdash \Phi_\omega, \quad (3.11)$$

and $Th_\omega \vdash \neg\Phi_\omega$. Then

$$Th_\omega \vdash \text{Pr}_{Th_\omega}([\neg\Phi_\omega]^c). \quad (3.12)$$

Note that $\text{Con}(Th_\omega)$ holds since $\exists M_\omega^{Th}$. Let Con_{Th_ω} be the formula

$$\begin{aligned} \text{Con}_{Th_\omega} &\leftrightarrow \forall t_1 \forall t_2 \forall t_3 (t_3 = [\Phi_\omega]^c) \\ &\neg \left[\text{Prov}_{Th_\omega}(t_1, [\Phi_\omega]^c) \wedge \text{Prov}_{Th_\omega}(t_2, \text{neg}([\Phi_\omega]^c)) \right] \\ &\leftrightarrow \neg \exists t_1 \neg \exists t_2 \neg \exists t_3 (t_3 = [\Phi_\omega]^c) \\ &\times \left[\text{Prov}_{Th_\omega}(t_1, [\Phi_\omega]^c) \wedge \text{Prov}_{Th_\omega}(t_2, \text{neg}([\Phi_\omega]^c)) \right]. \end{aligned} \quad (3.13)$$

where t_1, t_2, t_3 is a closed term. Note that for any ω -model M_ω^{Th} by the canonical observation one obtains the equivalence $\text{Con}(Th_\omega) \leftrightarrow \text{Con}_{Th_\omega}$ (see [2]). But the Formulae (3.11)-(3.12) contradicts the Formula (3.13). Therefore

$$Th_\omega \not\vdash \Phi_\omega, \not\vdash \text{Pr}_{Th_\omega}([\neg\Phi_\omega]^c) \text{ and } Th_\omega \not\vdash \neg\Phi_\omega.$$

Then theory $Th'_\omega = Th_\omega + \neg\Phi_\omega$ is consistent and from the above observation one obtains that:

$\text{Con}(Th'_\omega) \leftrightarrow \text{Con}_{Th'_\omega}$, where

$$\begin{aligned} \text{Con}_{Th'_\omega} &\leftrightarrow \neg \exists t_1 \neg \exists t_2 \neg \exists t_3 (t_3 = [\Phi_\omega]^c) \\ &\times \left[\text{Prov}_{Th'_\omega}(t_1, [\Phi_\omega]^c) \wedge \text{Prov}_{Th'_\omega}(t_2, \text{neg}([\Phi_\omega]^c)) \right]. \end{aligned} \quad (3.14)$$

On the other hand one obtains

$$Th'_\omega \vdash \text{Pr}_{Th'_\omega}([\Phi_\omega]^c), Th'_\omega \vdash \text{Pr}_{Th'_\omega}([\neg\Phi_\omega]^c). \quad (3.15)$$

But the Formulae (3.15) contradicts the Formula (3.14). This contradiction completed the proof. Proof (ii) similarly as the proof (i) above.

Definition 3.6.

Let Th be a theory such that the Assumption 1.1 is satisfied.

(a) Let $\Xi^{Th_\omega} \equiv \text{Con}(Th; M_\omega^{Th})$ be a sentence in Th asserting that Th has ω -model M_ω^{Th} .

(b) Let $\Xi^{Th_{SM}} \equiv \text{Con}(Th; M_{SM}^{Th})$ be a sentence in Th asserting that Th has standard model M_{SM}^{Th} .

Assumption 3.1. We assume that (i) a sentence Ξ^{Th_ω} is expressible in Th , i.e., a sentence Ξ^{Th_ω} is expressible by using the language \mathcal{L}_{Th} of the Th ; (ii) a sentence $\Xi^{Th_{SM}}$ is expressible in Th , i.e., a sentence $\Xi^{Th_{SM}}$ is expressible by using the language \mathcal{L}_{Th} of the Th .

Remark 3.3. Note that (i) for any ω -model M_ω^{Th} of the Th by the canonical observation (see [2]) one obtains the equivalence

$$\begin{aligned} \text{Con}(Th; M_\omega^{Th}) &\leftrightarrow \\ \text{Con}(Th[M_\omega^{Th}]) &\leftrightarrow \text{Con}_{Th[M_\omega^{Th}]}, \end{aligned} \quad (3.16)$$

(see remark 3.1) and the equivalence

$$\text{Con}_{Th[M_\omega^{Th}]} \leftrightarrow \neg \text{Pr}_{Th[M_\omega^{Th}]} \left(\left[F \left[M_\omega^{Th} \right] \right]^c \right) \quad (3.17)$$

(see remark 3.2), where F is a closed formula refutable in Th .

(ii) for any standard model M_ω^{Th} of the Th by the canonical observation (see [2] chapter), one obtains the equivalence

$$\text{Con}(Th; M_{SM}^{Th}) \leftrightarrow \text{Con}(Th[M_{SM}^{Th}]) \leftrightarrow \text{Con}_{Th[M_{SM}^{Th}]} \quad (3.18)$$

(see remark 3.1) and the equivalence

$$\text{Con}_{Th[M_{SM}^{Th}]} \leftrightarrow \neg \text{Pr}_{Th_{SM}} \left(\left[F \left[M_{SM}^{Th} \right] \right]^c \right) \square \quad (3.19)$$

(see remark 3.2), where F is a closed formula refutable in Th .

Lemma 3.4. (I) Assume that Th has ω -model M_ω^{Th} . Let Th_1 be a theory $Th_1 = Th + \Xi^{Th_\omega}$. Then Th_1 is consistent.

(II) Assume that Th has standard model SM^{Th} .

Let Th_2 be a theory $Th_2 = Th + \Xi^{Th_{SM}}$. Then Th_2 is consistent.

Proof. (I) Assume that a theory

$Th_1 = Th + \Xi^{Th_\omega} \equiv Th + \text{Con}(Th; M_\omega^{Th})$ is inconsistent: $\neg \text{Con}(Th_1)$. This means that there is no any model M^{Th} of Th in which $\text{Con}(Th; M_\omega^{Th})$ is true and in particular that is Th has no any ω -model $M_{1,\omega}^{Th}$ of Th in which $\text{Con}(Th; M_\omega^{Th})$ is true, i.e., $M_{1,\omega}^{Th} \not\models \Xi^{Th_\omega} [M_{1,\omega}^{Th}] \equiv \text{Con}(Th; M_\omega^{Th}) [M_{1,\omega}^{Th}]$ and therefore one obtains for any ω -model $M_{1,\omega}^{Th}$ of Th that

$$M_{1,\omega}^{Th} \models \neg \text{Con}(Th; M_\omega^{Th}) [M_{1,\omega}^{Th}], \quad (3.20)$$

and in particular

$$M_{1,\omega}^{Th} \models \neg \text{Con}(Th; M_{1,\omega}^{Th}) [M_{1,\omega}^{Th}], \quad (3.21)$$

From (3.21) using (3.16)-(3.17) and one obtains

$$M_{1,\omega}^{Th} \models \neg \text{Con}_{Th[M_{1,\omega}^{Th}]} [M_{1,\omega}^{Th}] \leftrightarrow \text{Pr}_{Th[M_{1,\omega}^{Th}]} \left(\left[F \left[M_{1,\omega}^{Th} \right] \right]^c \right). \quad (3.22)$$

From (3.22) and Theorem 3.2(I) one obtains

$$M_{1,\omega}^{Th} \models \left(\left[F \left[M_{1,\omega}^{Th} \right] \right]^c \right). \quad (3.23)$$

Obviously (3.23) contradicts to the assumption that Th has an ω -model M_ω^{Th} . This contradiction completed the proof.

Theorem 3.3. (I) Th has no any ω -model M_ω^{Th} .

(II) Th has no any standard model SM^{Th} .

Proof. (I) By Lemma 3.4(I) one obtains that $Th_1 \vdash \text{Con}(Th_1)$. But Godel's Second Incompleteness Theorem applied to Th_1 asserts that $\text{Con}(Th_1)$ is unprovable in Th_1 . This contradiction completed the proof.

Proof. (II) Similarly as above.

Remark 3.4. We emphasize that it is well known that axiom $\exists SM^{ZFC}$ a single statement in ZFC see [10], Ch. II, section 7. We denote this statement through all this paper by symbol $\text{Con}(ZFC; SM^{ZFC})$.

Theorem 3.4. ZFC has no any ω -model M_ω^{ZFC} .

Proof. Immediately follows from Theorem 3.3 (I) and Remark 3.4.

Theorem 3.5. ZFC has no any standard model. SM^{ZFC} .

Proof. Immediately follows from Theorem 3.3 (II) and Remark 3.4.

Theorem 3.6. ZFC is incompatible with all the usual large cardinal axioms [11] which imply the existence standard model of ZFC .

Proof. Theorem 3.6 immediately follows from Theorem 3.5.

Theorem 3.7. Let κ be an inaccessible cardinal. Then $\neg \text{Con}(ZFC + \exists \kappa)$.

Proof. Let H_κ be a set of all sets having hereditary size less than κ . It easy to see that H_κ forms standard model of ZFC . Therefore Theorem 3.7 immediately follows from Theorem 3.5.

4. Conclusion

In this paper we proved so-called strong reflection principles corresponding to formal theories Th which has ω -models M_ω^{Th} and in particular to formal theories Th , which has a standard models SM^{Th} . The assumption that there exists a standard model of Th is stronger than the assumption that there exists a model of Th . This paper examined some specified classes of the standard models of ZFC so-called strong standard models of ZFC . Such models correspond to large cardinals axioms. In particular we proved that theory $ZFC + \text{Con}(ZFC)$ is incompatible with existence of any inaccessible cardinal κ . Note that the statement: $\text{Con}(ZFC + \exists$ some inaccessible cardinal $\kappa)$ is Π_1^0 . Thus Theorem 3.6 asserts there exist numerical counterexample which would imply that a specific polynomial equation has at least one integer root.

REFERENCES

- [1] M. H. Löb, "Solution of a Problem of Leon Henkin," *The Journal of Symbolic Logic*, Vol. 20, No. 2, 1955, pp. 115-118. [doi:10.2307/2266895](https://doi.org/10.2307/2266895)
- [2] C. Smorynski, "Handbook of Mathematical Logic," North-Holland Publishing Company, 1977.
- [3] T. Drucker, "Perspectives on the History of Mathematical Logic," Birkhauser, Boston, 2008, p. 191.
- [4] A. Marcja and C. Toffalori, "A Guide to Classical and Modern Model Theory (Series: Trends in Logic)," Springer, Berlin, 2003, p. 371.

- [5] F. W. Lawvere, C. Maurer and G. C. Wraith, "Model Theory and Topoi," Springer, Berlin, 1975. pdf
- [6] D. Marker, "Model Theory: An Introduction (Graduate Texts in Mathematics)," Springer, Berlin, 2002.
- [7] J. Foukzon, "Generalized Löb's Theorem," 2013. <http://arxiv.org/abs/1301.5340>
- [8] J. Foukzon, "An Possible Generalization of the Löb's Theorem," *AMS Sectional Meeting AMS Special Session. Spring Western Sectional Meeting University of Colorado Boulder*, Boulder, 13-14 April 2013. http://www.ams.org/amsmtgs/2210_abstracts/1089-03-60.
- [9] P. Lindstrom, "First Order Predicate Logic with Generalized Quantifiers," *Theoria*, Vol. 32, No. 3, 1966, pp. 186-195.
- [10] P. Cohen, "Set theory and the continuum hypothesis," Reprint of the W. A. Benjamin, Inc., New York, 1966 edition; 1966. ISBN-13: 978-0486469218
- [11] A. Kanamori, "The Higher Infinite: Large Cardinals in Set Theory from Their Beginnings," 2nd Edition, Springer, Berlin, 2003.