

Algebraic aspects and coherence conditions for conjoined and disjoined conditionals

Angelo Gilio^{a,1,2}, Giuseppe Sanfilippo^{b,1,3,*}

^a*Department of Basic and Applied Sciences for Engineering, University of Rome “La Sapienza”, Via A. Scarpa 14, 00161 Roma, Italy*

^b*Department of Mathematics and Computer Science, Via Archirafi 34, 90123 Palermo, Italy*

Abstract

We deepen the study of conjoined and disjoined conditional events in the setting of coherence. These objects, differently from other approaches, are defined in the framework of conditional random quantities. We show that some well known properties, valid in the case of unconditional events, still hold in our approach to logical operations among conditional events. In particular we prove a decomposition formula and a related additive property. Then, we introduce the set of conditional constituents generated by n conditional events and we show that they satisfy the basic properties valid in the case of unconditional events. We obtain a generalized inclusion-exclusion formula and we prove a suitable distributivity property. Moreover, under logical independence of basic unconditional events, we give two necessary and sufficient coherence conditions. The first condition gives a geometrical characterization for the coherence of prevision assessments on a family \mathcal{F} constituted by n conditional events and all possible conjunctions among them. The second condition characterizes the coherence of prevision assessments defined on $\mathcal{F} \cup \mathcal{K}$, where \mathcal{K} is the set of conditional constituents associated with the conditional events in \mathcal{F} . Then, we give a further theoretical result and we examine some examples and counterexamples. Finally, we make a comparison with other approaches and we illustrate some theoretical aspects and applications.

Keywords: Coherence, Conditional random quantities, Conjunction and disjunction of conditionals, Decomposition formula, Conditional constituents, Inclusion-exclusion formula.

1. Introduction and motivations

The study of logical operations among conditional events is a relevant topic of research in many fields, such as probability logic, multi-valued logic, artificial intelligence, and psychology

*Int. J. Approx. Reason., <https://doi.org/10.1016/j.ijar.2020.08.004>

*Corresponding author

Email addresses: angelo.gilio@sbai.uniroma1.it (Angelo Gilio), giuseppe.sanfilippo@unipa.it (Giuseppe Sanfilippo)

¹Both authors equally contributed to this work

²Retired

³Also affiliated with INdAM-GNAMPA, Italy

of reasoning; it has been largely discussed and investigated by many authors (see, e.g., [2, 3, 16, 17, 20, 28, 41, 42, 45, 47]). We recall that in a pioneering paper, written in 1935, de Finetti ([25]) proposed a three-valued logic for conditional events, also studied by Lukasiewicz. Moreover, different authors (such as Adams, Belnap, Calabrese, de Finetti, Dubois, van Fraassen, McGee, Goodmann, Lewis, Nguyen, Prade, Schay) have given many contributions to research on three-valued logics and compounds of conditionals (for a survey see, e.g., [46]). Conditionals have been extensively studied also in [24, 45].

Usually, the result of the conjunction or the disjunction of conditionals, as defined in literature, is still a conditional; see e.g. [1, 8, 9, 11, 12, 41]. However, in this way classical probabilistic properties are lost; for instance, differently from the case of unconditional events, the lower and upper probability bounds for the conjunction of two conditional events are no more the Fréchet-Hoeffding bounds; in some cases trivially these bounds are 0 and 1, respectively. This aspect has been recently studied in [51].

A different approach, where the result of conjunction or disjunction of conditionals is not a three-valued object, has been given in [42, 45]. In [32, 33, 36] a related theory has been developed in the setting of coherence, with the advantage (among other things) of properly managing the case where some conditioning events have zero probability. In these papers, the results of conjunction and disjunction of conditional events are *conditional random quantities* with a finite number of possible values in the interval $[0, 1]$.

In addition, it has been proved that the Fréchet-Hoeffding probability bounds continue to hold for the conjunction of two conditional events ([36]). In this paper, we give a related result which concerns the conjunctions associated with two disjoint sub-families of a family of n conditional events.

We show that the conjunction $\mathcal{C}_{1\dots n}$ of n conditional events can be decomposed as the sum of its conjunctions with a further conditional event $E_{n+1}|H_{n+1}$ and the negation $\bar{E}_{n+1}|H_{n+1}$. This result generalizes the well known formula (for the indicators of two unconditional events A and B): $A = AB + A\bar{B}$.

We give a generalization of the inclusion-exclusion formula for the disjunction of a finite number of conditional events. Moreover, we prove the validity of a suitable distributivity property, by means of which we can directly obtain the inclusion-exclusion formula. A main motivation of the paper is that of introducing the conditional constituents for a finite family of conditional events, which can be looked at as a conditional counterpart of atoms of a Boolean algebra. We show that the conditional constituents satisfy the basic numerical and probabilistic properties of the (indicators of the) constituents associated with a finite family of unconditional events.

Under logical independence, we give a necessary and sufficient condition for coherence of a prevision assessment \mathcal{M} on a family \mathcal{F} containing n conditional events and all the $(2^n - n - 1)$ possible conjunctions among them. Such a characterization amounts to the solvability of a linear system which can be interpreted in geometrical terms. Then, the set of all coherent assessments on the family \mathcal{F} is represented by a list of linear inequalities on the components of each prevision assessment \mathcal{M} .

In addition, by considering the set \mathcal{K} of conditional constituents associated with the conditional events in \mathcal{F} , we give a result which under logical independence characterizes the coherence of prevision assessments on $\mathcal{F} \cup \mathcal{K}$. Then, given any coherent assessment \mathcal{M} on \mathcal{F} , we show that every

possible value of the random vector associated with \mathcal{F} is itself a particular coherent assessment on \mathcal{F} . To better illustrate our results, we examine some examples and counterexamples.

Finally, we make a comparison with other approaches, by giving a result related to the notion of atom of a Boolean algebra of conditionals introduced in [27, 28]. In this context, we discuss the significance of our theory by recalling some theoretical aspects and applications.

The paper is organized as follows: In Section 2 we recall some basic notions and results on coherence of conditional probability and prevision assessments. We also recall the definition of conjunction and disjunction among conditional events, and the notion of negation. In Section 3 we first give a result related to Fréchet-Hoeffding bounds; then we illustrate the decomposition formula for the conjunction of n conditional events. In Section 4 we introduce the set \mathcal{K} of conditional constituents for a family of n conditional events $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$. We show that, as in the case of unconditional events, the sum of the conditional constituents is equal to 1 and for each pair of them the conjunction is equal to 0. Then we show that, for each non empty subset $S \subseteq \{1, \dots, n\}$ the conjunction $\mathcal{C}_S = \bigwedge_{i \in S} E_i|H_i$ is the sum of suitable conditional constituents in \mathcal{K} ; hence the prevision of \mathcal{C}_S is the sum of the previsions of such conditional constituents. In Section 5 we give a generalization of the inclusion-exclusion formula for the disjunction of n conditional events. Then, we prove a suitable distributivity property and we examine related probabilistic results. In Section 6, under the hypothesis of logical independence of basic unconditional events, we characterize in terms of a suitable convex hull the set of all coherent prevision assessments on a family \mathcal{F} containing n conditional events and all the possible conjunctions among them. Such a characterization amounts to the solvability of a linear system. Then, we illustrate the set of all coherent assessments on the family \mathcal{F} by a list of linear inequalities on the components of each prevision assessment. We also characterize the coherence of prevision assessments on $\mathcal{F} \cup \mathcal{K}$. In Section 7, given any coherent assessment \mathcal{M} on \mathcal{F} , we show that every possible value of the random vector associated with \mathcal{F} is itself a particular coherent assessment on \mathcal{F} . In Section 8 we illustrate further aspects on coherence by examining some examples and counterexamples. In Section 9, after a comparison with other approaches, we give a result related to the notion of atom of a Boolean algebra of conditionals introduced in [27, 28] and we illustrate some theoretical aspects and applications of our theory. In Section 10 we give some conclusions.

2. Some preliminary notions and results

In this section we recall some basic notions and results which concern coherence (see, e.g., [4, 5, 7, 10, 15, 49, 50]) and logical operations among conditional events (see [32, 33, 36, 37, 39]).

2.1. Events and conditional events

An event E is an uncertain fact described by a (non ambiguous) logical proposition; in formal terms E is a two-valued logical entity which can be *true*, or *false*. The *indicator* of E , denoted by the same symbol, is 1, or 0, according to whether E is true, or false. The sure event and impossible event are denoted by Ω and \emptyset , respectively. Given two events E_1 and E_2 , we denote by $E_1 \wedge E_2$, or simply by $E_1 E_2$, (resp., $E_1 \vee E_2$) the logical conjunction (resp., the logical disjunction). The negation of E is denoted \bar{E} . We simply write $E_1 \subseteq E_2$ to denote that E_1 logically implies E_2 , that

is $E_1\bar{E}_2 = \emptyset$. We recall that n events E_1, \dots, E_n are logically independent when the number m of constituents, or possible worlds, generated by them is 2^n (in general $m \leq 2^n$).

Given two events E, H , with $H \neq \emptyset$, the conditional event $E|H$ is defined as a three-valued logical entity which is *true*, or *false*, or *void*, according to whether EH is true, or $\bar{E}H$ is true, or \bar{H} is true, respectively. Given a family $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$, we observe that, for each i , it holds that $E_iH_i \vee \bar{E}_iH_i \vee \bar{H}_i = \Omega$; then by expanding the expression $\bigwedge_{i=1}^n (E_iH_i \vee \bar{E}_iH_i \vee \bar{H}_i)$ we can represent Ω as the disjunction of 3^n logical conjunctions, some of which may be impossible. The remaining ones are the constituents generated by \mathcal{E} and, of course, are a partition of Ω . We denote by C_1, \dots, C_m the constituents which logically imply the event $\mathcal{H}_n = H_1 \vee \dots \vee H_n$. Moreover, (if $\mathcal{H}_n \neq \Omega$) we denote by C_0 the remaining constituent $\bar{\mathcal{H}}_n = \bar{H}_1 \cdots \bar{H}_n$. Thus

$$\mathcal{H}_n = C_1 \vee \dots \vee C_m, \quad \Omega = \bar{\mathcal{H}}_n \vee \mathcal{H}_n = C_0 \vee C_1 \vee \dots \vee C_m, \quad m + 1 \leq 3^n.$$

For instance, given four logically independent events E_1, E_2, H_1, H_2 , the constituents generated by $\mathcal{E} = \{E_1|H_1, E_2|H_2\}$ are $C_1 = E_1H_1E_2H_2$, $C_2 = E_1H_1\bar{E}_2H_2$, $C_3 = \bar{E}_1H_1E_2H_2$, $C_4 = \bar{E}_1H_1\bar{E}_2H_2$, $C_5 = \bar{H}_1E_2H_2$, $C_6 = \bar{H}_1\bar{E}_2H_2$, $C_7 = E_1H_1\bar{H}_2$, $C_8 = \bar{E}_1H_1\bar{H}_2$, $C_0 = \bar{H}_1\bar{H}_2$.

2.2. Coherent conditional prevision assessments for conditional random quantities

Given a (real) random quantity X and an event $H \neq \emptyset$, we denote by $\mathbb{P}(X|H)$ the prevision of X conditional on H . In the framework of coherence, to assess $\mathbb{P}(X|H) = \mu$ means that, for every real number s , you are willing to pay an amount $s\mu$ and to receive sX , or $s\mu$, according to whether H is true, or \bar{H} is true (the bet is called off), respectively. The random gain is $G = s(XH + \mu\bar{H}) - s\mu = sH(X - \mu)$.

As we will see, a conjunction of n conditional events is a conditional random quantity with a finite number of possible (numerical) values. Then, in what follows, for any given conditional random quantity $X|H$, we assume that, when H is true, the set of possible values of X is a finite subset of the set of real numbers \mathbb{R} . In this case we say that $X|H$ is a finite conditional random quantity. Given a prevision function \mathbb{P} defined on an arbitrary family \mathcal{K} of finite conditional random quantities, consider a finite subfamily $\mathcal{F} = \{X_1|H_1, \dots, X_n|H_n\} \subseteq \mathcal{K}$ and the vector $\mathcal{M} = (\mu_1, \dots, \mu_n)$, where $\mu_i = \mathbb{P}(X_i|H_i)$ is the assessed prevision for the conditional random quantity $X_i|H_i$, $i \in \{1, \dots, n\}$. With the pair $(\mathcal{F}, \mathcal{M})$ we associate the random gain $G = \sum_{i=1}^n s_i H_i (X_i - \mu_i)$. We denote by $\mathcal{G}_{\mathcal{H}_n}$ the set of values of G restricted to $\mathcal{H}_n = H_1 \vee \dots \vee H_n$. Then, by the *betting scheme* of de Finetti, the notion of coherence is defined as below.

Definition 1. The prevision function \mathbb{P} defined on \mathcal{K} is coherent if and only if, $\forall n \geq 1, \forall \mathcal{F} = \{X_1|H_1, \dots, X_n|H_n\} \subseteq \mathcal{K}$, it holds that: $\min \mathcal{G}_{\mathcal{H}_n} \leq 0 \leq \max \mathcal{G}_{\mathcal{H}_n}, \forall s_1, \dots, s_n$.

A conditional prevision assessment \mathbb{P} on \mathcal{K} is said incoherent if and only if there exists a finite combination of n bets such that $\min \mathcal{G}_{\mathcal{H}_n} \cdot \max \mathcal{G}_{\mathcal{H}_n} > 0$, that is such that the values in $\mathcal{G}_{\mathcal{H}_n}$ are all positive, or all negative (*Dutch Book*). In the particular case where \mathcal{K} is a family of conditional events, then Definition 1 becomes the well known definition of coherence for a probability function \mathbb{P} , denoted as P , defined on \mathcal{K} .

Given a family $\mathcal{F} = \{X_1|H_1, \dots, X_n|H_n\}$, for each $i = 1, \dots, n$, we denote by $\{x_{i1}, \dots, x_{ir_i}\}$ the set of possible (numerical) values for the restriction of X_i to H_i ; then, for each $i = 1, \dots, n$, and

$j = 1, \dots, r_i$, we set $A_{ij} = (X_i = x_{ij})$. Of course, for each i , the family $\{\bar{H}_i, A_{ij}H_i, j = 1, \dots, r_i\}$ is a partition of the sure event Ω , with $A_{ij}H_i = A_{ij}$ and $\bigvee_{j=1}^{r_i} A_{ij} = H_i$. Then, the constituents generated by the family \mathcal{F} are (the elements of the partition of Ω) obtained by expanding the expression $\bigwedge_{i=1}^n (A_{i1} \vee \dots \vee A_{ir_i} \vee \bar{H}_i)$. We set $C_0 = \bar{H}_1 \dots \bar{H}_n = \bar{\mathcal{H}}_n$ (it may be $C_0 = \emptyset$); moreover, we denote by C_1, \dots, C_m the constituents contained in \mathcal{H}_n . Hence $\bigwedge_{i=1}^n (A_{i1} \vee \dots \vee A_{ir_i} \vee \bar{H}_i) = \bigvee_{h=0}^m C_h$. With each $C_h, h = 1, \dots, m$, we associate a vector $Q_h = (q_{h1}, \dots, q_{hn})$, where $q_{hi} = x_{ij}$ if $C_h \subseteq A_{ij}, j = 1, \dots, r_i$, while $q_{hi} = \mu_i$ if $C_h \subseteq \bar{H}_i$; with C_0 it is associated $Q_0 = \mathcal{M} = (\mu_1, \dots, \mu_n)$. As, for each i, j , the quantities x_{ij}, μ_i are real numbers, it holds that $Q_h \in \mathbb{R}^n, h = 0, 1, \dots, m$.

Denoting by \mathcal{I} the convex hull of Q_1, \dots, Q_m , the condition $\mathcal{M} \in \mathcal{I}$ amounts to the existence of a vector $(\lambda_1, \dots, \lambda_m)$ such that: $\sum_{h=1}^m \lambda_h Q_h = \mathcal{M}, \sum_{h=1}^m \lambda_h = 1, \lambda_h \geq 0, \forall h$; in other words, $\mathcal{M} \in \mathcal{I}$ is equivalent to the solvability of the system (Σ) , associated with $(\mathcal{F}, \mathcal{M})$, given below.

$$(\Sigma) \begin{cases} \sum_{h=1}^m \lambda_h q_{hi} = \mu_i, & i = 1, \dots, n, \\ \sum_{h=1}^m \lambda_h = 1, & \lambda_h \geq 0, \quad h = 1, \dots, m. \end{cases} \quad (1)$$

Given the assessment $\mathcal{M} = (\mu_1, \dots, \mu_n)$ on $\mathcal{F} = \{X_1|H_1, \dots, X_n|H_n\}$, let S be the set of solutions $\Lambda = (\lambda_1, \dots, \lambda_m)$ of system (Σ) defined in (1). Then, the following characterization theorem for coherent assessments on finite families of conditional random quantities can be proved ([6]).

Theorem 1. [*Characterization of coherence*]. Given a family of n conditional random quantities $\mathcal{F} = \{X_1|H_1, \dots, X_n|H_n\}$, with finite sets of possible values, and a vector $\mathcal{M} = (\mu_1, \dots, \mu_n)$, the conditional prevision assessment $\mathbb{P}(X_1|H_1) = \mu_1, \dots, \mathbb{P}(X_n|H_n) = \mu_n$ is coherent if and only if, for every subset $J \subseteq \{1, \dots, n\}$, defining $\mathcal{F}_J = \{X_i|H_i, i \in J\}, \mathcal{M}_J = (\mu_i, i \in J)$, the system (Σ_J) associated with the pair $(\mathcal{F}_J, \mathcal{M}_J)$ is solvable.

As shown by Theorem 1, the solvability of system (Σ) (i.e., the condition $\mathcal{M} \in \mathcal{I}$) is a necessary (but not sufficient) condition for coherence of \mathcal{M} on \mathcal{F} . Given the assessment \mathcal{M} on \mathcal{F} , let S be the set of solutions $\Lambda = (\lambda_1, \dots, \lambda_m)$ of system (Σ) defined in (1). By assuming the system (Σ) solvable, that is $S \neq \emptyset$, we define:

$$I_0 = \{i : \max_{\Lambda \in S} \sum_{h: C_h \subseteq H_i} \lambda_h = 0\}, \quad \mathcal{F}_0 = \{X_i|H_i, i \in I_0\}, \quad \mathcal{M}_0 = (\mu_i, i \in I_0). \quad (2)$$

We observe that $i \in I_0$ if and only if the (unique) coherent extension of \mathcal{M} to $H_i|\mathcal{H}_n$ is zero. Then, the following theorem can be proved ([6, Theorem 3]):

Theorem 2. [*Operative characterization of coherence*] A conditional prevision assessment $\mathcal{M} = (\mu_1, \dots, \mu_n)$ on the family $\mathcal{F} = \{X_1|H_1, \dots, X_n|H_n\}$ is coherent if and only if the following conditions are satisfied:

(i) the system (Σ) defined in (1) is solvable; (ii) if $I_0 \neq \emptyset$, then \mathcal{M}_0 is coherent.

In order to illustrate the previous results, we examine an example.

Example 1. Let E, H, K be three events, with $HK = \emptyset$ and E logically independent of H and K . Moreover, let $\mathcal{P} = (x, y)$ be a probability assessment on the family $\mathcal{E} = \{E|H, E|K\}$, where $x = P(E|H)$ and $y = P(E|K)$. The constituents generated by \mathcal{E} are: $C_1 = EHK, C_2 = E\bar{H}K,$

$C_3 = \bar{E}H\bar{K}$, $C_4 = \bar{E}\bar{H}K$, $C_0 = \bar{H}\bar{K}$. Then, the associated points Q_h 's are: $Q_1 = (1, y)$, $Q_2 = (x, 1)$, $Q_3 = (0, y)$, $Q_4 = (x, 0)$, $Q_0 = \mathcal{P} = (x, y)$. The system (Σ) is

$$\left\{ \begin{array}{l} \lambda_1 + \lambda_2 x + \lambda_4 x = x, \quad \lambda_1 y + \lambda_2 + \lambda_3 y = y, \quad \lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 = 1, \quad \lambda_h \geq 0, \quad \forall h. \end{array} \right.$$

As it can be verified, for each $(x, y) \in [0, 1]^2$, the vector $(\lambda_1, \dots, \lambda_4) = (\frac{x}{2}, \frac{y}{2}, \frac{1-x}{2}, \frac{1-y}{2})$ is a solution of (Σ) . Moreover, for this solution it holds that

$$\sum_{C_h \subseteq H} \lambda_h = \lambda_1 + \lambda_3 = \frac{1}{2} > 0; \quad \sum_{C_h \subseteq K} \lambda_h = \lambda_2 + \lambda_4 = \frac{1}{2} > 0.$$

Then, $I_0 = \emptyset$ and by Theorem 2 the assessment (x, y) is coherent, for every $(x, y) \in [0, 1]^2$.

2.3. A deepening on the notion of conditional random quantity

We recall that, in the subjective approach to probability theory, given an event $H \neq \emptyset$ and a random quantity X by the betting metaphor the conditional prevision $\mathbb{P}(X|H)$ is defined as the amount μ you agree to pay, by knowing that you will receive the amount $XH + \mu\bar{H}$. This quantity coincides with X , if H is true, or with μ , if H is false (bet called off). Usually, in literature the conditional random quantity $X|H$ is defined as the *restriction* of X to H , which coincides with X , when H is true, and it is undefined when H is false. Under this point of view, (when H is false) $X|H$ does not coincide with $XH + \mu\bar{H}$. However, by coherence, it holds that $\mathbb{P}(XH + \mu\bar{H}) = \mathbb{P}(X|H)P(H) + \mu P(\bar{H}) = \mu P(H) + \mu P(\bar{H}) = \mu$. Then, we can extend the notion of $X|H$, by defining its value as equal to μ when H is false (for further details see [36]). In this way $X|H$ coincides with $XH + \mu\bar{H}$ and in the betting scheme it can be interpreted as the amount that you receive when you pay its prevision μ . In addition, the random gain G can be represented as $G = s(X|H - \mu)$. In particular, when X is the indicator of an event E , we obtain $X|H = EH + P(E|H)\bar{H}$ and it holds that

$$\mathbb{P}(X|H) = \mathbb{P}[EH + P(E|H)\bar{H}] = P(E|H)P(H) + P(E|H)P(\bar{H}) = P(E|H).$$

In this case $X|H$ is the indicator of the conditional event $E|H$ (which we denote by the same symbol) and, by defining $P(E|H) = x$, it holds that

$$E|H = EH + x\bar{H} = EH + x(1 - H) = \begin{cases} 1, & \text{if } EH \text{ is true,} \\ 0, & \text{if } \bar{E}H \text{ is true,} \\ x, & \text{if } \bar{H} \text{ is true.} \end{cases} \quad (3)$$

For related discussions, see also [14, 32, 43]. By Definition 1, the coherence of the assessment $P(E|H) = x$ is equivalent to $\min \mathcal{G}_H \leq 0 \leq \max \mathcal{G}_H$, $\forall s$, where \mathcal{G}_H is the set of values of G restricted to H . Then, the set Π of coherent assessments x on $E|H$ is: (i) $\Pi = [0, 1]$, when $\emptyset \neq EH \neq H$; (ii) $\Pi = \{0\}$, when $EH = \emptyset$; (iii) $\Pi = \{1\}$, when $EH = H$. Of course, the third value of the random quantity $E|H$ depends on the subjective assessment $P(E|H) = x$. Notice that, when $H \subseteq E$ (i.e., $EH = H$), by coherence $P(E|H) = 1$ and hence for the indicator it holds that $E|H = H + \bar{H} = 1$.

By exploiting our extended notion of conditional random quantity, we can develop some algebraic aspects ([32, Section 3],[36, Section 3.2]). For instance, we can show that:

- denoting by μ and ν the previsions of $X|H$ and $Y|K$, respectively, the sum $X|H + Y|K$ coincides with the conditional random quantity $(XH + \mu\bar{H} + YK + \nu\bar{K})|(H \vee K)$, with $\mathbb{P}(X|H + Y|K) = \mathbb{P}(X|H) + \mathbb{P}(Y|K) = \mu + \nu$;
- $a(X|H) + b(Y|K) = (aX)|H + (bY)|K$, where a, b are real numbers;
- $\mathbb{P}(XH|K) = P(H|K)\mathbb{P}(X|HK)$, which is the *compound prevision theorem*.

Moreover, as shown by the result below, if $X|H$ and $Y|K$ coincide when $H \vee K$ is true, then their previsions are equal and it follows that $X|H$ and $Y|K$ also coincide when $H \vee K$ is false, so that $X|H = Y|K$ in all cases ([36, Theorem 4]).

Theorem 3. Given any events $H \neq \emptyset, K \neq \emptyset$, and any r.q.'s X, Y , let Π be the set of the coherent prevision assessments $\mathbb{P}(X|H) = \mu, \mathbb{P}(Y|K) = \nu$.

(i) Assume that, for every $(\mu, \nu) \in \Pi$, $X|H = Y|K$ when $H \vee K$ is true; then $\mu = \nu$ for every $(\mu, \nu) \in \Pi$.

(ii) For every $(\mu, \nu) \in \Pi$, $X|H = Y|K$ when $H \vee K$ is true if and only if $X|H = Y|K$.

To better illustrate Theorem 3, we observe that

$$X|H - Y|K = (XH + \mu\bar{H} - YK - \nu\bar{K})|(H \vee K).$$

Now, assume that $X|H$ and $Y|K$ coincide when $H \vee K$ is true, so that $X|H - Y|K = 0$ when $H \vee K$ is true. If $H \vee K$ is false, that is $\bar{H}\bar{K}$ is true, it holds that $X|H = \mu$ and $Y|K = \nu$, so that $X|H - Y|K = \mu - \nu$. Then, in a conditional bet on the conditional random quantity $X|H - Y|K$, if you pay $\mu - \nu = \mathbb{P}(X|H - Y|K)$, you receive zero when $H \vee K$ is true, or you receive back $\mu - \nu$ when $H \vee K$ is false (bet called off). Then, by coherence, it must be $\mu - \nu = 0$, that is $\mu = \nu$, and hence $X|H = Y|K$.

Remark 1. Theorem 3 has been generalized in [39, Theorem 6] by replacing the symbol “=” by “ \leq ” in statements (i) and (ii). In other words, if $X|H \leq Y|K$ when $H \vee K$ is true, then $\mathbb{P}(X|H) \leq \mathbb{P}(Y|K)$ and hence $X|H \leq Y|K$ in all cases.

2.4. Logical operations among conditional events

We recall below the notions of conjunction and disjunction of two conditional events.

Definition 2. Given any pair of conditional events $E_1|H_1$ and $E_2|H_2$, with $P(E_1|H_1) = x_1$ and $P(E_2|H_2) = x_2$, their conjunction $(E_1|H_1) \wedge (E_2|H_2)$ is the conditional random quantity defined as

$$\begin{aligned} (E_1|H_1) \wedge (E_2|H_2) &= (E_1H_1E_2H_2 + x_1\bar{H}_1E_2H_2 + x_2\bar{H}_2E_1H_1)|(H_1 \vee H_2) = \\ &= \begin{cases} 1, & \text{if } E_1H_1E_2H_2 \text{ is true,} \\ 0, & \text{if } \bar{E}_1\bar{H}_1 \vee \bar{E}_2\bar{H}_2 \text{ is true,} \\ x_1, & \text{if } \bar{H}_1E_2H_2 \text{ is true,} \\ x_2, & \text{if } \bar{H}_2E_1H_1 \text{ is true,} \\ x_{12}, & \text{if } \bar{H}_1\bar{H}_2 \text{ is true,} \end{cases} \end{aligned} \quad (4)$$

where $x_{12} = \mathbb{P}[(E_1|H_1) \wedge (E_2|H_2)] = \mathbb{P}[(E_1H_1E_2H_2 + x_1\bar{H}_1E_2H_2 + x_2\bar{H}_2E_1H_1)|(H_1 \vee H_2)]$.

In betting terms, the prevision x_{12} represents the amount you agree to pay, with the proviso that you will receive the quantity $E_1H_1E_2H_2 + x_1\bar{H}_1E_2H_2 + x_2\bar{H}_2E_1H_1$, or you will receive back the quantity x_{12} , according to whether $H_1 \vee H_2$ is true, or $\bar{H}_1\bar{H}_2$ is true. In other words, by paying x_{12} you receive $E_1H_1E_2H_2 + x_1\bar{H}_1E_2H_2 + x_2\bar{H}_2E_1H_1 + x_{12}\bar{H}_1\bar{H}_2$, which assumes one of the following values:

- 1, if both conditional events are true;
- 0, if at least one of the conditional events is false;
- the probability of the conditional event that is void if one conditional event is void and the other one is true;
- x_{12} (the amount that you paid) if both conditional events are void.

Remark 2. By recalling (3), we again emphasize that there is a different indicator of a conditional event $E|H$ for each coherent evaluation of $P(E|H)$. The same comment applies to the conjunction $(E_1|H_1) \wedge (E_2|H_2)$; indeed, each different conjunction is associated to a different coherent assessment (x_1, x_2, x_{12}) . We also remark that Definition 2 is not circular because, after assessing (x_1, x_2) , the conjunction is completely specified once by the betting scheme you, coherently with (x_1, x_2) , decide the value $x_{12} = \mathbb{P}[(E_1H_1E_2H_2 + x_1\bar{H}_1E_2H_2 + x_2\bar{H}_2E_1H_1)|(H_1 \vee H_2)]$.

We recall a result which shows that Fréchet-Hoeffding bounds still hold for the conjunction of conditional events ([36, Theorem 7]).

Theorem 4. Given any coherent assessment (x_1, x_2) on $\{E_1|H_1, E_2|H_2\}$, with E_1, H_1, E_2, H_2 logically independent, $H_1 \neq \emptyset, H_2 \neq \emptyset$, the extension $x_{12} = \mathbb{P}[(E_1|H_1) \wedge (E_2|H_2)]$ is coherent if and only if the following Fréchet-Hoeffding bounds are satisfied:

$$\max\{x_1 + x_2 - 1, 0\} = x'_{12} \leq x_{12} \leq x''_{12} = \min\{x_1, x_2\}. \quad (5)$$

Remark 3. From Theorem 4, as the assessment (x_1, x_2) on $\{E_1|H_1, E_2|H_2\}$ is coherent for every $(x_1, x_2) \in [0, 1]^2$, the set Π of all coherent prevision assessments (x_1, x_2, x_{12}) on $\{E_1|H_1, E_2|H_2, (E_1|H_1) \wedge (E_2|H_2)\}$ is

$$\Pi = \{(x_1, x_2, x_{12}) : (x_1, x_2) \in [0, 1]^2, \max\{x_1 + x_2 - 1, 0\} \leq x_{12} \leq \min\{x_1, x_2\}\}, \quad (6)$$

which is the tetrahedron with vertices the points $(1, 1, 1), (1, 0, 0), (0, 1, 0), (0, 0, 0)$.

Other related approaches to compound conditionals have been developed in [42, 45]. However, in our coherence-based approach we can properly manage the case where the probability of some conditioning events is zero. Then, differently from other authors, we can compute lower and upper bounds for conjunction and disjunction only in terms of the probabilities of the two given conditional events. We recall below the notion of disjunction between two conditional events.

Definition 3. Given any pair of conditional events $E_1|H_1$ and $E_2|H_2$, with $P(E_1|H_1) = x_1$ and $P(E_2|H_2) = x_2$, their disjunction $(E_1|H_1) \vee (E_2|H_2)$ is the conditional random quantity defined as

$$\begin{aligned} (E_1|H_1) \vee (E_2|H_2) &= (E_1H_1 \vee E_2H_2 + x_1\bar{H}_1\bar{E}_2H_2 + x_2\bar{H}_2\bar{E}_1H_1)|(H_1 \vee H_2) = \\ &= \begin{cases} 1, & \text{if } E_1H_1 \vee E_2H_2 \text{ is true,} \\ 0, & \text{if } \bar{E}_1H_1\bar{E}_2H_2 \text{ is true,} \\ x_1, & \text{if } \bar{H}_1\bar{E}_2H_2 \text{ is true,} \\ x_2, & \text{if } \bar{H}_2\bar{E}_1H_1 \text{ is true,} \\ y_{12}, & \text{if } \bar{H}_1\bar{H}_2 \text{ is true,} \end{cases} \end{aligned} \quad (7)$$

where $y_{12} = \mathbb{P}[(E_1|H_1) \vee (E_2|H_2)] = \mathbb{P}[(E_1H_1 \vee E_2H_2 + x_1\bar{H}_1\bar{E}_2H_2 + x_2\bar{H}_2\bar{E}_1H_1)|(H_1 \vee H_2)]$.

Of course, the assessment (x_1, x_2, y_{12}) must be coherent. In betting terms, y_{12} represents the amount you agree to pay, with the proviso that you will receive the quantity $E_1H_1 \vee E_2H_2 + x_1\bar{H}_1\bar{E}_2H_2 + x_2\bar{H}_2\bar{E}_1H_1 + y_{12}\bar{H}_1\bar{H}_2$, which assumes one of the following values:

- 1, if at least one of the conditional events is true;
- 0, if both conditional events are false;
- the probability of the conditional event that is void if one conditional event is void and the other one is false;
- y_{12} (the amount that you paid) if both conditional events are void.

Notice that, differently from conditional events which are three-valued objects, the conjunction $(E_1|H_1) \wedge (E_2|H_2)$ and the disjunction $(E_1|H_1) \vee (E_2|H_2)$ are not any longer three-valued objects, but five-valued objects. Moreover, the comments of Remark 2 also apply in a dual way to disjunction.

We give below the notion of conjunction of n conditional events.

Definition 4. Let n conditional events $E_1|H_1, \dots, E_n|H_n$ be given. For each non-empty strict subset S of $\{1, \dots, n\}$, let x_S be a prevision assessment on $\bigwedge_{i \in S} (E_i|H_i)$. Then, the conjunction $(E_1|H_1) \wedge \dots \wedge (E_n|H_n)$ is the conditional random quantity $\mathcal{C}_{1\dots n}$ defined as

$$\begin{aligned} \mathcal{C}_{1\dots n} &= [\bigwedge_{i=1}^n E_iH_i + \sum_{\emptyset \neq S \subset \{1,2,\dots,n\}} x_S (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_iH_i)] | (\bigvee_{i=1}^n H_i) = \\ &= \begin{cases} 1, & \text{if } \bigwedge_{i=1}^n E_iH_i \text{ is true,} \\ 0, & \text{if } \bigvee_{i=1}^n \bar{E}_iH_i \text{ is true,} \\ x_S, & \text{if } (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_iH_i) \text{ is true, } \emptyset \neq S \subset \{1, 2, \dots, n\}, \\ x_{1\dots n}, & \text{if } \bigwedge_{i=1}^n \bar{H}_i \text{ is true,} \end{cases} \end{aligned} \quad (8)$$

where

$$x_{1\dots n} = x_{\{1,\dots,n\}} = \mathbb{P}(\mathcal{C}_{1\dots n}) = \mathbb{P}[(\bigwedge_{i=1}^n E_iH_i + \sum_{\emptyset \neq S \subset \{1,2,\dots,n\}} x_S (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_iH_i)) | (\bigvee_{i=1}^n H_i)].$$

For $n = 1$ we obtain $\mathcal{C}_1 = E_1|H_1$. In Definition 4 each possible value x_S of $\mathcal{C}_{1\dots n}$, $\emptyset \neq S \subset \{1, \dots, n\}$, is evaluated when defining (in a previous step) the conjunction $\mathcal{C}_S = \bigwedge_{i \in S} (E_i|H_i)$. Then, after the conditional prevision $x_{1\dots n}$ is evaluated, $\mathcal{C}_{1\dots n}$ is completely specified. Of course, we require coherence for the prevision assessment $(x_S, \emptyset \neq S \subseteq \{1, \dots, n\})$, so that $\mathcal{C}_{1\dots n} \in [0, 1]$. In the framework of the betting scheme, $x_{1\dots n}$ is the amount that you agree to pay with the proviso that you will receive:

- 1, if all conditional events are true;
- 0, if at least one of the conditional events is false;
- the prevision of the conjunction of that conditional events which are void, otherwise. In particular you receive back $x_{1\dots n}$ when all conditional events are void.

We observe that conjunction satisfies the monotonicity property ([39, Theorem7]), that is

$$\mathcal{C}_{1\dots n+1} \leq \mathcal{C}_{1\dots n}. \quad (9)$$

We recall the following result ([39, Theorem13]).

Theorem 5. Let n conditional events $E_1|H_1, \dots, E_n|H_n$ be given, with $x_i = P(E_i|H_i)$, $i = 1, \dots, n$ and $x_{1\dots n} = \mathbb{P}(\mathcal{C}_{1\dots n})$. Then: $\max\{x_1 + \dots + x_n - n + 1, 0\} \leq x_{1\dots n} \leq \min\{x_1, \dots, x_n\}$.

We give below the notion of disjunction of n conditional events.

Definition 5. Let n conditional events $E_1|H_1, \dots, E_n|H_n$ be given. For each non-empty strict subset S of $\{1, \dots, n\}$, let y_S be a prevision assessment on $\bigvee_{i \in S} (E_i|H_i)$. Then, the disjunction $(E_1|H_1) \vee \dots \vee (E_n|H_n)$ is the conditional random quantity $\mathcal{D}_{1\dots n}$ defined as

$$\begin{aligned} \mathcal{D}_{1\dots n} &= (\bigvee_{i=1}^n E_i H_i + \sum_{\emptyset \neq S \subset \{1, 2, \dots, n\}} y_S (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} \bar{E}_i H_i)) | (\bigvee_{i=1}^n H_i) = \\ &= \begin{cases} 1, & \text{if } \bigvee_{i=1}^n E_i H_i \text{ is true,} \\ 0, & \text{if } \bigwedge_{i=1}^n \bar{E}_i H_i \text{ is true,} \\ y_S, & \text{if } (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} \bar{E}_i H_i) \text{ is true, } \emptyset \neq S \subset \{1, 2, \dots, n\}, \\ y_{1\dots n}, & \text{if } \bigwedge_{i=1}^n \bar{H}_i \text{ is true,} \end{cases} \quad (10) \end{aligned}$$

where

$$y_{1\dots n} = y_{\{1, \dots, n\}} = \mathbb{P}(\mathcal{D}_{1\dots n}) = \mathbb{P}[(\bigvee_{i=1}^n E_i H_i + \sum_{\emptyset \neq S \subset \{1, 2, \dots, n\}} y_S (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} \bar{E}_i H_i)) | (\bigvee_{i=1}^n H_i)].$$

For $n = 1$ we obtain $\mathcal{D}_1 = E_1|H_1$. In the betting framework, you agree to pay $y_{1\dots n}$ with the proviso that you will receive:

- 1, if at least one of the conditional events is true;
- 0, if all conditional events are false;
- the prevision of the disjunction of that conditional events which are void, otherwise. In particular you receive back $y_{1\dots n}$ when all conditional events are void.

As we can see from (8) and (10), the conjunction $\mathcal{C}_{1\dots n}$ and the disjunction $\mathcal{D}_{1\dots n}$ are (in general) $(2^n + 1)$ -valued objects because the number of nonempty subsets S , and hence the number of possible values x_S , is $2^n - 1$. Of course, it may happen that the some of the possible values of $\mathcal{C}_{1\dots n}$ and $\mathcal{D}_{1\dots n}$ coincide.

Remark 4. Given a finite family \mathcal{E} of conditional events, their conjunction and disjunction are also denoted by $\mathcal{C}(\mathcal{E})$ and $\mathcal{D}(\mathcal{E})$, respectively. We recall that in [39], given two finite families of conditional events \mathcal{E}' and \mathcal{E}'' , the objects $\mathcal{C}(\mathcal{E}') \wedge \mathcal{C}(\mathcal{E}'')$ and $\mathcal{D}(\mathcal{E}') \vee \mathcal{D}(\mathcal{E}'')$ are defined as $\mathcal{C}(\mathcal{E}' \cup \mathcal{E}'')$ and $\mathcal{D}(\mathcal{E}' \cup \mathcal{E}'')$, respectively. Then, it is easy to verify the commutativity and associativity properties of conjunction and disjunction ([39, Propositions 1 and 2]). We recall below the notion of negation for conjoined and disjointed conditionals.

Definition 6. Given n conditional events $E_1|H_1, \dots, E_n|H_n$, the negations for the conjunction $\mathcal{C}_{1\dots n}$ and the disjunction $\mathcal{D}_{1\dots n}$ are defined as $\bar{\mathcal{C}}_{1\dots n} = 1 - \mathcal{C}_{1\dots n}$ and $\bar{\mathcal{D}}_{1\dots n} = 1 - \mathcal{D}_{1\dots n}$, respectively.

Of course, if $n = 1$ we obtain $\bar{\mathcal{C}}_1 = \bar{\mathcal{D}}_1 = \overline{E_1|H_1} = 1 - E_1|H_1 = \bar{E}_1|H_1$. We observe that conjunction and disjunction satisfy De Morgan's Laws ([39, Theorem 5]), that is

$$\bar{\mathcal{D}}_{1\dots n} = \mathcal{C}_{\bar{1}\dots\bar{n}} \quad (\text{i.e., } \mathcal{D}_{1\dots n} = \bar{\mathcal{C}}_{\bar{1}\dots\bar{n}}), \quad \bar{\mathcal{C}}_{1\dots n} = \mathcal{D}_{\bar{1}\dots\bar{n}} \quad (\text{i.e., } \mathcal{C}_{1\dots n} = \bar{\mathcal{D}}_{\bar{1}\dots\bar{n}}), \quad (11)$$

where $\mathcal{C}_{\bar{1}\dots\bar{n}} = \bigwedge_{i=1}^n \bar{E}_i|H_i$ and $\mathcal{D}_{\bar{1}\dots\bar{n}} = \bigvee_{i=1}^n \bar{E}_i|H_i$. As shown in formula (11), by exploiting negation, disjunction could be equivalently defined as $\mathcal{D}_{1\dots n} = \bar{\mathcal{C}}_{\bar{1}\dots\bar{n}} = 1 - \mathcal{C}_{\bar{1}\dots\bar{n}}$.

3. A decomposition formula for conjunctions

In this section we show that the conjunction $\mathcal{C}_{1\dots n}$ of n conditional events can be represented as the sum of two suitable conjunctions of $n + 1$ conditional events. We first give a preliminary result, which is related to Theorem 4, and a remark.

Theorem 6. Let n conditional events $E_1|H_1, \dots, E_k|H_k, \dots, E_n|H_n$ be given, with $E_1, H_1, \dots, E_n, H_n$ logically independent, and a coherent prevision assessment $\mathcal{M} = (x_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ on the family $\mathcal{F} = \{\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, n\}\}$. For every $1 \leq k \leq n - 1$ it holds that

$$\max\{0, x_{1\dots k} + x_{k+1\dots n} - 1\} \leq x_{1\dots n} \leq \min\{x_{1\dots k}, x_{k+1\dots n}\},$$

where

$$x_{1\dots k} = \mathbb{P}(\mathcal{C}_{1\dots k}), \quad x_{k+1\dots n} = \mathbb{P}(\mathcal{C}_{k+1\dots n}), \quad x_{1\dots n} = \mathbb{P}(\mathcal{C}_{1\dots n}).$$

Proof. We set $\mathcal{M}_3 = (x_{1\dots k}, x_{k+1\dots n}, x_{1\dots n})$. Moreover, we observe that

$$\mathcal{C}_{1\dots k} \in \{1, 0, x_{S'}; S' \subseteq \{1, \dots, k\}\}, \quad \mathcal{C}_{k+1\dots n} \in \{1, 0, x_{S''}; S'' \subseteq \{k+1, \dots, n\}\}.$$

The possible values Q_h 's of the random vector $(\mathcal{C}_{1\dots k}, \mathcal{C}_{k+1\dots n}, \mathcal{C}_{1\dots n})$ are given in Table 1. We

| C_h | $\mathcal{C}_{1\dots k}$ | $\mathcal{C}_{k+1\dots n}$ | $\mathcal{C}_{1\dots n}$ | Q_h |
|---|--------------------------|----------------------------|--------------------------|--|
| $\bigwedge_{i=1}^n E_i H_i$ | 1 | 1 | 1 | $(1, 1, 1)$ |
| $(\bigwedge_{i=1}^k E_i H_i)(\bigvee_{i=k+1}^n \bar{E}_i H_i)$ | 1 | 0 | 0 | $(1, 0, 0)$ |
| $(\bigvee_{i=1}^k \bar{E}_i H_i)(\bigwedge_{i=k+1}^n E_i H_i)$ | 0 | 1 | 0 | $(0, 1, 0)$ |
| $(\bigvee_{i=1}^k \bar{E}_i H_i)(\bigvee_{i=k+1}^n \bar{E}_i H_i)$ | 0 | 0 | 0 | $(0, 0, 0)$ |
| $(\bigwedge_{i \notin S''} E_i H_i)(\bigwedge_{i \in S''} \bar{H}_i)$ | 1 | $x_{S''}$ | $x_{S''}$ | $(1, x_{S''}, x_{S''})$ |
| $(\bigvee_{i=1}^k \bar{E}_i H_i)(\bigwedge_{i \in \{k+1, \dots, n\} \setminus S''} E_i H_i)(\bigwedge_{i \in S''} \bar{H}_i)$ | 0 | $x_{S''}$ | 0 | $(0, x_{S''}, 0)$ |
| $(\bigwedge_{i \notin S'} E_i H_i)(\bigwedge_{i \in S'} \bar{H}_i)$ | $x_{S'}$ | 1 | $x_{S'}$ | $(x_{S'}, 1, x_{S'})$ |
| $(\bigvee_{i=k+1}^n \bar{E}_i H_i)(\bigwedge_{i \in \{1, \dots, k\} \setminus S'} E_i H_i)(\bigwedge_{i \in S'} \bar{H}_i)$ | $x_{S'}$ | 0 | 0 | $(x_{S'}, 0, 0)$ |
| $(\bigwedge_{i \notin S' \cup S''} E_i H_i)(\bigwedge_{i \in S' \cup S''} \bar{H}_i)$ | $x_{S'}$ | $x_{S''}$ | $x_{S' \cup S''}$ | $(x_{S'}, x_{S''}, x_{S' \cup S''})$ |
| $\bigwedge_{i=1}^n \bar{H}_i$ | $x_{1\dots k}$ | $x_{k+1\dots n}$ | $x_{1\dots n}$ | $(x_{1\dots k}, x_{k+1\dots n}, x_{1\dots n})$ |

Table 1: Possible values Q_h 's of the random vector $(\mathcal{C}_{1\dots k}, \mathcal{C}_{k+1\dots n}, \mathcal{C}_{1\dots n})$, where $\emptyset \neq S' \subseteq \{1, \dots, k\}$, $\emptyset \neq S'' \subseteq \{k+1, \dots, n\}$, $S' \cup S'' \neq \{1, \dots, n\}$, and $(x_{1\dots k}, x_{k+1\dots n}, x_{1\dots n}) = Q_0 = \mathcal{M}_3$.

denote by \mathcal{T} the tetrahedron with vertices $(1, 1, 1)$, $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 0)$, that is

$$\mathcal{T} = \{(x, y, z) : (x, y) \in [0, 1]^2, \max\{0, x + y - 1\} \leq z \leq \min\{x, y\}\}.$$

We observe that \mathcal{T} is the convex hull of $(1, 1, 1)$, $(1, 0, 0)$, $(0, 1, 0)$, $(0, 0, 0)$. We also observe that the points $(1, x_{S''}, x_{S''})$, $(0, x_{S''}, 0)$, $(x_{S'}, 1, x_{S'})$, $(x_{S'}, 0, 0)$ belong to \mathcal{T} because

$$\begin{aligned} (1, x_{S''}, x_{S''}) &= x_{S''}(1, 1, 1) + (1 - x_{S''})(1, 0, 0), & (0, x_{S''}, 0) &= x_{S''}(0, 1, 0) + (1 - x_{S''})(0, 0, 0), \\ (x_{S'}, 1, x_{S'}) &= x_{S'}(1, 1, 1) + (1 - x_{S'})(0, 1, 0), & (x_{S'}, 0, 0) &= x_{S'}(1, 0, 0) + (1 - x_{S'})(0, 0, 0). \end{aligned}$$

We recall that coherence of \mathcal{M} implies coherence of the sub-assessment (x_i, x_j, x_{ij}) , with $i \neq j$, on the sub-family $\{E_i|H_i, E_j|H_j, \mathcal{C}_{ij}\}$. By formula (6), the coherence of (x_i, x_j, x_{ij}) amounts to the condition $(x_i, x_j, x_{ij}) \in \mathcal{T}$. Now, let us assume by induction that the point $(x_{S'}, x_{S''}, x_{S' \cup S''})$ belongs to \mathcal{T} , for every pair of nonempty subsets $S' \subseteq \{1, \dots, k\}$, $S'' \subseteq \{k+1, \dots, n\}$, with $S' \cup S'' \subset \{1, \dots, n\}$. Under this inductive hypothesis, the convex hull of the points Q_h 's, with $Q_h \neq Q_0$, is the tetrahedron \mathcal{T} . Coherence of \mathcal{M}_3 requires that \mathcal{M}_3 belongs to the convex hulls of all the points Q_h 's ($h \neq 0$), that is $\mathcal{M}_3 \in \mathcal{T}$. Then, the inequalities

$$\max\{0, x_{1\dots k} + x_{k+1\dots n} - 1\} \leq x_{1\dots n} \leq \min\{x_{1\dots k}, x_{k+1\dots n}\},$$

are satisfied. \square

Remark 5. Given the conjunction $\mathcal{C}_{1\dots n}$ of n conditional events and a further conditional event $E_{n+1}|H_{n+1}$, it holds that

$$\mathcal{C}_{1\dots n+1} = \mathcal{C}_{1\dots n} \wedge (E_{n+1}|H_{n+1}) = \begin{cases} \mathcal{C}_{1\dots n}, & \text{if } E_{n+1}H_{n+1} \text{ is true,} \\ 0, & \text{if } \bar{E}_{n+1}H_{n+1} \text{ is true,} \\ x_{S \cup \{n+1\}}, & \text{if } (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_i H_i) \wedge \bar{H}_{n+1} \text{ is true.} \end{cases} \quad (12)$$

In particular

$$\mathcal{C}_{1\dots n} \wedge 0 = 0, \quad \mathcal{C}_{1\dots n} \wedge 1 = \mathcal{C}_{1\dots n}. \quad (13)$$

Indeed, if $E_{n+1}H_{n+1} = \emptyset$, it holds that $P(E_{n+1}|H_{n+1}) = x_{n+1} = 0$ and hence $E_{n+1}|H_{n+1} = E_{n+1}H_{n+1} + x_{n+1}\bar{H}_{n+1} = 0$. As, by (9), $\mathcal{C}_{1\dots n+1} \leq E_{n+1}|H_{n+1} = 0$, it follows that $\mathcal{C}_{1\dots n+1} = \mathcal{C}_{1\dots n} \wedge 0 = 0$.

If $H_{n+1} \subseteq E_{n+1}$, i.e., $E_{n+1}H_{n+1} = H_{n+1}$, it holds that $x_{n+1} = 1$ and hence $E_{n+1}|H_{n+1} = E_{n+1}H_{n+1} + x_{n+1}\bar{H}_{n+1} = H_{n+1} + \bar{H}_{n+1} = 1$; then (12) becomes

$$\mathcal{C}_{1\dots n+1} = \begin{cases} \mathcal{C}_{1\dots n}, & \text{if } H_{n+1} \text{ is true,} \\ x_{S \cup \{n+1\}}, & \text{if } (\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_i H_i) \wedge \bar{H}_{n+1} \text{ is true.} \end{cases}$$

For every nonempty subset $S \subseteq \{1, \dots, n\}$, by Theorem 6 it holds that

$$\max\{0, x_S + x_{n+1} - 1\} = x_S \leq x_{S \cup \{n+1\}} \leq x_S = \min\{x_S, x_{n+1}\}.$$

Then, $x_{S \cup \{n+1\}} = x_S$, which is the value of $\mathcal{C}_{1\dots n}$ when $(\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_i H_i)$ is true. Thus $\mathcal{C}_{1\dots n+1} = \mathcal{C}_{1\dots n} \wedge 1 = \mathcal{C}_{1\dots n}$.

Concerning the decomposition formula, we first examine the case $n = 1$ (see also [53, Proposition 1]). We recall that, given a conditional event $\mathcal{C}_1 = E_1|H_1$, we denote its indicator by the same symbol. Then, given a further conditional event $E_2|H_2$, we show that (the indicator) \mathcal{C}_1 can be decomposed as the sum of the conjunctions $\mathcal{C}_{12} = (E_1|H_1) \wedge (E_2|H_2)$ and $\mathcal{C}_{1\bar{2}} = (E_1|H_1) \wedge (\bar{E}_2|H_2)$. We set $P(E_1|H_1) = x_1$, $P(E_2|H_2) = x_2$, $\mathbb{P}(\mathcal{C}_{12}) = x_{12}$, $\mathbb{P}(\mathcal{C}_{1\bar{2}}) = x_{1\bar{2}}$. The next result shows the decomposition of \mathcal{C}_1 .

Theorem 7. The conditionals $\mathcal{C}_1, \mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}$ satisfy the relation

$$\mathcal{C}_1 = \mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}. \quad (14)$$

Proof. Table 2 shows, under logical independence of the events E_1, E_2, H_1, H_2 , the possible values for the random vector $(\mathcal{C}_1, \mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{12} + \mathcal{C}_{1\bar{2}})$ associated with the constituents C_h 's generated by the family $\{\mathcal{C}_1, \mathcal{C}_2\}$. We observe that both \mathcal{C}_{12} and $\mathcal{C}_{1\bar{2}}$ are conditional random quantities with the

| | C_h | \mathcal{C}_1 | \mathcal{C}_{12} | $\mathcal{C}_{1\bar{2}}$ | $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$ |
|-------|-------------------------------|-----------------|--------------------|--------------------------|---|
| C_1 | $E_1 H_1 E_2 H_2$ | 1 | 1 | 0 | 1 |
| C_2 | $E_1 H_1 \bar{E}_2 H_2$ | 1 | 0 | 1 | 1 |
| C_3 | $\bar{E}_1 H_1 E_2 H_2$ | 0 | 0 | 0 | 0 |
| C_4 | $\bar{E}_1 H_1 \bar{E}_2 H_2$ | 0 | 0 | 0 | 0 |
| C_5 | $\bar{H}_1 E_2 H_2$ | x_1 | x_1 | 0 | x_1 |
| C_6 | $\bar{H}_1 \bar{E}_2 H_2$ | x_1 | 0 | x_1 | x_1 |
| C_7 | $E_1 H_1 \bar{H}_2$ | 1 | x_2 | $1 - x_2$ | 1 |
| C_8 | $\bar{E}_1 H_1 \bar{H}_2$ | 0 | 0 | 0 | 0 |
| C_0 | $\bar{H}_1 \bar{H}_2$ | x_1 | x_{12} | $x_{1\bar{2}}$ | $x_{12} + x_{1\bar{2}}$ |

Table 2: Numerical values of the random vector $(\mathcal{C}_1, \mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{12} + \mathcal{C}_{1\bar{2}})$.

same conditioning event $H_1 \vee H_2$ and hence $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$ is still a conditional random quantity with conditioning event $H_1 \vee H_2$. As shown in Table 2, for each $C_h \subseteq H_1 \vee H_2$ (i.e., $h = 1, \dots, 8$), if C_h is true then \mathcal{C}_1 coincides with $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$. In other words, \mathcal{C}_1 coincides with $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$ when $H_1 \vee H_2$ is true. Thus, by Theorem 3, it holds that

$$x_1 = \mathbb{P}(\mathcal{C}_1) = \mathbb{P}(\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}) = \mathbb{P}(\mathcal{C}_{12}) + \mathbb{P}(\mathcal{C}_{1\bar{2}}) = x_{12} + x_{1\bar{2}};$$

then \mathcal{C}_1 coincides with $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$ when C_0 is true. Therefore \mathcal{C}_1 and $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$ coincide in all cases; that is $\mathcal{C}_1 = \mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$. In case of some logical dependencies, some constituent C_h may be impossible; but, of course, the relation $\mathcal{C}_1 = \mathcal{C}_{12} + \mathcal{C}_{1\bar{2}}$ is still valid. \square

By the same reasoning, $\mathcal{C}_{\bar{1}} = \mathcal{C}_{\bar{1}2} + \mathcal{C}_{\bar{1}\bar{2}}$, where $\mathcal{C}_{\bar{1}2} = \mathcal{C}_{\bar{1}} \wedge \mathcal{C}_2$, $\mathcal{C}_{\bar{1}\bar{2}} = \mathcal{C}_{\bar{1}} \wedge \mathcal{C}_{\bar{2}}$.

We observe that by Remark 4, given $n + 1$ conditional events $E_1|H_1, \dots, E_{n+1}|H_{n+1}$, their conjunction $\mathcal{C}_{1\dots n+1}$ coincides with $\mathcal{C}_{1\dots n} \wedge (E_{n+1}|H_{n+1})$. Likewise, $\mathcal{C}_{1\dots n\bar{n+1}}$ coincides with $\mathcal{C}_{1\dots n} \wedge (\bar{E}_{n+1}|H_{n+1})$. The next result shows the decomposition for the conjunction of n conditional events.

Theorem 8. Let $n + 1$ conditional events $E_1|H_1, \dots, E_{n+1}|H_{n+1}$ be given. It holds that

$$\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n+1}}. \quad (15)$$

Proof. We recall that $x_{1\dots n} = \mathbb{P}(\mathcal{C}_{1\dots n})$, $x_{n+1} = P(E_{n+1}|H_{n+1})$, $x_{1\dots n+1} = \mathbb{P}(\mathcal{C}_{1\dots n+1})$, and $x_{1\dots n\bar{n+1}} = \mathbb{P}(\mathcal{C}_{1\dots n\bar{n+1}})$. Moreover, given any nonempty strict subset $S = \{i_1, \dots, i_k\}$ of $\{1, 2, \dots, n\}$, we set

$$\mathcal{C}_S = \mathcal{C}_{i_1 \dots i_k} = \bigwedge_{j \in S} (E_j|H_j), \quad \mathcal{C}_{S \cup \{n+1\}} = \mathcal{C}_S \wedge E_{n+1}|H_{n+1}, \quad \mathcal{C}_{S \cup \{\bar{n+1}\}} = \mathcal{C}_S \wedge \bar{E}_{n+1}|H_{n+1},$$

and

$$x_S = \mathbb{P}(\mathcal{C}_S), \quad x_{S \cup \{n+1\}} = \mathbb{P}(\mathcal{C}_{S \cup \{n+1\}}), \quad x_{S \cup \{\bar{n}+1\}} = \mathbb{P}(\mathcal{C}_{S \cup \{\bar{n}+1\}}).$$

We prove the theorem by induction on the cardinality of S , denoted by s . By Theorem 7 the equality (15) holds for $n = 1$. We assume that (15) holds for each integer $s < n$, that is: $\mathcal{C}_S = \mathcal{C}_{S \cup \{n+1\}} + \mathcal{C}_{S \cup \{\bar{n}+1\}}$; then, we prove that (15) holds for $s = n$, that is: $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$. We first assume logical independence of the events $E_i, H_i, i = 1, \dots, n+1$. We distinguish the following cases: (i) $E_{n+1}H_{n+1}$ true; (ii) $\bar{E}_{n+1}H_{n+1}$ true; (iii) \bar{H}_{n+1} true.

Case (i). From (12) it holds that $\mathcal{C}_{1\dots n+1} = \mathcal{C}_{1\dots n}$ and $\mathcal{C}_{1\dots n\bar{n}+1} = 0$, so that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$.

Case (ii). From (12) it holds that $\mathcal{C}_{1\dots n+1} = 0$ and $\mathcal{C}_{1\dots n\bar{n}+1} = \mathcal{C}_{1\dots n}$, so that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$.

Case (iii). We distinguish the following subcases: (a) $\bigwedge_{i=1}^n E_i H_i$ true; (b) $\bigvee_{i=1}^n \bar{E}_i H_i$ true; (c) $(\bigwedge_{i \in S} \bar{H}_i) \wedge (\bigwedge_{i \notin S} E_i H_i)$ true, for some nonempty $S \subset \{1, \dots, n\}$; (d) $\bigwedge_{i=1}^{n+1} \bar{H}_i$ true.

In the subcase (a) it holds that $\mathcal{C}_{1\dots n} = 1$, $\mathcal{C}_{1\dots n+1} = x_{n+1}$, and $\mathcal{C}_{1\dots n\bar{n}+1} = 1 - x_{n+1}$; so that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$.

In the subcase (b) it holds that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} = \mathcal{C}_{1\dots n\bar{n}+1} = 0$; so that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$.

In the subcase (c) it holds that $\mathcal{C}_{1\dots n} = x_S$, $\mathcal{C}_{1\dots n+1} = x_{S \cup \{n+1\}}$, and $\mathcal{C}_{1\dots n\bar{n}+1} = x_{S \cup \{\bar{n}+1\}}$. By the inductive hypothesis it follows that $x_S = x_{S \cup \{n+1\}} + x_{S \cup \{\bar{n}+1\}}$, so that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$.

In the subcase (d) it holds that $\mathcal{C}_{1\dots n} = x_{1\dots n}$, $\mathcal{C}_{1\dots n+1} = x_{1\dots n+1}$, and $\mathcal{C}_{1\dots n\bar{n}+1} = x_{1\dots n\bar{n}+1}$. We observe that $\mathcal{C}_{1\dots n}$ is a conditional random quantity with conditioning event $H_1 \vee \dots \vee H_n$. Moreover, both $\mathcal{C}_{1\dots n+1}$ and $\mathcal{C}_{1\dots n\bar{n}+1}$ are conditional random quantities with the same conditioning event $H_1 \vee \dots \vee H_{n+1}$ and hence $\mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$ is still a conditional random quantity with conditioning event $H_1 \vee \dots \vee H_{n+1}$. Finally, we observe that $\mathcal{C}_{1\dots n}$ and $\mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$ coincide when $H_1 \vee \dots \vee H_{n+1}$ is true. Then, by applying Theorem 3 with $X|H = \mathcal{C}_{1\dots n}$ and $Y|K = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$, it holds that $x_{1\dots n} = x_{1\dots n+1} + x_{1\dots n\bar{n}+1}$, so that $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$. In conclusion, $\mathcal{C}_{1\dots n}$ and $\mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$ coincide in all cases; that is $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$ (see also Table 3). In case of some logical dependencies, some constituent C_h may be impossible; but, of course, the relation $\mathcal{C}_{1\dots n} = \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$ is still valid. \square

| C_h | $\mathcal{C}_{1\dots n}$ | $\mathcal{C}_{1\dots n+1}$ | $\mathcal{C}_{1\dots n\bar{n}+1}$ | $\mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$ |
|--|--------------------------|----------------------------|-----------------------------------|--|
| $E_{n+1}H_{n+1}$ | $\mathcal{C}_{1\dots n}$ | $\mathcal{C}_{1\dots n}$ | 0 | $\mathcal{C}_{1\dots n}$ |
| $\bar{E}_{n+1}H_{n+1}$ | $\mathcal{C}_{1\dots n}$ | 0 | $\mathcal{C}_{1\dots n}$ | $\mathcal{C}_{1\dots n}$ |
| $(\bigwedge_{i=1}^n E_i H_i) \bar{H}_{n+1}$ | 1 | x_{n+1} | $1 - x_{n+1}$ | 1 |
| $(\bigvee_{i=1}^n \bar{E}_i H_i) \bar{H}_{n+1}$ | 0 | 0 | 0 | 0 |
| $(\bigwedge_{i \in S} \bar{H}_i \bigwedge_{i \notin S} E_i H_i) \bar{H}_{n+1}$ | x_S | $x_{S \cup \{n+1\}}$ | $x_{S \cup \{\bar{n}+1\}}$ | x_S |
| $\bigwedge_{i=1}^{n+1} \bar{H}_i$ | $x_{1\dots n}$ | $x_{1\dots n+1}$ | $x_{1\dots n\bar{n}+1}$ | $x_{1\dots n}$ |

Table 3: Numerical values of the conditional random quantities $\mathcal{C}_{1\dots n}, \mathcal{C}_{1\dots n+1}, \mathcal{C}_{1\dots n\bar{n}+1}, \mathcal{C}_{1\dots n+1} + \mathcal{C}_{1\dots n\bar{n}+1}$. Each S is a nonempty strict subset of $\{1, \dots, n\}$.

Given any integer $n \geq 1$ and n conditional events $E_1|H_1, \dots, E_n|H_n$, we set $\mathcal{C}_{1^* \dots n^*} = \bigwedge_{i=1}^n E_i^* | H_i$, where for each index i it holds that $i^* \in \{i, \bar{i}\}$ and $E_i^* = E_i$, or $E_i^* = \bar{E}_i$, according to whether $i^* = i$, or $i^* = \bar{i}$, respectively. In particular $\mathcal{C}_{1^*} = \mathcal{C}_1 = E_1|H_1$ when $1^* = 1$ and

$\mathcal{C}_{1^*} = \mathcal{C}_{\bar{1}} = \bar{E}_1|H_1$ when $1^* = \bar{1}$. Moreover, given any subset $\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}$, by defining $\{i_{h+1}, \dots, i_n\} = \{1, \dots, n\} \setminus \{i_1, \dots, i_h\}$, we set

$$\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = (E_{i_1}|H_{i_1}) \wedge \dots \wedge (E_{i_h}|H_{i_h}) \wedge (\bar{E}_{i_{h+1}}|H_{i_{h+1}}) \wedge \dots \wedge (\bar{E}_{i_n}|H_{i_n}). \quad (16)$$

We recall that by definition the value of $\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$, when the conditioning events H_1, \dots, H_n are all false, is its prevision $\mathbb{P}(\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n})$. We set

$$\mathbb{P}(\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}) = x_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}. \quad (17)$$

Notice that, as the operation of conjunction is commutative, for each conjunction $\mathcal{C}_{1^* \dots n^*}$ it holds that $\mathcal{C}_{1^* \dots n^*} = \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$, for a suitable subset $\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}$. Then, given a further conditional event $E_{n+1}|H_{n+1}$, by the same reasoning of Theorem 8 it holds that

$$\mathcal{C}_{1^* \dots n^*} = \mathcal{C}_{1^* \dots n^* n+1} + \mathcal{C}_{1^* \dots n^* \bar{n+1}}, \quad \forall (1^*, \dots, n^*) \in \{1, \bar{1}\} \times \dots \times \{n, \bar{n}\}, \quad (18)$$

or equivalently

$$\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_{n+1}} + \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n \bar{n+1}}, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}. \quad (19)$$

For instance, it holds that: $\mathcal{C}_{\bar{1}\bar{2}\bar{3}} = \mathcal{C}_{\bar{1}\bar{2}\bar{3}4} + \mathcal{C}_{\bar{1}\bar{2}\bar{3}\bar{4}}$; $\mathcal{C}_{\bar{1}\bar{2}4} = \mathcal{C}_{\bar{1}\bar{2}43} + \mathcal{C}_{\bar{1}\bar{2}\bar{4}3} = \mathcal{C}_{\bar{1}\bar{2}\bar{3}4} + \mathcal{C}_{\bar{1}\bar{2}\bar{3}\bar{4}}$, and so on.

4. The set of conditional constituents

In this section we show that a notion of “constituent”, which we call *conditional constituent*, can be introduced for the case of n conditional events $E_1|H_1, \dots, E_n|H_n$. We recall that, given n (unconditional) events E_1, \dots, E_n and denoting the set of their constituents by $\{C_h, h = 1, \dots, m\}$, where $m \leq 2^n$ (with $m = 2^n$ in case of logical independence), it holds that

$$(i) \quad C_h \wedge C_k = \emptyset, \quad \forall h \neq k; \quad (ii) \quad \bigvee_{h=1}^m C_h = \Omega. \quad (20)$$

In terms of indicators (denoted by the same symbols) formula (20) becomes:

$$(i)' \quad C_h \wedge C_k = 0, \quad \forall h \neq k; \quad (ii)' \quad \sum_{h=1}^m C_h = 1, \quad (21)$$

with $C_h \geq 0$, $h = 1, \dots, m$. Then, it holds that:

$$E_j = \sum_{h: C_h \subseteq E_j} C_h, \quad P(E_j) = \sum_{h: C_h \subseteq E_j} P(C_h); \quad j = 1, \dots, n. \quad (22)$$

We introduce the set of conditional constituents associated with n conditional events, by obtaining some properties which are analogous to those valid for the unconditional events. Indeed, we will show that properties (i)' and (ii)' in (21), still hold if we replace events, and their constituents, by conditional events, and their conditional constituents, respectively. In other words, the conditional constituents are *incompatible* (i.e., their conjunction is 0) and their sum is 1.

Moreover, likewise formula (22), we will show that the indicator of each conditional event, and its prevision, can be decomposed as the sum of suitable conditional constituents, and their previsions, respectively.

In addition, as in the case of unconditional events, the conditional constituents associated with a family of n conditional events $\{E_1|H_1, \dots, E_n|H_n\}$ are all the (non zero) conjunctions $(A_1|H_1) \wedge \dots \wedge (A_n|H_n)$, where $A_i \in \{E_i, \bar{E}_i\}$, $i = 1, \dots, n$.

Definition 7. The set of conditional constituents, or *c-constituents*, associated with a family of n conditional events $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$ is

$$\mathcal{K} = \{\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} : \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}, \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} \neq 0\},$$

where each c-constituent $\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$ is a conjunction, as defined in (16).

Notice that the cardinality of \mathcal{K} is 2^n when the events $E_1, \dots, E_n, H_1, \dots, H_n$ are logically independent. In the presence of some logical dependencies it may be that $\mathcal{C}_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = 0$ for some $\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}$, as shown in the example below. If $\mathcal{C}_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$ coincides with 0, then it is not included in the set \mathcal{K} .

Example 2. Given two logically independent events E, H let us consider the family $\mathcal{E} = \{E_1|H_1, E_2|H_2\}$, where $E_1 = E, E_2 = \bar{E}, H_1 = H_2 = H$. We observe that $\mathcal{K} \subseteq \{\mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{\bar{1}2}, \mathcal{C}_{\bar{1}\bar{2}}\}$, where, by recalling that $E|H = EH + P(E|H)\bar{H}$ and hence $\emptyset|H = 0$, it holds that

$$\mathcal{C}_{12} = (E|H) \wedge (\bar{E}|H) = \emptyset|H = 0 = \mathcal{C}_{\bar{1}\bar{2}}, \mathcal{C}_{1\bar{2}} = (E|H) \wedge (E|H) = E|H, \mathcal{C}_{\bar{1}2} = (\bar{E}|H) \wedge (\bar{E}|H) = \bar{E}|H.$$

As we can see, in this case there are two c-constituents which are not zero; that is: $\mathcal{K} = \{\mathcal{C}_{1\bar{2}}, \mathcal{C}_{\bar{1}2}\} = \{E|H, \bar{E}|H\} = \mathcal{E}$.

In the next result we show that the properties (i)' and (ii)' in (21), relative to unconditional events, still hold for the case of conditional events.

Theorem 9. Given a family of n conditional events $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$, let \mathcal{K} be the set of c-constituents associated with \mathcal{E} . It holds that

$$\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} \wedge \mathcal{C}_{j_1 \dots j_k \bar{j}_{k+1} \dots \bar{j}_n} = \emptyset|(H_1 \vee \dots \vee H_n) = 0, \quad \forall \{i_1, i_2, \dots, i_h\} \neq \{j_1, j_2, \dots, j_k\}. \quad (23)$$

Proof. As $\{i_1, \dots, i_h\} \neq \{j_1, \dots, j_k\}$, the set $(\{i_1, \dots, i_h\} \setminus \{j_1, \dots, j_k\}) \cup (\{j_1, \dots, j_k\} \setminus \{i_1, \dots, i_h\})$ is non empty. Let r be one of its elements. For the sake of simplicity, we assume that $r = i_1 = j_{k+1}$. Then

$$(E_{i_1}|H_{i_1}) \wedge (\bar{E}_{j_{k+1}}|H_{j_{k+1}}) = (E_{i_1}|H_{i_1}) \wedge (\bar{E}_{i_1}|H_{i_1}) = (E_{i_1} \wedge \bar{E}_{i_1})|H_{i_1} = \emptyset|H_{i_1} = 0,$$

and hence

$$\begin{aligned} \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} \wedge \mathcal{C}_{j_1 \dots j_k \bar{j}_{k+1} \dots \bar{j}_n} &= (E_{i_1}|H_{i_1}) \wedge (\bar{E}_{j_{k+1}}|H_{j_{k+1}}) \wedge \mathcal{C}_{i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} \wedge \mathcal{C}_{j_1 \dots j_k \bar{j}_{k+2} \dots \bar{j}_n} = \\ &= (\emptyset|H_{i_1}) \wedge \mathcal{C}_{i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} \wedge \mathcal{C}_{j_1 \dots j_k \bar{j}_{k+2} \dots \bar{j}_n} = \emptyset|(H_1 \vee \dots \vee H_n) = 0. \end{aligned}$$

□

Theorem 10. Given a family of n conditional events $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$, let \mathcal{K} be the set of c-constituents associated with \mathcal{E} . For each $1 \leq k \leq n$, it holds that

$$\sum_{\{i_1, \dots, i_h\} \subseteq \{1, \dots, k\}} \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k} = \sum_{(1^* \dots k^*) \in \{1, \bar{1}\} \times \dots \times \{k, \bar{k}\}} \mathcal{C}_{1^* \dots k^*} = 1, \quad (24)$$

$$\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k} \geq 0, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, k\}.$$

Proof. First of all, as $P(\Omega|H_1) = 1$, we observe that

$$\mathcal{C}_1 + \mathcal{C}_{\bar{1}} = E_1|H_1 + \bar{E}_1|H_1 = (E_1 \vee \bar{E}_1)|H_1 = \Omega|H_1 = 1,$$

that is (24) holds when $k = 1$. Moreover, from (14), $\mathcal{C}_{12} + \mathcal{C}_{1\bar{2}} + \mathcal{C}_{\bar{1}2} + \mathcal{C}_{\bar{1}\bar{2}} = \mathcal{C}_1 + \mathcal{C}_{\bar{1}} = 1$, that is (24) holds when $k = 2$. Then, by induction, assuming (24) valid for $k - 1$, from (18) it follows that

$$\begin{aligned} & \sum_{\{i_1, \dots, i_h\} \subseteq \{1, \dots, k\}} \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k} = \sum_{(1^*, \dots, k^*) \in \{1, \bar{1}\} \times \dots \times \{k, \bar{k}\}} \mathcal{C}_{1^* \dots k^*} = \\ & = \sum_{(1^*, \dots, (k-1)^*) \in \{1, \bar{1}\} \times \dots \times \{k-1, \bar{k-1}\}} (\mathcal{C}_{1^* \dots (k-1)^* k} + \mathcal{C}_{1^* \dots (k-1)^* \bar{k}}) = \\ & = \sum_{(1^*, \dots, (k-1)^*) \in \{1, \bar{1}\} \times \dots \times \{k-1, \bar{k-1}\}} \mathcal{C}_{1^* \dots (k-1)^*} = 1, \end{aligned}$$

that is formula (24) is valid for k . Finally, the inequalities $\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k} \geq 0$, $\forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, k\}$, hold because the prevision assessments used when defining conjunctions are assumed to be coherent. \square

We observe in particular that from Definition 4 and Theorem 10 it holds that

$$\sum_{\{i_1, i_2, \dots, i_h\} \subseteq \{1, 2, \dots, n\}} x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = 1, \quad x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} \geq 0, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}. \quad (25)$$

In other words the prevision of each conditional constituent is nonnegative and the sum of all these previsions is equal to 1. In addition, we show that the properties in (22) still hold for conditional events. Indeed, by Theorem 8, it follows that

$$\begin{aligned} \mathcal{C}_1 &= \mathcal{C}_{12} + \mathcal{C}_{1\bar{2}} = \mathcal{C}_{123} + \mathcal{C}_{12\bar{3}} + \mathcal{C}_{\bar{1}23} + \mathcal{C}_{\bar{1}2\bar{3}} = \dots = \\ &= \mathcal{C}_{12 \dots n} + \mathcal{C}_{12 \dots n-1 \bar{n}} + \dots + \mathcal{C}_{1\bar{2} \dots n-1 n} + \mathcal{C}_{1\bar{2} \dots \bar{n}} = \\ &= \sum_{(2^*, \dots, n^*) \in \{2, \bar{2}\} \times \dots \times \{n, \bar{n}\}} \mathcal{C}_{12^* \dots n^*} = \sum_{\{1\} \subseteq \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}, \end{aligned}$$

and in general, for each $j \in \{1, \dots, n\}$ it holds that

$$\mathcal{C}_j = \sum_{\{(1^*, \dots, (j-1)^*, (j+1)^*, \dots, n^*)\}} \mathcal{C}_{1^* \dots (j-1)^* j (j+1)^* \dots n^*} = \sum_{\{j\} \subseteq \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} \mathcal{C}_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}, \quad (26)$$

where the symbol $\{(1^*, \dots, (j-1)^*, (j+1)^*, \dots, n^*)\}$ denotes the following set

$$\{(1^*, \dots, (j-1)^*, (j+1)^*, \dots, n^*) \in \{1, \bar{1}\} \times \dots \times \{j-1, \bar{j-1}\} \times \{j+1, \bar{j+1}\} \times \dots \times \{n, \bar{n}\}.$$

Moreover, concerning the probability x_j of $E_j|H_j$, from (26) it holds that

$$\begin{aligned} P(\mathcal{C}_j) &= x_j = \sum_{\{(1^*, \dots, (j-1)^*, (j+1)^*, \dots, n^*)\}} x_{1^* \dots (j-1)^* j (j+1)^* \dots n^*} = \\ &= \sum_{\{j\} \subseteq \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} x_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = \sum_{\{j\} \subseteq \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} \mathbb{P}(\mathcal{C}_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}). \end{aligned} \quad (27)$$

More in general, for the conjunction $\mathcal{C}_S = \bigwedge_{j \in S} (E_j|H_j)$ it holds that

$$\mathcal{C}_S = \sum_{\{i_1, \dots, i_h\} \supseteq S} \mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}, \quad \forall \emptyset \neq S \subseteq \{1, \dots, n\}, \quad (28)$$

and hence

$$\mathbb{P}(\mathcal{C}_S) = x_S = \sum_{\{i_1, \dots, i_h\} \supseteq S} x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}, \quad \forall \emptyset \neq S \subseteq \{1, \dots, n\}. \quad (29)$$

Moreover,

$$\mathcal{C}_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k} = \sum_{\{j_1, \dots, j_r\} \subseteq \{i_{k+1}, \dots, i_n\}} \mathcal{C}_{i_1 i_2 \dots i_h j_1 \dots j_r \bar{i}_{h+1} \dots \bar{i}_k \bar{j}_{r+1} \dots \bar{j}_{n-k-r}}, \quad \forall 0 \leq h \leq k \leq n, \quad (30)$$

and for its prevision $x_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k}$ it holds that

$$x_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k} = \sum_{\{j_1, \dots, j_r\} \subseteq \{i_{k+1}, \dots, i_n\}} x_{i_1 i_2 \dots i_h j_1 \dots j_r \bar{i}_{h+1} \dots \bar{i}_k \bar{j}_{r+1} \dots \bar{j}_{n-k-r}}, \quad \forall 0 \leq h \leq k \leq n. \quad (31)$$

5. The inclusion-exclusion principle and the distributivity property

In this section we show that the well known inclusion-exclusion formula, which holds for the disjunction of n unconditional events (and its probability), still holds for the disjunction of n conditional events (and its prevision). This result, and other related formulas, will be used in Section 6. We also prove a distributivity property by means of which we can directly derive the inclusion-exclusion formula. We first give a preliminary result.

Theorem 11. Given $n + 1$ conditional events $E_1 | H_1, \dots, E_{n+1} | H_{n+1}$, it holds that

$$\mathcal{C}_{\bar{1} \dots \bar{h} i_1 \dots i_k n+1} = \mathcal{C}_{i_1 \dots i_k n+1} - \sum_{j=1}^h \mathcal{C}_{j i_1 \dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h} \mathcal{C}_{j_1 j_2 i_1 \dots i_k n+1} + \dots + (-1)^h \mathcal{C}_{1 \dots h i_1 \dots i_k n+1}, \quad 1 \leq h \leq n, \quad \{i_1, \dots, i_k\} \subseteq \{h+1, \dots, n\}. \quad (32)$$

Proof. Formula (32) is satisfied for $h = 1$ because, by the decomposition formula (19), it holds that

$$\mathcal{C}_{\bar{1} i_1 \dots i_k n+1} = \mathcal{C}_{i_1 \dots i_k n+1} - \mathcal{C}_{1 i_1 \dots i_k n+1}, \quad \forall \{i_1, \dots, i_k\} \subseteq \{2, \dots, n\}.$$

By assuming that (32) is satisfied for $h \leq n - 1$, we prove that (32) is also satisfied for $h + 1$. By (19), it holds that

$$\mathcal{C}_{\bar{1} \dots \bar{h} h+1 i_1 \dots i_k n+1} = \mathcal{C}_{\bar{1} \dots \bar{h} h+1 i_1 \dots i_k n+1} + \mathcal{C}_{\bar{1} \dots \bar{h} h+1 i_1 \dots i_k n+1}, \quad \forall \{i_1, \dots, i_k\} \subseteq \{h+2, \dots, n\}.$$

Moreover, by the hypothesis, it holds that

$$\mathcal{C}_{\bar{1} \dots \bar{h} i_1 \dots i_k n+1} = \mathcal{C}_{i_1 \dots i_k n+1} - \sum_{j=1}^h \mathcal{C}_{j i_1 \dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h} \mathcal{C}_{j_1 j_2 i_1 \dots i_k n+1} + \dots + (-1)^h \mathcal{C}_{1 \dots h i_1 \dots i_k n+1}$$

and

$$\mathcal{C}_{\bar{1} \dots \bar{h} h+1 i_1 \dots i_k n+1} = \mathcal{C}_{h+1 i_1 \dots i_k n+1} - \sum_{j=1}^h \mathcal{C}_{j h+1 i_1 \dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h} \mathcal{C}_{j_1 j_2 h+1 i_1 \dots i_k n+1} + \dots + (-1)^h \mathcal{C}_{1 \dots h h+1 i_1 \dots i_k n+1}.$$

Then, by (19), for all $\{i_1, \dots, i_k\} \subseteq \{h+2, \dots, n\}$ it follows that

$$\begin{aligned} \mathcal{C}_{\bar{1}\dots\bar{h+1}i_1\dots i_k n+1} &= \mathcal{C}_{\bar{1}\dots\bar{h}i_1\dots i_k n+1} - \mathcal{C}_{\bar{1}\dots\bar{h}h+1i_1\dots i_k n+1} = \\ &= [\mathcal{C}_{i_1\dots i_k n+1} - \sum_{j=1}^h \mathcal{C}_{ji_1\dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h} \mathcal{C}_{j_1 j_2 i_1\dots i_k n+1} + \dots + (-1)^h \mathcal{C}_{1\dots h i_1\dots i_k n+1}] + \\ &\quad - [\mathcal{C}_{h+1 i_1\dots i_k n+1} - \sum_{j=1}^h \mathcal{C}_{jh+1 i_1\dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h} \mathcal{C}_{j_1 j_2 h+1 i_1\dots i_k n+1} + \dots + (-1)^h \mathcal{C}_{1\dots h+1 i_1\dots i_k n+1}] = \\ &= \mathcal{C}_{i_1\dots i_k n+1} - \sum_{j=1}^{h+1} \mathcal{C}_{ji_1\dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h+1} \mathcal{C}_{j_1 j_2 i_1\dots i_k n+1} + \dots + (-1)^{h+1} \mathcal{C}_{1\dots h+1 i_1\dots i_k n+1}. \end{aligned}$$

Then, formula (32) follows by iterating the previous reasoning from $h = 0$ to $h = n - 1$. \square

We give below some examples where formula (32) is obtained.

$$\begin{aligned} \mathcal{C}_{\bar{1}2} &= \mathcal{C}_2 - \mathcal{C}_{12}, \quad \mathcal{C}_{\bar{1}23} = \mathcal{C}_{23} - \mathcal{C}_{123}, \quad \mathcal{C}_{\bar{1}\bar{2}3} = \mathcal{C}_{\bar{1}3} - \mathcal{C}_{\bar{1}23} = \mathcal{C}_3 - \mathcal{C}_{13} - \mathcal{C}_{23} + \mathcal{C}_{123}, \\ \mathcal{C}_{\bar{1}\bar{2}34} &= \mathcal{C}_{\bar{1}34} - \mathcal{C}_{\bar{1}234} = \mathcal{C}_{34} - \mathcal{C}_{134} - \mathcal{C}_{234} + \mathcal{C}_{1234}, \\ \mathcal{C}_{\bar{1}\bar{2}\bar{3}4} &= \mathcal{C}_{\bar{1}\bar{2}4} - \mathcal{C}_{\bar{1}\bar{2}34} = \mathcal{C}_{\bar{1}4} - \mathcal{C}_{\bar{1}24} - \mathcal{C}_{\bar{1}34} + \mathcal{C}_{\bar{1}234} = \mathcal{C}_4 - \mathcal{C}_{14} - \mathcal{C}_{24} + \mathcal{C}_{124} - \mathcal{C}_{34} + \mathcal{C}_{134} + \mathcal{C}_{234} - \mathcal{C}_{1234}. \end{aligned}$$

In the next result we obtain the inclusion-exclusion formula for the disjunction of n conditional events.

Theorem 12. Given n conditional events $E_1|H_1, \dots, E_n|H_n$, it holds that

$$\mathcal{D}_{1\dots n} = \sum_{h=1}^n (-1)^{h+1} \sum_{1 \leq i_1 < \dots < i_h \leq n} \mathcal{C}_{i_1\dots i_h} = \sum_{i=1}^n \mathcal{C}_i - \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \dots + (-1)^{n+1} \mathcal{C}_{1\dots n}.$$

Proof. From (13), it holds that $\mathcal{C}_{\bar{1}\dots\bar{n}} \wedge 1 = \mathcal{C}_{\bar{1}\dots\bar{n}}$. Then, by applying (32) with $h = n$ and $\mathcal{C}_{n+1} = 1$, it follows that

$$\mathcal{C}_{\bar{1}\dots\bar{n}} = 1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \dots + (-1)^n \mathcal{C}_{1\dots n}. \quad (33)$$

Finally, by recalling (11), we obtain

$$\mathcal{D}_{1\dots n} = 1 - \mathcal{C}_{\bar{1}\dots\bar{n}} = \sum_{i=1}^n \mathcal{C}_i - \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \dots + (-1)^{n+1} \mathcal{C}_{1\dots n}.$$

\square

In the next result we prove the validity of a suitable distributivity property.

Theorem 13. Let $\mathcal{C}_1, \dots, \mathcal{C}_{n+1}$ be $n+1$ conditional events. Then, the following distributivity property is satisfied:

$$\begin{aligned} &[1 - \sum_{i=1}^h \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq h} \mathcal{C}_{i_1 i_2} + \dots + (-1)^h \mathcal{C}_{1\dots h}] \wedge \mathcal{C}_{i_1\dots i_k n+1} = \\ &= 1 \wedge \mathcal{C}_{i_1\dots i_k n+1} - \sum_{i=1}^h \mathcal{C}_i \wedge \mathcal{C}_{i_1\dots i_k n+1} + \sum_{1 \leq i_1 < i_2 \leq h} \mathcal{C}_{i_1 i_2} \wedge \mathcal{C}_{i_1\dots i_k n+1} + \dots + (-1)^h \mathcal{C}_{1\dots h} \wedge \mathcal{C}_{i_1\dots i_k n+1}, \\ &1 \leq h \leq n, \quad \{i_1, \dots, i_k\} \subseteq \{h+1, \dots, n\}. \end{aligned} \quad (34)$$

Proof. By recalling Remark 4, formulas (32), and (33), it holds that

$$\begin{aligned}
& [1 - \sum_{i=1}^h \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq h} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^h \mathcal{C}_{1 \dots h}] \wedge \mathcal{C}_{i_1 \dots i_k n+1} = \mathcal{C}_{\bar{1} \dots \bar{h}} \wedge \mathcal{C}_{i_1 \dots i_k n+1} = \mathcal{C}_{\bar{1} \dots \bar{h} i_1 \dots i_k n+1} = \\
& = \mathcal{C}_{i_1 \dots i_k n+1} - \sum_{j=1}^h \mathcal{C}_{j i_1 \dots i_k n+1} + \sum_{1 \leq j_1 < j_2 \leq h} \mathcal{C}_{j_1 j_2 i_1 \dots i_k n+1} + \cdots + (-1)^h \mathcal{C}_{1 \dots h i_1 \dots i_k n+1} = \\
& = 1 \wedge \mathcal{C}_{i_1 \dots i_k n+1} - \sum_{i=1}^h \mathcal{C}_i \wedge \mathcal{C}_{i_1 \dots i_k n+1} + \sum_{1 \leq i_1 < i_2 \leq h} \mathcal{C}_{i_1 i_2} \wedge \mathcal{C}_{i_1 \dots i_k n+1} + \cdots + (-1)^h \mathcal{C}_{1 \dots h} \wedge \mathcal{C}_{i_1 \dots i_k n+1}.
\end{aligned}$$

□

The next result shows a further aspect of the distributivity property.

Theorem 14. Let $\mathcal{C}_1, \dots, \mathcal{C}_{n+1}$ be $n + 1$ conditional events. Then,

$$\begin{aligned}
& [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge (1 - \mathcal{C}_{n+1}) = \\
& = [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge 1 + \\
& - [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge \mathcal{C}_{n+1}.
\end{aligned} \tag{35}$$

Proof. We observe that, by Theorem 13, when $h = n$ it holds that

$$\begin{aligned}
& [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge \mathcal{C}_{n+1} = \\
& = 1 \wedge \mathcal{C}_{n+1} - \sum_{i=1}^n \mathcal{C}_i \wedge \mathcal{C}_{n+1} + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} \wedge \mathcal{C}_{n+1} + \cdots + (-1)^n \mathcal{C}_{1 \dots n} \wedge \mathcal{C}_{n+1}.
\end{aligned} \tag{36}$$

In particular, if $\mathcal{C}_{n+1} = 1$ it follows that

$$\begin{aligned}
& [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge 1 = \\
& = 1 \wedge 1 - \sum_{i=1}^n \mathcal{C}_i \wedge 1 + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} \wedge 1 + \cdots + (-1)^n \mathcal{C}_{1 \dots n} \wedge 1.
\end{aligned} \tag{37}$$

Based on (33), (36), and (37) it follows that

$$\begin{aligned}
& [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge (1 - \mathcal{C}_{n+1}) = \mathcal{C}_{\bar{1} \dots \bar{n}} \wedge \mathcal{C}_{n+1} = \\
& = \mathcal{C}_{\bar{1} \dots \bar{n+1}} = 1 - \sum_{i=1}^{n+1} \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n+1} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^{n+1} \mathcal{C}_{1 \dots n+1} = \\
& = [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] + \\
& - [\mathcal{C}_{n+1} - \sum_{i=1}^n \mathcal{C}_{i n+1} + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2 n+1} + \cdots + (-1)^{n+1} \mathcal{C}_{1 \dots n+1}] = \\
& = [1 \wedge 1 - \sum_{i=1}^n \mathcal{C}_i \wedge 1 + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} \wedge 1 + \cdots + (-1)^n \mathcal{C}_{1 \dots n} \wedge 1] + \\
& - [1 \wedge \mathcal{C}_{n+1} - \sum_{i=1}^n \mathcal{C}_i \wedge \mathcal{C}_{n+1} + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} \wedge \mathcal{C}_{n+1} + \cdots + (-1)^{n+1} \mathcal{C}_{1 \dots n} \wedge \mathcal{C}_{n+1}] = \\
& = [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge 1 + \\
& - [1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}] \wedge \mathcal{C}_{n+1}.
\end{aligned}$$

□

We remark that, by the relation $\mathcal{D}_{1 \dots n} = 1 - \mathcal{C}_{\bar{1} \dots \bar{n}}$, the inclusion-exclusion formula also follows by directly computing $\mathcal{C}_{\bar{1} \dots \bar{n}}$ by means of the distributivity property, as shown below.

$$\begin{aligned}
& \mathcal{C}_{\bar{1} \dots \bar{n}} = \bigwedge_{i=1}^n \mathcal{C}_{\bar{i}} = \bigwedge_{i=1}^n (1 - \mathcal{C}_i) = (1 - \mathcal{C}_1 - \mathcal{C}_2 + \mathcal{C}_{12}) \wedge (1 - \mathcal{C}_3) \wedge \cdots \wedge (1 - \mathcal{C}_n) = \\
& = (1 - \mathcal{C}_1 - \mathcal{C}_2 - \mathcal{C}_3 + \mathcal{C}_{12} + \mathcal{C}_{13} + \mathcal{C}_{23} - \mathcal{C}_{123}) \wedge (1 - \mathcal{C}_4) \wedge \cdots \wedge (1 - \mathcal{C}_n) = \\
& = \cdots = 1 - \sum_{i=1}^n \mathcal{C}_i + \sum_{1 \leq i_1 < i_2 \leq n} \mathcal{C}_{i_1 i_2} + \cdots + (-1)^n \mathcal{C}_{1 \dots n}.
\end{aligned} \tag{38}$$

Then, for each nonempty subset $\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}$, by taking into account (34) it holds that

$$\begin{aligned} \mathcal{C}_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}} &= \mathcal{C}_{\overline{i_{h+1}} \dots \overline{i_n}} \wedge \mathcal{C}_{i_1 \dots i_h} = \\ &= [1 - \sum_{j \in \{i_{h+1}, \dots, i_n\}} \mathcal{C}_j + \sum_{\{j_1, j_2\} \subseteq \{i_{h+1}, \dots, i_n\}} \mathcal{C}_{j_1 j_2} + \dots + (-1)^{n-h} \mathcal{C}_{i_{h+1} \dots i_n}] \wedge \mathcal{C}_{i_1 \dots i_h} = \\ &= \mathcal{C}_{i_1 \dots i_h} - \sum_{j \in \{i_{h+1}, \dots, i_n\}} \mathcal{C}_{i_1 \dots i_h j} + \sum_{\{j_1, j_2\} \subseteq \{i_{h+1}, \dots, i_n\}} \mathcal{C}_{i_1 \dots i_h j_1 j_2} + \dots + (-1)^{n-h} \mathcal{C}_{i_1 \dots i_n}. \end{aligned} \quad (39)$$

When $\{i_1, \dots, i_h\} = \emptyset$, that is $h = 0$, formula (39) continues to hold because, if we set by convention that $\mathcal{C}_{\{i_1, \dots, i_h\}} = \mathcal{C}_{\emptyset} = 1$, it reduces to formula (33). Then, in general, it holds that

$$\mathcal{C}_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}} = \sum_{k=0}^{n-h} (-1)^k \sum_{\{j_1, \dots, j_k\} \subseteq \{i_{h+1}, \dots, i_n\}} \mathcal{C}_{i_1 \dots i_h j_1 \dots j_k}, \quad \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}. \quad (40)$$

Remark 6. We observe that, concerning the probabilistic aspects, by recalling (17) from coherence it holds that

$$\begin{aligned} x_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}} &= x_{i_1 \dots i_h} - \sum_{j \in \{i_{h+1}, \dots, i_n\}} x_{i_1 \dots i_h j} + \sum_{\{j_1, j_2\} \subseteq \{i_{h+1}, \dots, i_n\}} x_{i_1 \dots i_h j_1 j_2} + \dots + (-1)^{n-h} x_{i_1 \dots i_n} = \\ &= \sum_{k=0}^{n-h} (-1)^k \sum_{\{j_1, \dots, j_k\} \subseteq \{i_{h+1}, \dots, i_n\}} x_{i_1 \dots i_h j_1 \dots j_k}, \end{aligned} \quad (41)$$

where by convention we set $x_{i_1 \dots i_h} = x_{\emptyset} = 1$ when $\{i_1, \dots, i_h\} = \emptyset$. Moreover, as each conditional constituent $\mathcal{C}_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}}$ is a nonnegative conditional random quantity, by coherence it must be

$$x_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}} = \sum_{k=0}^{n-h} (-1)^k \sum_{\{j_1, \dots, j_k\} \subseteq \{i_{h+1}, \dots, i_n\}} x_{i_1 \dots i_h j_1 \dots j_k} \geq 0, \quad \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}, \quad (42)$$

with

$$\sum_{\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} x_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}} = \sum_{\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} \sum_{k=0}^{n-h} (-1)^k \sum_{\{j_1, \dots, j_k\} \subseteq \{i_{h+1}, \dots, i_n\}} x_{i_1 \dots i_h j_1 \dots j_k} = 1, \quad (43)$$

as it also follows by observing that $\sum_{\{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}} \mathcal{C}_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}} = 1$.

Notice that, given a coherent prevision assessment $(x_{i_1 \dots i_h}; \emptyset \neq \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\})$ on the family $\{\mathcal{C}_{i_1 \dots i_h}; \emptyset \neq \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}\}$, where $\mathcal{C}_{i_1 \dots i_h} = (E_{i_1} | H_{i_1}) \wedge \dots \wedge (E_{i_h} | H_{i_h})$, as shown by formula (42) for every nonempty subset $\{i_1, \dots, i_h\}$ there exists a unique coherent extension $x_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}}$ for the prevision of the conditional constituent $\mathcal{C}_{i_1 \dots i_h \overline{i_{h+1}} \dots \overline{i_n}}$.

6. Necessary and sufficient conditions for coherence

In this section we obtain, under logical independence, two necessary and sufficient coherence conditions. Let a family of n conditional events $\mathcal{E} = \{E_1 | H_1, \dots, E_n | H_n\}$ be given, with $E_1, \dots, E_n, H_1, \dots, H_n$ logically independent. We denote by $\mathcal{M} = (x_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ a prevision assessment on $\mathcal{F} = \{\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, n\}\}$, where $\mathcal{C}_S = \bigwedge_{i \in S} (E_i | H_i)$ and $x_S = \mathbb{P}(\mathcal{C}_S)$. We observe that \mathcal{F} is the family of all $2^n - 1$ possible conjunctions among the conditional events in \mathcal{E} . The first condition characterizes the coherence of \mathcal{M} and will be represented in

geometrical terms by a suitable convex hull. The second condition characterizes the coherence of a prevision assessment on $\mathcal{F} \cup \mathcal{K}$, where \mathcal{K} is the set of conditional constituents associated with \mathcal{E} .

We denote by $C_0, C_1, \dots, C_{3^n-1}$, the constituents associated with the family \mathcal{E} , that is the elements of the partition of Ω obtained by expanding the expression

$$\bigwedge_{i=1}^n (E_i H_i \vee \bar{E}_i H_i \vee \bar{H}_i),$$

where $C_0 = \bar{H}_1 \cdots \bar{H}_n$. With each C_h we associate a point

$$Q_h = (q_{hS} : \emptyset \neq S \subseteq \{1, \dots, n\}), \quad (44)$$

where q_{hS} is the value of \mathcal{C}_S when C_h is true. In particular with C_0 it is associated $Q_0 = \mathcal{M}$. We notice that Q_h is the value of the random vector $(\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ when C_h is true. By discarding Q_0 , we denote by \mathcal{Q} the set of remaining points Q_h 's associated with the pair $(\mathcal{F}, \mathcal{M})$ and by $\mathcal{I}_{\mathcal{Q}}$ the convex hull of the set \mathcal{Q} . We denote by \mathcal{B} the subset of \mathcal{Q} , constituted by 2^n binary points Q_1, \dots, Q_{2^n} , defined as

$$\mathcal{B} = \{Q_1, \dots, Q_{2^n}\} = \{Q_h \in \mathcal{Q} : q_{hS} \in \{0, 1\}, S = \{i\}, i = 1, \dots, n\}. \quad (45)$$

We observe that the points Q_1, \dots, Q_{2^n} are associated with the 2^n constituents C_h 's obtained by expanding the expression

$$\bigwedge_{i=1}^n (E_i H_i \vee \bar{E}_i H_i),$$

which coincides with $\bigwedge_{i=1}^n H_i$. Notice that, given any C_h such that $Q_h \in \mathcal{B}$, the sub-vector $(q_{hS}, S = \{i\}, i = 1, \dots, n)$ is a vertex of the unit hypercube $[0, 1]^n$ and it is the value assumed by the random vector $(E_1|H_1, \dots, E_n|H_n)$ when C_h is true. We also remark that, from the definition of conjunction it follows that

$$Q_h \in \mathcal{B} \implies q_{hS} \in \{0, 1\}, \quad \forall \emptyset \neq S \subseteq \{1, \dots, n\}. \quad (46)$$

Then, the set \mathcal{B} can be equivalently defined as

$$\mathcal{B} = \{Q_h \in \mathcal{Q} : q_{hS} \in \{0, 1\}, \emptyset \neq S \subseteq \{1, \dots, n\}\}.$$

We denote by $\mathcal{I}_{\mathcal{B}}$ the convex hull of the set \mathcal{B} ; of course $\mathcal{I}_{\mathcal{B}} \subseteq \mathcal{I}_{\mathcal{Q}}$. Then we have

Theorem 15. Given a family of n conditional events $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$, let $\mathcal{M} = (x_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ be a prevision assessment on the family $\mathcal{F} = \{\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, n\}\}$, where $\mathcal{C}_S = \bigwedge_{i \in S} (E_i|H_i)$. Under the assumption of logical independence of $E_1, \dots, E_n, H_1, \dots, H_n$, the prevision assessment \mathcal{M} on \mathcal{F} is coherent if and only if \mathcal{M} belongs to the convex hull $\mathcal{I}_{\mathcal{B}}$ of the 2^n binary points Q_1, \dots, Q_{2^n} .

Proof. (\Rightarrow) Assume that \mathcal{M} is coherent. Then, all the inequalities in (42) are satisfied. We observe that the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$ is satisfied if there exist suitable nonnegative coefficients λ_h 's, with $\sum_{h=1}^{2^n} \lambda_h = 1$, such that $\mathcal{M} = \sum_{h=1}^{2^n} \lambda_h Q_h$. This means that for each component x_S of \mathcal{M} it must be $x_S = \sum_{h=1}^{2^n} \lambda_h q_{hS} = \sum_{h:q_{hS}=1} \lambda_h$. We observe that with each $Q_h \in \mathcal{B}$ it is associated a unique subset $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$ such that, when $S = \{i\}$, $i = 1, \dots, n$, it holds that $q_{hS} = q_{h\{i\}} = 1$ if $i \in \{i_1, \dots, i_k\}$ and $q_{hS} = q_{h\{i\}} = 0$ if $i \in \{i_{k+1}, \dots, i_n\} = \{1, \dots, n\} \setminus \{i_1, \dots, i_k\}$. Then, by changing notations, the point Q_h associated with $\{i_1, \dots, i_k\}$ will be denoted by the symbol $Q_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}$ and the coefficient λ_h will be denoted by $\lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}$. By this change of notations, the binary quantity q_{hS} becomes $q_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n S}$, with

$$q_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n S} = \begin{cases} 1, & \text{if } S \subseteq \{i_1, \dots, i_k\}, \\ 0, & \text{if } S \not\subseteq \{i_1, \dots, i_k\}. \end{cases}$$

Then the equality $x_S = \sum_{h=1}^{2^n} \lambda_h q_{hS}$ becomes $x_S = \sum_{\{i_1, \dots, i_k\} \supseteq S} \lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}$. Then, more explicitly, the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$ is satisfied if there exists a vector, with components, $\Lambda = (\lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}; \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\})$ which is a solution of the system below.

$$(\Sigma_{\mathcal{B}}) \begin{cases} x_S = \sum_{\{i_1, \dots, i_k\} \supseteq S} \lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}, & \emptyset \neq S \subseteq \{1, 2, \dots, n\}, \\ \sum_{\{i_1, \dots, i_k\} \subseteq \{1, 2, \dots, n\}} \lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n} = 1, \\ \lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n} \geq 0, & \{i_1, \dots, i_k\} \subseteq \{1, 2, \dots, n\}. \end{cases} \quad (47)$$

We observe that Λ has 2^n (nonnegative) components and $(\Sigma_{\mathcal{B}})$ has 2^n equations. By coherence of \mathcal{M} , from (42) and (43) we can compute the quantities $x_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}$, for all $\{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$, which are nonnegative and with their sum equal to 1. Moreover, by (29), for each subset S it holds that $x_S = \sum_{\{i_1, \dots, i_h\} \supseteq S} x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$, which has the same structure of the equation $x_S = \sum_{\{i_1, \dots, i_h\} \supseteq S} \lambda_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$ in $(\Sigma_{\mathcal{B}})$. Then, $(\Sigma_{\mathcal{B}})$ is solvable and the (unique) solution is the vector Λ with components

$$\lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n} = x_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n} = \mathbb{P}(\mathcal{C}_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}), \quad \forall \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}.$$

Thus, the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$ is satisfied.

(\Leftarrow) Assume that $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$. Then, $\mathcal{M} \in \mathcal{I}_{\mathcal{Q}}$ because $\mathcal{B} \subset \mathcal{Q}$ and hence the system (Σ) is solvable. Moreover, $\mathcal{I}_0 = \emptyset$ because all the coefficients $\lambda_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}$'s are associated with the constituents, which we denote by $C_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n}$'s, such that for every subset $\{i_1, \dots, i_k\}$ it holds that $C_{i_1 \dots i_k \bar{i}_{k+1} \dots \bar{i}_n} \subseteq H_i$, for every $i = 1, \dots, n$. Thus, by Theorem 2 the prevision assessment \mathcal{M} is coherent. \square

Remark 7. As shown by Theorem 15, under logical independence of the basic events $E_1, \dots, E_n, H_1, \dots, H_n$, the coherence of \mathcal{M} amounts to the solvability of system $(\Sigma_{\mathcal{B}})$. Moreover, $(\Sigma_{\mathcal{B}})$ is solvable if and only if the following inequalities are satisfied

$$\sum_{k=0}^{n-h} (-1)^k \sum_{\{j_1, \dots, j_k\} \subseteq \{i_{h+1}, \dots, i_n\}} x_{i_1 \dots i_h j_1 \dots j_k} \geq 0, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}, \quad (48)$$

where we recall that, by (42), the first member of (48) is the prevision $x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$ of $\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$. Therefore, under logical independence, the set of all coherent assessments on the family \mathcal{F} is the set of assessments \mathcal{M} , with components x_S which satisfy the list of linear inequalities (48). Indeed, $x_{i_1 \dots i_h j_1 \dots j_k}$ coincides with the component x_S , where $S = \{i_1, \dots, i_h, j_1, \dots, j_k\}$.

We will now give another result on coherence under logical independence. We denote by Δ the $(2^n - 1)$ -dimensional simplex of \mathbb{R}^{2^n} , that is the set of vectors $\mathcal{V} = (v_{\{i_1, \dots, i_h\}}, \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\})$ such that

$$\sum_{\{i_1, \dots, i_h\} \subseteq \{1, 2, \dots, n\}} v_{\{i_1, \dots, i_h\}} = 1, \quad v_{\{i_1, \dots, i_h\}} \geq 0, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}.$$

We observe that, given any $\mathcal{V} \in \Delta$, we can construct a prevision assessment $\mathcal{M} = (x_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ on $\mathcal{F} = \{\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, n\}\}$, where each x_S is obtained by applying (29) with $x_{i_1 i_2 \dots i_h \bar{i}_{h+1} \dots \bar{i}_k}$ replaced by $v_{\{i_1, \dots, i_h\}}$. Moreover, concerning the set \mathcal{K} of the conditional constituents $\{\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} : \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\}\}$, each $\mathcal{C}_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}$ is obtained from suitable elements of \mathcal{F} by applying (40). In this way, we also obtain a prevision assessment $\mathcal{P} = (x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n}, \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\})$ on \mathcal{K} , with $x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = v_{\{i_1, \dots, i_h\}}$; thus $\mathcal{P} = \mathcal{V}$. The next result shows that, under logical independence, by the simplex Δ we obtain all the coherent prevision assessments on $\mathcal{F} \cup \mathcal{K}$. For the sake of simplicity, even if both \mathcal{M} and \mathcal{P} contain the element $x_{1 \dots n} = \mathbb{P}(\mathcal{C}_{1 \dots n})$, we denote by $(\mathcal{M}, \mathcal{P}) = (\mathcal{M}, \mathcal{V})$ the prevision assessment on $\mathcal{F} \cup \mathcal{K}$ associated with \mathcal{V} .

Theorem 16. Let a family of n conditional events $\mathcal{E} = \{E_1|H_1, \dots, E_n|H_n\}$ be given, with $E_1, \dots, E_n, H_1, \dots, H_n$ logically independent. A prevision assessment $(\mathcal{M}, \mathcal{P})$ on $\mathcal{F} \cup \mathcal{K}$ is coherent if and only if it is associated to a vector $\mathcal{V} \in \Delta$.

Proof. If $(\mathcal{M}, \mathcal{P})$ is a coherent prevision assessment on $\mathcal{F} \cup \mathcal{K}$, then \mathcal{M} is obtained from \mathcal{P} by means of (29) and from (25) it holds that $\mathcal{P} \in \Delta$. Then $(\mathcal{M}, \mathcal{P})$ is associated with the vector $\mathcal{V} = \mathcal{P} \in \Delta$.

Conversely, if $(\mathcal{M}, \mathcal{P})$ is associated to some $\mathcal{V} \in \Delta$, then $\mathcal{P} = \mathcal{V}$. Moreover, as shown in the proof of Theorem 15, under logical independence, by setting

$$\lambda_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = x_{i_1 \dots i_h \bar{i}_{h+1} \dots \bar{i}_n} = v_{\{i_1, \dots, i_h\}}, \quad \forall \{i_1, \dots, i_h\} \subseteq \{1, \dots, n\},$$

the system $(\Sigma_{\mathcal{B}})$ is solvable, that is $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$; thus \mathcal{M} is coherent. Finally, if we extend the assessment \mathcal{M} , defined on \mathcal{F} , to the family \mathcal{K} , by recalling (29) and (42) the extension coincides with \mathcal{P} . Hence, $(\mathcal{M}, \mathcal{P})$ is coherent. \square

7. Some further aspects

In this section we examine some further aspects which are related with Theorem 15. We observe that each Q_h defined as in (44) is itself a prevision assessment on \mathcal{F} and hence, by Theorem 15, Q_h is coherent if and only if $Q_h \in \mathcal{I}_{\mathcal{B}}$. In the next result we prove that, under logical independence, coherence of \mathcal{M} requires coherence of all the points Q_h 's. In other words, if \mathcal{M} is coherent, then for each h it holds that $Q_h \in \mathcal{I}_{\mathcal{B}}$, even if $Q_h \notin \mathcal{B}$.

Theorem 17. Let n conditional events $E_1|H_1, \dots, E_n|H_n$ be given, with $E_1, H_1, \dots, E_n, H_n$ logically independent. Given a coherent prevision assessment $\mathcal{M} = (x_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ on the family $\mathcal{F} = \{\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, n\}\}$, for every point Q_h it holds that Q_h is a coherent assessment on \mathcal{F} , or equivalently $Q_h \in \mathcal{I}_{\mathcal{B}}$.

Proof. Of course, for each $Q_h \in \mathcal{B}$ it holds that $Q_h \in \mathcal{I}_{\mathcal{B}}$, that is Q_h is coherent. Let us consider any point Q_h , $h \neq 0$, associated with a constituent C_h , such that $Q_h \notin \mathcal{B}$. Without loss of generality we assume that

$$C_h \subset \bar{H}_1 \cdots \bar{H}_k H_{k+1} \cdots H_n, \quad 1 \leq k < n.$$

Then, for the components q_{hS} of Q_h , with $S = \{i\}$, $i = 1, \dots, n$, it holds that

$$q_{hS} = \begin{cases} x_i, & \text{if } S = \{i\}, i = 1, \dots, k; \\ b_i \in \{0, 1\}, & \text{if } S = \{i\}, i = k + 1, \dots, n. \end{cases}$$

More in general we have

$$q_{hS} = \begin{cases} x_S, & \text{if } S \subseteq \{1, \dots, k\}, \\ 1, & \text{if } S \subseteq \{k + 1, \dots, n\} \text{ and } b_i = 1, \forall i \in S, \\ 0, & \text{if } S \subseteq \{k + 1, \dots, n\} \text{ and } b_i = 0, \text{ for some } i \in S, \\ x_{S'} & \text{if } S \cap \{1, \dots, k\} = S' \neq \emptyset \text{ and } b_i = 1, \forall i \in S \setminus S' \neq \emptyset, \\ 0 & \text{if } S \cap \{1, \dots, k\} = S' \neq \emptyset \text{ and } b_i = 0, \text{ for some } i \in S \setminus S' \neq \emptyset. \end{cases} \quad (49)$$

We denote by \mathcal{M}_k the sub-assessment of \mathcal{M} defined as

$$\mathcal{M}_k = (x_S : \emptyset \neq S \subseteq \{1, \dots, k\})$$

on the sub-family \mathcal{F}_k of \mathcal{F} defined as

$$\mathcal{F}_k = (\mathcal{C}_S : \emptyset \neq S \subseteq \{1, \dots, k\}).$$

The coherence of \mathcal{M} implies the coherence of the sub-assessment \mathcal{M}_k . We observe that, as the events E_i, H_i , $i = 1, \dots, n$ are logically independent, the extension \mathcal{M}_k^* of \mathcal{M}_k on $\mathcal{F}_k \cup \{E_i | H_i, i = k + 1, \dots, n\}$, such that $P(E_i | H_i) = b_i \in \{0, 1\}$ for $i = k + 1, \dots, n$, is coherent. Moreover, there exists a unique extension \mathcal{M}^* of \mathcal{M}_k^* on the family \mathcal{F} because, for the assessment

$$\mathcal{M}^* = (x_S^* : \emptyset \neq S \subseteq \{1, \dots, n\}),$$

each component x_S^* is uniquely determined by \mathcal{M}_k^* . Indeed, it holds that

$$x_S^* = \begin{cases} x_S, & \text{if } S \subseteq \{1, \dots, k\}, \\ 1, & \text{if } S \subseteq \{k + 1, \dots, n\} \text{ and } b_i = 1, \forall i \in S, \\ 0, & \text{if } S \subseteq \{k + 1, \dots, n\} \text{ and } b_i = 0, \text{ for some } i \in S, \\ x_{S'} & \text{if } S \cap \{1, \dots, k\} = S' \neq \emptyset \text{ and } b_i = 1, \forall i \in S'' = S \setminus S' \neq \emptyset, \\ 0 & \text{if } S \cap \{1, \dots, k\} = S' \neq \emptyset \text{ and } b_i = 0, \text{ for some } i \in S'' = S \setminus S' \neq \emptyset. \end{cases} \quad (50)$$

The uniqueness of the extension $x_S^* = 0$, or $x_S^* = 1$ shown in the second and third lines of (50), follows from Theorem 5. Moreover, the uniqueness of the extension $x_S^* = x_{S'}$ follows because, by Theorem 6, it holds that

$$\max\{x_{S'} + x_{S''} - 1, 0\} \leq x_S^* \leq \min\{x_{S'}, x_{S''}\}, \quad (51)$$

and, from Theorem 5, it holds that $x_{S''} = \mathbb{P}(\mathcal{C}_{S \setminus S'}) = 1$; thus (51) becomes

$$\max\{x_{S'} + x_{S''} - 1, 0\} = x_{S'} \leq x_S^* \leq x_{S'} = \min\{x_{S'}, x_{S''}\}.$$

Finally, the uniqueness of the extension $x_S^* = 0$ in the last line of (50) follows because, from Theorem 5, it holds that $x_{S''} = \mathbb{P}(\mathcal{C}_{S \setminus S'}) = 0$; thus (51) becomes

$$\max\{x_{S'} + x_{S''} - 1, 0\} = 0 \leq x_S^* \leq 0 = \min\{x_{S'}, x_{S''}\}.$$

Of course, as the extension \mathcal{M}^* of \mathcal{M}_k^* is unique, coherence of \mathcal{M}_k^* implies coherence of \mathcal{M}^* . Then by Theorem 15, it holds that $\mathcal{M}^* \in \mathcal{I}_{\mathcal{B}}$. Finally, from (49) and (50) it follows that $x_S^* = q_{hS} \forall S \neq \emptyset$. Therefore $\mathcal{M}^* = \mathcal{Q}_h$, so that \mathcal{Q}_h is coherent, or equivalently $\mathcal{Q}_h \in \mathcal{I}_{\mathcal{B}}$. \square

Remark 8. We recall that each \mathcal{Q}_h associated with the pair $(\mathcal{F}, \mathcal{M})$ represents the value of the random vector $(C_S : \emptyset \neq S \subseteq \{1, \dots, n\})$ when C_h is true. Then coherence of \mathcal{M} implies that, as for the case of unconditional events, each possible value \mathcal{Q}_h of the random vector is itself a particular coherent assessment on \mathcal{F} .

8. Some examples and counterexamples

As shown by Theorem 15, under logical independence of $E_1, \dots, E_n, H_1, \dots, H_n$, coherence of \mathcal{M} amounts to condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$, that is to validity of all inequalities in formula (42). We examine this aspect for $n = 2$ and $n = 3$ in the examples below.

Example 3. In this example we obtain the lower and upper bounds given in Theorem 4 by using the conditional constituents. We consider $\mathcal{E} = \{E_1|H_1, E_2|H_2\}$ and $\mathcal{F} = \{E_1|H_1, E_2|H_2, (E_1|H_1) \wedge (E_2|H_2)\}$, with E_1, E_2, H_1, H_2 logically independent. Then, let $\mathcal{M} = (x_1, x_2, x_{12})$ be a prevision assessment on \mathcal{F} . The set of conditional constituents is $\mathcal{K} = \{\mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{\bar{1}2}, \mathcal{C}_{\bar{1}\bar{2}}\}$, where $\mathcal{C}_{12} = (E_1|H_1) \wedge (E_2|H_2)$, $\mathcal{C}_{1\bar{2}} = (E_1|H_1) \wedge (\bar{E}_2|H_2)$, $\mathcal{C}_{\bar{1}2} = (\bar{E}_1|H_1) \wedge (E_2|H_2)$, $\mathcal{C}_{\bar{1}\bar{2}} = (\bar{E}_1|H_1) \wedge (\bar{E}_2|H_2)$. As made in the proof of Theorem 15, we change notations for the points \mathcal{Q}_h 's of the set \mathcal{B} . In this example $n = 2$, then $\mathcal{B} = \{\mathcal{Q}_{12}, \mathcal{Q}_{1\bar{2}}, \mathcal{Q}_{\bar{1}2}, \mathcal{Q}_{\bar{1}\bar{2}}\}$, where

$$\mathcal{Q}_{12} = (1, 1, 1), \quad \mathcal{Q}_{1\bar{2}} = (1, 0, 0), \quad \mathcal{Q}_{\bar{1}2} = (0, 1, 0), \quad \mathcal{Q}_{\bar{1}\bar{2}} = (0, 0, 0).$$

The previsions of the conditional constituents $\mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{\bar{1}2}, \mathcal{C}_{\bar{1}\bar{2}}$ are, respectively,

$$x_{12}, \quad x_{1\bar{2}} = x_1 - x_{12}, \quad x_{\bar{1}2} = x_2 - x_{12}, \quad x_{\bar{1}\bar{2}} = 1 - x_1 - x_2 + x_{12}.$$

These previsions are the coefficients which allow to represent \mathcal{M} as a linear convex combinations of the points of the set \mathcal{B} . By Remark 7, coherence of \mathcal{M} amounts to the inequalities in (48), that is

$$x_{12} \geq 0, \quad x_1 - x_{12} \geq 0, \quad x_2 - x_{12} \geq 0, \quad 1 - x_1 - x_2 + x_{12} \geq 0,$$

which are equivalent to the following conditions

$$(x_1, x_2) \in [0, 1]^2, \quad \max\{0, x_1 + x_2 - 1\} \leq x_{12} \leq \min\{x_1, x_2\}.$$

Notice that, by recalling Theorem 16, each vector \mathcal{V} of the 3-dimensional simplex Δ determines a coherent prevision assessments on $\mathcal{F} \cup \mathcal{K} = \{E_1|H_1, E_2|H_2, \mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{\bar{1}2}, \mathcal{C}_{\bar{1}\bar{2}}\}$. For instance, with the vector $\mathcal{V} = (v_{\{1,2\}}, v_{\{1\}}, v_{\{2\}}, v_{\emptyset}) = (\frac{1}{6}, \frac{1}{6}, \frac{1}{3}, \frac{1}{3})$ it is associated the assessment $(\mathcal{M}, \mathcal{P}) = (x_1, x_2, x_{12}, x_{1\bar{2}}, x_{\bar{1}2}, x_{\bar{1}\bar{2}})$, where

$$\begin{aligned} x_1 &= v_{\{1,2\}} + v_{\{1\}} = \frac{1}{3}, & x_2 &= v_{\{1,2\}} + v_{\{2\}} = \frac{1}{2}, & x_{12} &= v_{\{1,2\}} = \frac{1}{6}, \\ x_{1\bar{2}} &= v_{\{1\}} = \frac{1}{6}, & x_{\bar{1}2} &= v_{\{2\}} = \frac{1}{3}, & x_{\bar{1}\bar{2}} &= v_{\{\emptyset\}} = \frac{1}{3}. \end{aligned}$$

Example 4. In this example, by using the set of conditional constituents, we obtain the same result given in [39, Corollary 1]. We start by a family $\mathcal{E} = \{E_1|H_1, E_2|H_2, E_3|H_3\}$, where $E_1, E_2, E_3, H_1, H_2, H_3$ are logically independent. The conditional constituents are $\mathcal{C}_{123} = (E_1|H_1) \wedge (E_2|H_2) \wedge (E_3|H_3), \dots, \mathcal{C}_{\bar{1}\bar{2}\bar{3}} = (\bar{E}_1|H_1) \wedge (\bar{E}_2|H_2) \wedge (\bar{E}_3|H_3)$. Let $\mathcal{M} = (x_1, x_2, x_3, x_{12}, x_{13}, x_{23}, x_{123})$ be a prevision assessment on the family $\mathcal{F} = \{E_1|H_1, E_2|H_2, E_3|H_3, \mathcal{C}_{12}, \mathcal{C}_{13}, \mathcal{C}_{23}, \mathcal{C}_{123}\}$, where $\mathcal{C}_{ij} = E_i|H_i \wedge E_j|H_j$. By logical independence and by Theorem 15, coherence of \mathcal{M} amounts to the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$, where \mathcal{B} is the set of points

$$\begin{aligned} \mathcal{Q}_{123} &= (1, 1, 1, 1, 1, 1), & \mathcal{Q}_{12\bar{3}} &= (1, 1, 0, 1, 0, 0, 0), & \mathcal{Q}_{\bar{1}23} &= (1, 0, 1, 0, 1, 0, 0), \\ \mathcal{Q}_{1\bar{2}3} &= (1, 0, 0, 0, 0, 0, 0), & \mathcal{Q}_{\bar{1}2\bar{3}} &= (0, 1, 1, 0, 0, 1), & \mathcal{Q}_{\bar{1}\bar{2}3} &= (0, 1, 0, 0, 0, 0, 0), \\ \mathcal{Q}_{1\bar{2}\bar{3}} &= (0, 0, 1, 0, 0, 0, 0), & \mathcal{Q}_{\bar{1}\bar{2}\bar{3}} &= (0, 0, 0, 0, 0, 0, 0). \end{aligned}$$

By recalling (42), the previsions of the conditional constituents, which are the coefficients in the representation of \mathcal{M} as a linear convex combinations of the points of the set \mathcal{B} , are

$$x_{123}, x_{12\bar{3}} = x_{12} - x_{123}, \quad x_{\bar{1}23} = x_{13} - x_{123}, \quad x_{1\bar{2}3} = x_{1\bar{2}} - x_{1\bar{2}3} = x_1 - x_{12} - x_{13} + x_{123},$$

$$x_{\bar{1}2\bar{3}} = x_{23} - x_{123}, \quad x_{1\bar{2}\bar{3}} = x_{\bar{1}2} - x_{1\bar{2}\bar{3}} = x_2 - x_{12} - x_{23} + x_{123}, \quad x_{\bar{1}\bar{2}3} = x_{\bar{1}3} - x_{\bar{1}23} = x_3 - x_{13} - x_{23} + x_{123},$$

$$x_{\bar{1}\bar{2}\bar{3}} = x_{\bar{1}\bar{2}} - x_{\bar{1}\bar{2}\bar{3}} = (1 - x_1 - x_2 + x_{12}) - (x_3 - x_{13} - x_{23} + x_{123}) = 1 - x_1 - x_2 - x_3 + x_{12} + x_{13} + x_{23} - x_{123}.$$

By Remark 7, coherence of \mathcal{M} amounts to the inequalities in (48), that is

$$x_{123} \geq 0, \quad x_{12} - x_{123} \geq 0, \quad x_{13} - x_{123} \geq 0, \quad x_1 - x_{12} - x_{13} + x_{123} \geq 0, \quad x_{23} - x_{123} \geq 0,$$

$$x_2 - x_{12} - x_{23} + x_{123} \geq 0, \quad x_3 - x_{13} - x_{23} + x_{123} \geq 0, \quad 1 - x_1 - x_2 - x_3 + x_{12} + x_{13} + x_{23} - x_{123} \geq 0,$$

which can be written as $(x_1, x_2, x_3) \in [0, 1]^3$, $x'_{123} \leq x_{123} \leq x''_{123}$, where

$$x'_{123} = \max\{0, x_{12} + x_{13} - x_1, x_{12} + x_{23} - x_2, x_{13} + x_{23} - x_3\},$$

$$x''_{123} = \min\{x_{12}, x_{13}, x_{23}, 1 - x_1 - x_2 - x_3 + x_{12} + x_{13} + x_{23}\}.$$

Notice that there are inequalities which hold, even if they are not evident. Indeed, as $x''_{123} \geq x'_{123}$, it holds for instance that $1 - x_1 - x_2 - x_3 + x_{12} + x_{13} + x_{23} \geq x_{12} + x_{13} - 1$, that is $x_{23} \geq x_2 + x_3 - 1$, and so on. Of course, if $x''_{123} < x'_{123}$ (because for instance $x_{12} < x_{12} + x_{13} - x_1$, that is $x_{13} > x_1$), then the assessment is not coherent. Moreover, by recalling Theorem 16, each vector \mathcal{V} of the 7-dimensional simplex Δ determines a coherent prevision assessments on $\mathcal{F} \cup \mathcal{K}$.

We also remark that, in case of some logical dependencies, Theorem 15 is no more valid; that is, coherence is not equivalent to the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$. We give below two examples; in the first one the set \mathcal{B} is empty.

Example 5. Let three events A, H, K be given, with $HK = \emptyset$ and A logically independent of H and K . Moreover, let $\mathcal{M} = (x, y, z)$ be a prevision assessment on the family $\mathcal{F} = \{A|H, A|K, (A|H) \wedge (A|K)\}$. The constituents generated by $\{A|H, A|K\}$ are

$$C_1 = AH\bar{K}, C_2 = A\bar{H}K, C_3 = \bar{A}H\bar{K}, C_4 = \bar{A}\bar{H}K, C_0 = \bar{H}\bar{K}.$$

The associated points Q_h 's for the pair $(\mathcal{F}, \mathcal{M})$ are

$$Q_1 = (1, y, y), Q_2 = (x, 1, x), Q_3 = (0, y, 0), Q_4 = (x, 0, 0), Q_0 = \mathcal{M} = (x, y, z).$$

As we can see, it holds that $\mathcal{B} = \emptyset$ and hence $\mathcal{I}_{\mathcal{B}} = \emptyset$; then, to check coherence of \mathcal{M} we cannot use the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$, which is meaningless. Instead, in order to check coherence we need to start by checking the condition $\mathcal{M} \in \mathcal{I}_Q$, which amounts to solvability of the system below.

$$\begin{cases} x = \lambda_1 + \lambda_2x + \lambda_4x, \\ y = \lambda_1y + \lambda_2 + \lambda_3y, \\ z = \lambda_1y + \lambda_2x, \\ \lambda_1 + \dots + \lambda_4 = 1, \quad \lambda_h \geq 0, \quad h = 1, 2, 3, 4, \end{cases}$$

which can be written as

$$\begin{cases} xy = \lambda_1y + \lambda_2xy + \lambda_4xy, \\ xy = \lambda_1xy + \lambda_2x + \lambda_3xy, \\ z = \lambda_1y + \lambda_2x, \\ \lambda_1 + \dots + \lambda_4 = 1, \quad \lambda_h \geq 0, \quad h = 1, 2, 3, 4. \end{cases}$$

By summing the first two equations, we obtain: $z = xy$; then, the unique coherent extension of (x, y) to the conditional constituent $(A|H) \wedge (A|K)$ is $z = xy$ (see also [38]). We observe that the assessment (x, y, z) uniquely determines the extensions to the other conditional constituents, $(A|H) \wedge (\bar{A}|K)$, $(\bar{A}|H) \wedge (A|K)$, and $(\bar{A}|H) \wedge (\bar{A}|K)$, given by $x(1-y)$, $(1-x)y$, and $(1-x)(1-y)$, respectively.

Remark 9. Example 5 shows that in general, given two conditional events $E_1|H_1, E_2|H_2$, in order a prevision assessment $(x_{12}, x_{1\bar{2}}, x_{\bar{1}2}, x_{\bar{1}\bar{2}})$ on the family of conditional constituents $\{\mathcal{C}_{12}, \mathcal{C}_{1\bar{2}}, \mathcal{C}_{\bar{1}2}, \mathcal{C}_{\bar{1}\bar{2}}\}$ be coherent, it is not sufficient that the conditions given in (25), that is

$$x_{12} + x_{1\bar{2}} + x_{\bar{1}2} + x_{\bar{1}\bar{2}} = 1, \quad x_{12} \geq 0, x_{1\bar{2}} \geq 0, x_{\bar{1}2} \geq 0, x_{\bar{1}\bar{2}} \geq 0,$$

be satisfied. Indeed, even if the previous conditions imply that $x_{12} + x_{1\bar{2}} = x_1$, and $x_{12} + x_{\bar{1}2} = x_2$, in Example 5 coherence also requires that the conditions $x_{12} = x_1x_2$, $x_{1\bar{2}} = x_1(1-x_2)$, $x_{\bar{1}2} = (1-x_1)x_2$, and $x_{\bar{1}\bar{2}} = (1-x_1)(1-x_2)$ be satisfied, and this is not guaranteed. Then, the assessment $(x_{12}, x_{1\bar{2}}, x_{\bar{1}2}, x_{\bar{1}\bar{2}})$ could be incoherent. The same remark holds more in general when we consider the c-constituents associated with n conditional events. In other words, the conditions in (25) are necessary but not sufficient for coherence.

Example 6. We examine the previous example, by assuming $HK \neq \emptyset$. In this case the constituents generated by $\{A|H, A|K\}$ are

$$C_1 = AH\bar{K}, C_2 = A\bar{H}K, C_3 = \bar{A}H\bar{K}, C_4 = \bar{A}\bar{H}K, C_5 = AHK, C_6 = \bar{A}HK, C_0 = \bar{H}\bar{K},$$

and the points Q_h 's for the pair $(\mathcal{F}, \mathcal{M})$ are

$$Q_1 = (1, y, y), Q_2 = (x, 1, x), Q_3 = (0, y, 0), Q_4 = (x, 0, 0), Q_5 = (1, 1, 1), Q_6 = (0, 0, 0),$$

and $Q_0 = \mathcal{M} = (x, y, z)$. In this case $\mathcal{B} = \{Q_5, Q_6\} = \{(1, 1, 1), (0, 0, 0)\}$, that is \mathcal{B} is non empty, but its cardinality is less than $2^2 = 4$ as there are logical dependencies ($E_1 = E_2 = A$). Then, to check coherence of \mathcal{M} we cannot use the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{B}}$, but we still need to start by checking the condition $\mathcal{M} \in \mathcal{I}_{\mathcal{Q}}$, which amounts to solvability of the system below.

$$\begin{cases} x = \lambda_1 + \lambda_2x + \lambda_4x + \lambda_5, \\ y = \lambda_1y + \lambda_2 + \lambda_3y + \lambda_5, \\ z = \lambda_1y + \lambda_2x + \lambda_5, \\ \lambda_1 + \dots + \lambda_5 = 1, \quad \lambda_h \geq 0, \quad h = 1, 2, 3, 4, 5, \end{cases}$$

where it is immediate to verify that $z \leq x$ and $z \leq y$. Moreover, the system can be written as

$$\begin{cases} xy = \lambda_1y + \lambda_2xy + \lambda_4xy + \lambda_5y, \\ xy = \lambda_1xy + \lambda_2x + \lambda_3xy + \lambda_5x, \\ z = \lambda_1y + \lambda_2x + \lambda_5, \\ \lambda_1 + \dots + \lambda_5 = 1, \quad \lambda_h \geq 0, \quad h = 1, 2, 3, 4, 5. \end{cases}$$

By summing the first two equations, we obtain:

$$xy = z - \lambda_5(1 - x)(1 - y) - \lambda_6xy;$$

that is

$$z = xy + \lambda_5(1 - x)(1 - y) + \lambda_6xy \geq xy.$$

Then, the set of coherent extensions z of (x, y) to the conditional constituent $(A|H) \wedge (A|K)$ is the set $\{z : xy \leq z \leq \min\{x, y\}\}$ (see also [38, Theorem 5]).

9. Some comparison with other approaches

Usually in literature the notion of conjunction has been defined as a suitable conditional event; for some of these notions the lower and upper probability bounds have been computed in [51]. However, by defining compound conditionals as tri-valued entities, some basic probabilistic properties are not satisfied. Within our approach the conjunction of conditional events is no longer a tri-valued entity, but it is a suitable conditional random quantity with a finite number of possible values in the unit interval. Anyway, this lack of closure does not seem a high price to pay because by our definition we preserve relevant probabilistic properties. On the other hand, there is often a lack of closure with respect to mathematical operations. This happens, for instance, by considering the ratio of integer numbers.

In the next subsection we make a comparison between quasi conjunction and conjunction.

9.1. A comparison between quasi conjunction and conjunction

We recall below the notion of quasi conjunction ([1], see also [8, 23, 55]), which coincides with Sobociński conjunction ([11]), defined as

$$\begin{aligned} Q(E_1|H_1, E_2|H_2) &= [(\bar{H}_1 \vee E_1H_1) \wedge (\bar{H}_2 \vee E_2H_2)]|(H_1 \vee H_2) = \\ &= (E_1H_1E_2H_2 + \bar{H}_1E_2H_2 + \bar{H}_2E_1H_1)|(H_1 \vee H_2). \end{aligned} \quad (52)$$

Concerning the lower and upper bounds on quasi conjunction, the assessment (x_1, x_2) on $\{E_1|H_1, E_2|H_2\}$, with E_1, H_1, E_2, H_2 logically independent, propagates to the interval $[z', z'']$ on the probability of $Q(E_1|H_1, E_2|H_2)$, where ([29, 35])

$$z' = \max\{x_1 + x_2 - 1, 0\}, \quad z'' = \begin{cases} \frac{x_1 + x_2 - 2x_1x_2}{1 - x_1x_2}, & (x_1, x_2) \neq (1, 1), \\ 1, & (x_1, x_2) = (1, 1). \end{cases}$$

Notice that $z'' \geq \min\{x_1, x_2\}$, that is the upper bound for the quasi conjunction is greater than or equal to the Fréchet-Hoeffding upper bound. For instance, when $x_1 = x_2 = \frac{1}{2}$ it follows that $z'' = \frac{2}{3} > \frac{1}{2} = \min\{\frac{1}{2}, \frac{1}{2}\}$. Thus our notion of conjunction preserves Fréchet-Hoeffding bounds, while quasi conjunction does not. Table 4 illustrates the numerical values of quasi conjunction and conjunction of two conditional events. As shown in Table 4, the value of the conjunction is less

| | C_h | $E_1 H_1$ | $E_2 H_2$ | $(E_1 H_1) \wedge (E_2 H_2)$ | $Q(E_1 H_1, E_2 H_2)$ |
|-------|----------------------------|-----------|-----------|------------------------------|-----------------------|
| C_1 | $E_1H_1E_2H_2$ | 1 | 1 | 1 | 1 |
| C_2 | $E_1H_1\bar{E}_2H_2$ | 1 | 0 | 0 | 0 |
| C_3 | $E_1H_1\bar{H}_2$ | 1 | x_2 | x_2 | 1 |
| C_4 | $\bar{E}_1H_1E_2H_2$ | 0 | 1 | 0 | 0 |
| C_5 | $\bar{E}_1H_1\bar{E}_2H_2$ | 0 | 0 | 0 | 0 |
| C_6 | $\bar{E}_1H_1\bar{H}_2$ | 0 | x_2 | 0 | 0 |
| C_7 | $\bar{H}_1E_2H_2$ | x_1 | 1 | x_1 | 1 |
| C_8 | $\bar{H}_1\bar{E}_2H_2$ | x_1 | 0 | 0 | 0 |
| C_0 | $\bar{H}_1\bar{H}_2$ | x_1 | x_2 | x_{12} | z |

Table 4: Numerical values of the conjunctions. The values x_1, x_2, x_{12}, z denote $P(E_1|H_1)$, $P(E_2|H_2)$, $\mathbb{P}[(E_1|H_1) \wedge (E_2|H_2)]$ and $P[Q(E_1|H_1, E_2|H_2)]$, respectively.

than or equal to the value of the quasi conjunction when $H_1 \vee H_2$ is true. Then, by Remark 1, it holds that

$$\mathbb{P}[(E_1|H_1) \wedge (E_2|H_2)] = x_{12} \leq z = P[Q(E_1|H_1, E_2|H_2)]$$

and hence $(E_1|H_1) \wedge (E_2|H_2) \leq Q(E_1|H_1, E_2|H_2)$ also when $\bar{H}_1\bar{H}_2$ is true. Thus, in all cases it holds that

$$(E_1|H_1) \wedge (E_2|H_2) \leq Q(E_1|H_1, E_2|H_2). \quad (53)$$

We observe that, in the particular cases where $E_1H_1\bar{H}_2 = \bar{H}_1E_2H_2 = \emptyset$, or $x_1 = x_2 = 1$, by Theorem 3 it holds that $z = x_{12}$ and $(E_1|H_1) \wedge (E_2|H_2) = Q(E_1|H_1, E_2|H_2)$. More precisely $x_1 = x_2 = 1$ implies $z = x_{12} = 1$. Moreover, $E_1H_1\bar{H}_2 = \bar{H}_1E_2H_2 = \emptyset$ implies that

$$(E_1|H_1) \wedge (E_2|H_2) = Q(E_1|H_1, E_2|H_2) = E_1H_1E_2H_2|(H_1 \vee H_2),$$

with $z = x_{12} = P(E_1H_1E_2H_2|(H_1 \vee H_2))$. In this case, $(E_1|H_1) \wedge (E_2|H_2)$ also coincides with the Kleene-Lukasiewicz-Heyting conjunction $E_1H_1E_2H_2|(E_1H_1E_2H_2 \vee \bar{E}_1H_1 \vee \bar{E}_2H_2)$ (see [51, Table 2]). We recall that the Kleene-Lukasiewicz-Heyting conjunction coincides with the logical product between tri-events given in [25] (see also [46]). In addition, we observe that

$$Q(E_1|H_1, E_2|H_2) - (E_1|H_1) \wedge (E_2|H_2) = [(1 - x_1)\bar{H}_1E_2H_2 + (1 - x_2)E_1H_1\bar{H}_2]|(H_1 \vee H_2) \geq 0,$$

and

$$z - x_{12} = (1 - x_1)P(\bar{H}_1E_2H_2|(H_1 \vee H_2)) + (1 - x_2)P(E_1H_1\bar{H}_2|(H_1 \vee H_2)) \geq 0.$$

Then, to assess $z = x_{12}$ amounts to

$$(1 - x_1)P(\bar{H}_1E_2H_2|(H_1 \vee H_2)) = (1 - x_2)P(E_1H_1\bar{H}_2|(H_1 \vee H_2)) = 0,$$

that is $(P(\bar{E}_1|H_1) = 1 - x_1 = 0$ or $P(\bar{H}_1E_2H_2|(H_1 \vee H_2)) = 0)$ and $(P(\bar{E}_2|H_2) = 1 - x_2 = 0$ or $P(E_1H_1\bar{H}_2|(H_1 \vee H_2)) = 0)$. In addition, it is true that in a conditional bet on quasi conjunction we receive a random amount greater than or equal to the random amount received in a conditional bet on conjunction, but in these bets we pay two different amounts z and x_{12} , with $z \geq x_{12}$. Moreover, $z = x_{12}$ only in extreme cases where some suitable conditional probabilities are zero.

We also recall the notion of logical inclusion relation among conditional events given in [40] (see also [48] for an extension to conditional gambles). Given two conditional events $E_1|H_1$ and $E_2|H_2$, we say that $E_1|H_1$ implies $E_2|H_2$, denoted by $E_1|H_1 \subseteq E_2|H_2$, iff E_1H_1 true implies E_2H_2 true and \bar{E}_2H_2 true implies \bar{E}_1H_1 true; i.e., iff $E_1H_1 \subseteq E_2H_2$ and $\bar{E}_2H_2 \subseteq \bar{E}_1H_1$. Then, we remark that given two conditional events $E_1|H_1, E_2|H_2$, with $E_1|H_1 \subseteq E_2|H_2$, for the quasi conjunction it holds that ([34])

$$E_1|H_1 \subseteq Q(E_1|H_1, E_2|H_2) \subseteq E_2|H_2,$$

while in our approach one has

$$(E_1|H_1) \wedge (E_2|H_2) = E_1|H_1.$$

Moreover, if $E_1|H_1 \subseteq E_2|H_2 \subseteq E_3|H_3$ then

$$(E_1|H_1) \wedge (E_2|H_2) \wedge (E_3|H_3) = E_1|H_1,$$

while

$$E_1|H_1 \subseteq Q(E_1|H_1, E_2|H_2) \subseteq Q(E_1|H_1, E_2|H_2, E_3|H_3) \subseteq E_3|H_3,$$

and so on (see also [35, Theorem 9]). We also observe that from $E_1|H_1 \subseteq E_2|H_2$, it follows that $P(E_1|H_1) \leq P(E_2|H_2)$ and $E_1|H_1 \leq E_2|H_2$. This property of conditional monotony of conditional probability, as shown in Remark 1, holds more in general for the conditional previsions of conditional random quantities. For instance, given $n + 1$ conditional events $E_1|H_1, \dots, E_{n+1}|H_{n+1}$, by applying Remark 1 with $X|H = \mathcal{C}_{1\dots n+1}$ and $Y|K = \mathcal{C}_{1\dots n}$, it holds that $\mathcal{C}_{1\dots n+1} \leq \mathcal{C}_{1\dots n}$ when $H_1 \vee \dots \vee H_{n+1}$ is true; then $\mathbb{P}(\mathcal{C}_{1\dots n+1}) \leq \mathbb{P}(\mathcal{C}_{1\dots n})$ and hence $\mathcal{C}_{1\dots n+1} \leq \mathcal{C}_{1\dots n}$ in all cases; a dual result is valid for disjunctions ([39, theorems 7 and 8]). As we can see, the property of conditional monotony of conditional previsions is satisfied.

9.2. On Boolean algebras of conditionals

Boolean algebras of conditionals have been studied in [26, 27], where the authors characterize the atomic structure of the algebra of conditionals and introduce the logic of Boolean conditionals. In their work the notions of conjunction \sqcap and disjunction \sqcup are not (completely) specified, but it is assumed that some basic properties are satisfied. For instance, given three events A, B, C , it is required that ([27, Proposition 1], see also [28, Proposition 3.3])

$$(A|B) \sqcap (B|C) = A|C, \text{ when } A \subseteq B \subseteq C. \quad (54)$$

In our approach we do not start by an algebra of events, by means of which an algebra of conditionals is constructed, but we consider arbitrary families of conditional events. Then we determine the associated constituents and directly define the notions of conjunction and disjunction, by verifying the properties. For instance, in our approach formula (54) holds. Indeed, by assuming that $A \subseteq B \subseteq C$, we obtain

$$(A|B) \wedge (B|C) = \begin{cases} 1, & \text{if } A \text{ is true,} \\ 0, & \text{if } \bar{A}C \text{ is true,} \\ x_{12}, & \text{if } \bar{C} \text{ is true,} \end{cases} \quad (55)$$

where $x_{12} = \mathbb{P}[(A|B) \wedge (B|C)]$. Moreover,

$$A|C = \begin{cases} 1, & \text{if } A \text{ is true,} \\ 0, & \text{if } \bar{A}C \text{ is true,} \\ z, & \text{if } \bar{C} \text{ is true,} \end{cases} \quad (56)$$

where $z = P(A|C)$. Then, by Theorem 3, it follows that $x_{12} = z$ and hence $(A|B) \wedge (B|C) = A|C$. Moreover, it holds that

$$\mathbb{P}[(A|B) \wedge (B|C)] = P(A|C) = P(AB|C) = P(A|BC)P(B|C) = P(A|B)P(B|C),$$

which is the well known compound probability theorem.

Our notion of conjunction satisfies another property which is related to the atoms of the Boolean algebra of conditionals studied in [27, 28]. This property is described in the result below (where it is not assumed that the conditioning events have positive probability).

Theorem 18. Let H_1, \dots, H_n be n pairwise incompatible events. Then,

$$(H_1|\Omega) \wedge (H_2|\bar{H}_1) \wedge \dots \wedge (H_n|\bar{H}_1 \dots \bar{H}_{n-1}) = P(H_2|\bar{H}_1) \dots P(H_n|\bar{H}_1 \dots \bar{H}_{n-1}) H_1, \quad (57)$$

so that

$$\mathbb{P}[(H_1|\Omega) \wedge (H_2|\bar{H}_1) \wedge \dots \wedge (H_n|\bar{H}_1 \dots \bar{H}_{n-1})] = P(H_1)P(H_2|\bar{H}_1) \dots P(H_n|\bar{H}_1 \dots \bar{H}_{n-1}).$$

Proof. We set $P(H_1) = x_1$ and $P(H_j|\bar{H}_1 \dots \bar{H}_{j-1}) = x_j$, $j = 2, \dots, n$, and $\mathbb{P}[(H_1|\Omega) \wedge (H_2|\bar{H}_1) \wedge \dots \wedge (H_n|\bar{H}_1 \dots \bar{H}_{n-1})] = x_{1\dots n}$. Formula (57) holds for $n = 2$ and $n = 3$. Indeed, for $n = 2$ it holds that

$$(H_1|\Omega) \wedge (H_2|\bar{H}_1) = \begin{cases} x_2, & \text{if } H_1 \text{ is true,} \\ 0, & \text{if } \bar{H}_1 \text{ is true,} \end{cases} = x_2 H_1, \quad (58)$$

so that

$$\mathbb{P}[(H_1|\Omega) \wedge (H_2|\bar{H}_1)] = x_{12} = x_2P(H_1) = x_1x_2 = P(H_1)P(H_2|\bar{H}_1).$$

Moreover, based on (58) and on Definition 2, for $n = 3$ we obtain

$$\begin{aligned} (H_1|\Omega) \wedge (H_2|\bar{H}_1) \wedge (H_3|\bar{H}_1\bar{H}_2) &= x_2H_1 \wedge (H_3|\bar{H}_1\bar{H}_2) = \\ &= x_2[(H_1H_3\bar{H}_2\bar{H}_1 + x_1\bar{\Omega}H_3\bar{H}_2\bar{H}_1 + x_3(H_1 \vee H_2)H_1)](\Omega \vee \bar{H}_1\bar{H}_2) = x_2(x_3H_1|\Omega) = x_2x_3H_1. \end{aligned}$$

We assume by induction that (57) holds for $n - 1$, that is

$$(H_1|\Omega) \wedge (H_2|\bar{H}_1) \cdots \wedge (H_{n-1}|\bar{H}_1 \cdots \bar{H}_{n-2}) = x_2 \cdots x_{n-1}H_1, \quad (59)$$

then we prove that it holds for n . Indeed, from (59) we obtain

$$(H_1|\Omega) \wedge (H_2|\bar{H}_1) \wedge \cdots \wedge (H_n|\bar{H}_1 \cdots \bar{H}_{n-1}) = x_2 \cdots x_{n-1}H_1 \wedge (H_n|\bar{H}_1 \cdots \bar{H}_{n-1}).$$

Moreover, by Definition 2, it holds that

$$\begin{aligned} H_1 \wedge (H_n|\bar{H}_1 \cdots \bar{H}_{n-1}) &= \\ &= (H_1H_n\bar{H}_1 \cdots \bar{H}_{n-1} + x_1\bar{\Omega}H_n\bar{H}_1 \cdots \bar{H}_{n-1} + x_n(H_1 \vee \cdots \vee H_{n-1})H_1)(\Omega \vee \bar{H}_1\bar{H}_2) = \\ &= x_nH_1|\Omega = x_nH_1. \end{aligned}$$

Finally,

$$(H_1|\Omega) \wedge (H_2|\bar{H}_1) \wedge \cdots \wedge (H_n|\bar{H}_1 \cdots \bar{H}_{n-1}) = x_2 \cdots x_{n-1}x_nH_1,$$

and hence $x_{1\dots n} = x_1 \cdots x_n$. □

9.3. Some theoretical aspects and applications of conjunction

In this section we recall some theoretical aspects and applications of our approach to compound conditionals.

- All the basic properties valid for the unconditional events are satisfied in our theory of compound conditionals. For instance, (generalized) De Morgan's Laws are satisfied; moreover the formula $P(E_1 \vee E_2) = P(E_1) + P(E_2) - P(E_1E_2)$ becomes $\mathbb{P}[(E_1|H_1) \vee (E_2|H_2)] = P(E_1|H_1) + P(E_2|H_2) - \mathbb{P}[(E_1|H_1) \wedge (E_2|H_2)]$.

-The Fréchet-Hoeffding lower and upper prevision bounds for the conjunction (and for the disjunction) of two conditional events still hold.

- A generalized inclusion-exclusion formula for the disjunction of conditional events holds in our approach to compound conditionals.

- We can introduce the notion of conditional constituents, with properties analogous to the case of unconditional events, which allow to characterize coherence when the basic events are logically independent.

- Conjoined conditionals have been applied to probabilistic nonmonotonic reasoning ([31, 39]), by obtaining a characterization for the property of probabilistic entailment of Adams ([1]). In particular, in [39] it has been shown that a conditional event $E_{n+1}|H_{n+1}$ is p-entailed from a p-consistent family of n conditional events $E_1|H_1, \dots, E_n|H_n$ if and only if the conjunction $\mathcal{C}_{1\dots n+1}$ of the premises and the conclusion coincides with the conjunction $\mathcal{C}_{1\dots n}$ of the premises. Another

equivalent condition is that $\mathcal{C}_{1\dots n} \leq E_{n+1}|H_{n+1}$. Moreover, by exploiting a suitable notion of iterated conditional, in [31] it has been shown that a family $\{E_1|H_1, E_2|H_2\}$ p-entails a conditional event $E_3|H_3$ if and only if the iterated conditional $(E_3|H_3)|((E_1|H_1) \wedge (E_2|H_2))$ is constant and coincides with 1.

- Compound conditionals have been also applied to the psychology of the probabilistic reasoning, where by exploiting the notion of iterated conditional, the probabilistic modus ponens has been generalized to conditional events ([53]).

- Another application to one-premise and two-premise centering inferences has been given in [30, 54], by also determining the lower and upper prevision bounds for the conclusion of the rules.

- We remark that, like in [1, 42] and differently from [45], the Import-Export Principle is not valid in our theory of compound and iterated conditionals. Then, as proved in [36] (see also [52, 54]), we avoid Lewis' triviality results ([44]). In addition, within our theory, we can explain some intuitive probabilistic assessments discussed in [19], by suitably formalizing different kinds of latent information ([52]).

10. Conclusions

In this paper we deepened the study of conjunctions and disjunctions among conditional events in the framework of conditional random quantities. We proved that the Fréchet-Hoeffding bounds are a necessary coherence condition for the prevision assessments on $\{\mathcal{C}_{1\dots k}, \mathcal{C}_{k+1\dots n}, \mathcal{C}_{1\dots n}\}$, for every $1 \leq k \leq n-1$. We obtained a decomposition formula for the conjunction and we introduced the set of (non negative) conditional constituents \mathcal{K} for a family \mathcal{E} of n conditional events.

We showed that, as in the case of unconditional events, the sum of the conditional constituents is equal to 1 and for each pair of them the conjunction is equal to 0. We verified that, for each non empty subset S , the conjunction \mathcal{C}_S is the sum of suitable conditional constituents in \mathcal{K} and hence the prevision of \mathcal{C}_S is the sum of the previsions of such conditional constituents.

We obtained a generalized inclusion-exclusion formula for the disjunction of n conditional events; we proved a suitable distributivity property and we examined some related probabilistic results.

Under logical independence, we characterized in terms of a suitable convex hull the set of all coherent prevision assessments on a family \mathcal{F} containing n conditional events and all the possible conjunctions among them. We showed that such a characterization amounts to the solvability of a linear system and we described the set of all coherent prevision assessments on \mathcal{F} by a list of linear inequalities. Based on the $(2^n - 1)$ -dimensional simplex Δ , we characterized (still under logical independence) the set of all coherent prevision assessments on $\mathcal{F} \cup \mathcal{K}$.

Then, given a coherent assessment \mathcal{M} on \mathcal{F} , we showed that every possible value Q_h of the random vector associated with \mathcal{F} is itself a particular coherent assessment on \mathcal{F} . We deepened some aspects of coherence by illustrating examples and counterexamples.

We made a comparison with other approaches, by obtaining a result related to the notion of atom of a Boolean algebra of conditionals introduced in [27, 28]. Finally, we discussed the significance and perspectives of our theory by illustrating basic theoretical aspects and some applications to nonmonotonic reasoning and to the psychology of probabilistic reasoning.

Future work should concern in particular the study of necessary and sufficient conditions of coherence in the general case of logical dependencies among the basic unconditional events, by exploiting the set of conditional constituents.

Further future work could concern the study of compound conditionals in the setting of imprecise probabilities and gambles. Indeed, indicators of conditional events are ternary gambles and our conjunction builds n -ary gambles from ternary ones. Another interesting aspect that could be deepened is the study of the role of our compound conditionals in the framework of fuzzy logic and information fusion (see, e.g., [13, 18, 21, 22]).

Declaration of competing interest

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

Acknowledgments

We thank the three anonymous reviewers for their careful reading of our manuscript. Their many insightful comments and suggestions were very helpful in improving this paper. Giuseppe Sanfilippo has been partially supported by the INdAM–GNAMPA Project (2020 Grant U-UFMBAZ-2020-000819).

References

- [1] Adams, E.W., 1975. The logic of conditionals. Reidel, Dordrecht.
- [2] Baratgin, J., Politzer, G., Over, D., Takahashi, T., 2018. The psychology of uncertainty and three-valued truth tables. *Frontiers in Psychology* 9, 1479. doi:10.3389/fpsyg.2018.01479.
- [3] Benferhat, S., Dubois, D., Prade, H., 1997. Nonmonotonic reasoning, conditional objects and possibility theory. *Artificial Intelligence* 92, 259–276. doi:10.1016/S0004-3702(97)00012-X.
- [4] Berti, P., Miranda, E., Rigo, P., 2017. Basic ideas underlying conglomerability and disintegrability. *International Journal of Approximate Reasoning* 88, 387 – 400. doi:10.1016/j.ijar.2017.06.009.
- [5] Biazzo, V., Gilio, A., Lukasiewicz, T., Sanfilippo, G., 2005. Probabilistic logic under coherence: Complexity and algorithms. *Annals of Mathematics and Artificial Intelligence* 45, 35–81. doi:10.1007/s10472-005-9005-y.
- [6] Biazzo, V., Gilio, A., Sanfilippo, G., 2008. Generalized coherence and connection property of imprecise conditional previsions, in: *Proc. IPMU 2008, Malaga, Spain, June 22 - 27*, pp. 907–914. URL: <http://www.gimac.uma.es/ipmu08/proceedings/html/120.html>.
- [7] Biazzo, V., Gilio, A., Sanfilippo, G., 2012. Coherent conditional previsions and proper scoring rules, in: *Advances in Computational Intelligence. IPMU 2012*. Springer Heidelberg. volume 300 of *CCIS*, pp. 146–156. doi:10.1007/978-3-642-31724-8_16.
- [8] Calabrese, P., 1987. An algebraic synthesis of the foundations of logic and probability. *Information Sciences* 42, 187 – 237. doi:10.1016/0020-0255(87)90023-5.
- [9] Calabrese, P., 2017. *Logic and Conditional Probability: A Synthesis*. College Publications.
- [10] Capotorti, A., Lad, F., Sanfilippo, G., 2007. Reassessing accuracy rates of median decisions. *American Statistician* 61, 132–138. doi:10.1198/000313007X190943.
- [11] Ciucci, D., Dubois, D., 2012. Relationships between Connectives in Three-Valued Logics, in: *Advances on Computational Intelligence*. Springer. volume 297 of *CCIS*, pp. 633–642. doi:10.1007/978-3-642-31709-5_64.

- [12] Ciucci, D., Dubois, D., 2013. A map of dependencies among three-valued logics. *Information Sciences* 250, 162 – 177. doi:10.1016/j.ins.2013.06.040.
- [13] Coletti, G., Petturiti, D., Vantaggi, B., 2017. Fuzzy memberships as likelihood functions in a possibilistic framework. *International Journal of Approximate Reasoning* 88, 547 – 566. doi:10.1016/j.ijar.2016.11.017.
- [14] Coletti, G., Scozzafava, R., 1999. Conditioning and inference in intelligent systems. *Soft Computing* 3, 118–130. doi:10.1007/s005000050060.
- [15] Coletti, G., Scozzafava, R., 2002. Probabilistic logic in a coherent setting. Kluwer, Dordrecht.
- [16] Coletti, G., Scozzafava, R., Vantaggi, B., 2013. Coherent conditional probability, fuzzy inclusion and default rules, in: Yager, R., Abbasov, A.M., Reformat, M.Z., Shahbazova, S.N. (Eds.), *Soft Computing: State of the Art Theory and Novel Applications*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 193–208. doi:10.1007/978-3-642-34922-5_14.
- [17] Coletti, G., Scozzafava, R., Vantaggi, B., 2015. Possibilistic and probabilistic logic under coherence: Default reasoning and System P. *Mathematica Slovaca* 65, 863–890. doi:10.1515/ms-2015-0060.
- [18] Coletti, G., Vantaggi, B., 2018. Coherent Conditional Plausibility: A Tool for Handling Fuzziness and Uncertainty Under Partial Information. Springer International Publishing, Cham. pp. 129–152. doi:10.1007/978-3-319-60207-3_9.
- [19] Douven, I., Dietz, R., 2011. A puzzle about Stalnaker’s hypothesis. *Topoi*, 31–37doi:10.1007/s11245-010-9082-3.
- [20] Douven, I., Elqayam, S., Singmann, H., van Wijnbergen-Huitink, J., 2019. Conditionals and inferential connections: toward a new semantics. *Thinking & Reasoning*, 1–41doi:10.1080/13546783.2019.1619623.
- [21] Dubois, D., Faux, F., Prade, H., 2020. Prejudice in uncertain information merging: Pushing the fusion paradigm of evidence theory further. *International Journal of Approximate Reasoning* 121, 1 – 22. doi:10.1016/j.ijar.2020.02.012.
- [22] Dubois, D., Liu, W., Ma, J., Prade, H., 2016. The basic principles of uncertain information fusion. An organised review of merging rules in different representation frameworks. *Information Fusion* 32, 12 – 39. doi:10.1016/j.inffus.2016.02.006.
- [23] Dubois, D., Prade, H., 1994. Conditional objects as nonmonotonic consequence relationships. *IEEE Trans. on Syst. Man and Cybernetics*, 24, 1724 –1740. doi:10.1109/21.328930.
- [24] Edgington, D., 1995. On conditionals. *Mind* 104, 235–329.
- [25] de Finetti, B., 1936. La logique de la probabilité, in: *Actes du Congrès International de Philosophie Scientifique*, Paris, 1935, pp. IV 1–IV 9.
- [26] Flaminio, T., Godo, L., Hosni, H., 2015. On the algebraic structure of conditional events, in: Destercke, S., Denoëux, T. (Eds.), *Symbolic and Quantitative Approaches to Reasoning with Uncertainty (ECSQARU 2015)*. Springer LNAI 9161, Dordrecht, pp. 106–116. doi:10.1007/978-3-319-20807-7_10.
- [27] Flaminio, T., Godo, L., Hosni, H., 2017. On boolean algebras of conditionals and their logical counterpart, in: *ECSQARU 2017*. Springer. volume 10369 of *LNCS*, pp. 246–256. doi:10.1007/978-3-319-61581-3_23.
- [28] Flaminio, T., Godo, L., Hosni, H., 2020. Boolean algebras of conditionals, probability and logic. *Artificial Intelligence* 286, 103347. doi:10.1016/j.artint.2020.103347.
- [29] Gilio, A., 2012. Generalizing inference rules in a coherence-based probabilistic default reasoning. *International Journal of Approximate Reasoning* 53, 413–434. doi:10.1016/j.ijar.2011.08.004.
- [30] Gilio, A., Over, D., Pfeifer, N., Sanfilippo, G., 2017. Centering and compound conditionals under coherence, in: *Soft Methods for Data Science*. Springer. volume 456 of *AISC*, pp. 253–260.
- [31] Gilio, A., Pfeifer, N., Sanfilippo, G., 2020. Probabilistic entailment and iterated conditionals, in: Elqayam, S., Douven, I., Evans, J.S.B.T., Cruz, N. (Eds.), *Logic and Uncertainty in the Human Mind: A Tribute to David E. Over*. Routledge, Oxon, pp. 71–101. doi:10.4324/9781315111902-6.
- [32] Gilio, A., Sanfilippo, G., 2013a. Conditional random quantities and iterated conditioning in the setting of coherence, in: van der Gaag, L.C. (Ed.), *ECSQARU 2013*. Springer, Berlin, Heidelberg. volume 7958 of *LNCS*, pp. 218–229. doi:10.1007/978-3-642-39091-3_19.
- [33] Gilio, A., Sanfilippo, G., 2013b. Conjunction, disjunction and iterated conditioning of conditional events, in: *Synergies of Soft Computing and Statistics for Intelligent Data Analysis*. Springer, Berlin. volume 190 of *AISC*,

- pp. 399–407. doi:10.1007/978-3-642-33042-1_43.
- [34] Gilio, A., Sanfilippo, G., 2013c. Probabilistic entailment in the setting of coherence: The role of quasi conjunction and inclusion relation. *International Journal of Approximate Reasoning* 54, 513–525. doi:10.1016/j.ijar.2012.11.001.
- [35] Gilio, A., Sanfilippo, G., 2013d. Quasi conjunction, quasi disjunction, t-norms and t-conorms: Probabilistic aspects. *Information Sciences* 245, 146–167. doi:10.1016/j.ins.2013.03.019.
- [36] Gilio, A., Sanfilippo, G., 2014. Conditional random quantities and compounds of conditionals. *Studia Logica* 102, 709–729. doi:10.1007/s11225-013-9511-6.
- [37] Gilio, A., Sanfilippo, G., 2017. Conjunction and disjunction among conditional events, in: Benferhat, S., Tabia, K., Ali, M. (Eds.), *IEA/AIE 2017, Part II*. Springer, Cham. volume 10351 of *LNCS*, pp. 85–96. doi:10.1007/978-3-319-60045-1_11.
- [38] Gilio, A., Sanfilippo, G., 2019a. Conjunction of conditional events and t-norms, in: Kern-Isberner, G., Ognjanović, Z. (Eds.), *Symbolic and Quantitative Approaches to Reasoning with Uncertainty, ECSQARU 2019*. Springer International Publishing. volume 11726 of *LNCS*, pp. 199–211. doi:10.1007/978-3-030-29765-7_17.
- [39] Gilio, A., Sanfilippo, G., 2019b. Generalized logical operations among conditional events. *Applied Intelligence* 49, 79–102. doi:10.1007/s10489-018-1229-8.
- [40] Goodman, I.R., Nguyen, H.T., 1988. Conditional Objects and the Modeling of Uncertainties, in: Gupta, M.M., Yamakawa, T. (Eds.), *Fuzzy Computing*. North-Holland, pp. 119–138.
- [41] Goodman, I.R., Nguyen, H.T., Walker, E.A., 1991. *Conditional Inference and Logic for Intelligent Systems: A Theory of Measure-Free Conditioning*. North-Holland. URL: www.dtic.mil/dtic/tr/fulltext/u2/a241568.pdf.
- [42] Kaufmann, S., 2009. Conditionals right and left: Probabilities for the whole family. *Journal of Philosophical Logic* 38, 1–53. doi:10.1007/s10992-008-9088-0.
- [43] Lad, F., 1996. *Operational subjective statistical methods: A mathematical, philosophical, and historical introduction*. Wiley, New York.
- [44] Lewis, D., 1976. Probabilities of conditionals and conditional probabilities. *The Philosophical Review* 85, 297–315.
- [45] McGee, V., 1989. Conditional probabilities and compounds of conditionals. *Philosophical Review* 98, 485–541. doi:<http://dx.doi.org/10.2307/2185116>.
- [46] Milne, P., 1997. Bruno de Finetti and the Logic of Conditional Events. *British Journal for the Philosophy of Science* 48, 195–232. URL: <http://www.jstor.org/stable/687745>.
- [47] Nguyen, H.T., Walker, E.A., 1994. A history and introduction to the algebra of conditional events and probability logic. *IEEE Transactions on Systems, Man, and Cybernetics* 24, 1671–1675. doi:10.1109/21.328924.
- [48] Pelessoni, R., Vicig, P., 2014. The Goodman-Nguyen relation within imprecise probability theory. *International Journal of Approximate Reasoning* 55, 1694–1707. doi:10.1016/j.ijar.2014.06.002.
- [49] Petturiti, D., Vantaggi, B., 2017. Envelopes of conditional probabilities extending a strategy and a prior probability. *International Journal of Approximate Reasoning* 81, 160 – 182. doi:10.1016/j.ijar.2016.11.014.
- [50] Pfeifer, N., Sanfilippo, G., 2017. Probabilistic squares and hexagons of opposition under coherence. *International Journal of Approximate Reasoning* 88, 282–294. doi:10.1016/j.ijar.2017.05.014.
- [51] Sanfilippo, G., 2018. Lower and upper probability bounds for some conjunctions of two conditional events, in: *SUM 2018*. Springer International Publishing, Cham. volume 11142 of *LNCS*, pp. 260–275. doi:10.1007/978-3-030-00461-3_18.
- [52] Sanfilippo, G., Gilio, A., Over, D., Pfeifer, N., 2020. Probabilities of conditionals and previsions of iterated conditionals. *International Journal of Approximate Reasoning* 121, 150 – 173. doi:10.1016/j.ijar.2020.03.001.
- [53] Sanfilippo, G., Pfeifer, N., Gilio, A., 2017. Generalized probabilistic modus ponens, in: Antonucci, A., Cholvy, L., Papini, O. (Eds.), *ECSQARU 2017*. Springer. volume 10369 of *LNCS*, pp. 480–490. doi:10.1007/978-3-319-61581-3_43.
- [54] Sanfilippo, G., Pfeifer, N., Over, D., Gilio, A., 2018. Probabilistic inferences from conjoined to iterated conditionals. *International Journal of Approximate Reasoning* 93, 103 – 118. doi:10.1016/j.ijar.2017.10.027.

- [55] Schay, G., 1968. An algebra of conditional events. *Journal of Mathematical Analysis and Applications* 24, 334–344. doi:10.1016/0022-247X(68)90035-8.