

Simultaneity in wavepacket reduction

Arthur Jabs

Alumnus, Technical University Berlin.
Voßstr. 9, 10117 Berlin, Germany
arthur.jabs@alumni.tu-berlin.de

(11 June 2015)

Abstract. It is conjectured that the effects of reduction, which occur with the individual wavepackets in a system of two separated though entangled packets, occur simultaneously in the center frame (properly defined) of the entangled packets. Existing experiments are compatible with this type of nonlocality. Further experimental tests are suggested.

Keywords: entanglement, reduction, simultaneity, preferred frame, spooky action at a distance

That the effects of reduction (collapse) on two spatially separated but entangled wavepackets occur at spacelike distances in the laboratory frame has been shown in many experiments with two photon packets. This is the quantum mechanical nonlocality. The experiments [1] - [4] moreover showed that any hypothetical influence of one reduction effect on the other in some other frame would have to proceed at a speed that exceeds the speed of light c by at least a factor of about 10^4 .

What is the exact value of that speed? At present there is no answer to that question. An interesting proposal is that the speed is infinite, in other words: that there is a reference frame where the two reduction effects occur simultaneously [5]. Indeed several candidates have been considered in the literature. In [7] the lab frame, the frame of the massive device that triggers the reduction, and the cosmic background radiation frame were considered, without definite conclusion however. As quantum nonlocality anyway implies some violation of special relativity theory a rather drastic proposal is to replace the Lorentz transformation by the Tangherlini transformation [8]. Then simultaneity becomes frame independent, and the question of a preferred reference frame becomes meaningless.

In this note another proposal is advanced, which is motivated by the concept that a system of entangled wavepackets (an entangled system, for short) is a fundamental region of space [6, Secs. 2.3, 3.1]: simultaneity occurs in the ‘center frame’ S_C of the entangled wavepackets and is restricted to the interior of the entangled system. The origin of this frame is defined as the mean position, with a wave function which describes the entangled system, for example:

$$\mathbf{r}_C := \langle \mathbf{r}(t) \rangle = \frac{1}{2} \int \mathbf{r} \left| \psi_1(\mathbf{r}, a, t) \psi_2(\mathbf{r}, a, t) \pm \psi_1(\mathbf{r}, b, t) \psi_2(\mathbf{r}, b, t) \right|^2 d^3r. \quad (1)$$

If in the center frame S_C the two reduction effects are connected by a hypothetical velocity \mathbf{u}_C forming an angle α_C with the relative velocity \mathbf{v} between S_C and the lab frame S_L , then within the lab frame the hypothetical velocity \mathbf{u}_L forming an angle α_L with \mathbf{v} is [9]:

$$u_L^2 = \frac{u_C^2 + v^2 + 2u_C v \cos \alpha_C - [(u_C v/c) \sin \alpha_C]^2}{[1 + (u_C v/c^2) \cos \alpha_C]^2} \quad (2)$$

and

$$\tan \alpha_L = \tan \alpha_C \frac{\sqrt{1 - (v/c)^2}}{1 + v/(u_C \cos \alpha_C)}. \quad (3)$$

Simultaneity in the center frame means $u_C = \infty$, and formulas (2) and (3) turn into

$$u_L^2 = \frac{c^4}{v^2} \frac{1 - [(v/c) \sin \alpha_C]^2}{\cos^2 \alpha_C} \quad (4)$$

$$\tan \alpha_L = \tan \alpha_C \sqrt{1 - (v/c)^2}, \quad (v \neq c). \quad (5)$$

The inverse formulas are obtained by replacing v by $-v$.

Some special cases are:

- (i) $\alpha_C = 0^\circ \quad \implies u_L = c^2/v, \alpha_L = 0^\circ,$ de Broglie waves ('waves of simultaneity' [10]).
- (ii) $v = c \quad \implies u_L = c,$
- (iii) $v = 0 \quad \implies u_L = \infty, \alpha_L = \alpha_C,$
- (iv) $v \neq c, \alpha_C = 90^\circ \implies u_L = \infty, \alpha_L = 90^\circ,$ ('transverse simultaneity').

The hypothetical velocity u_L in the lab frame lies between c and ∞ , depending on the values of v and α_C . The values of v and α_L are determined by the direction of each of the two wavepackets at the moment of the first reduction effect in the lab frame. In the actual experiments [1 - 4] $u_L > c$ is confirmed, and there is no indication of an upper limit to it. In [3] it was $v = 0$, so that (iii) yields indeed $u_L = \infty$. In [1, 2, 4] the angle α_L is 90° due to the symmetric arrangement of the receivers with respect to the source. Thus according to (iv) α_C is also 90° and $u_L = \infty$.

Though the experiments do not contradict the present proposal, the available data do not suffice to definitely confirm it. If it is possible to determine u_L , one way to achieve a confirmation would be to calculate $u_L(\alpha_L)$ by (4) and (5) as a function of v and α_C and to compare this kind of dependence with the measured values.

Notes and References

- [1] Zbinden, H., Brendel, J., Tittel, W. et al.: Experimental Test of Relativistic Quantum State Collapse with Moving Reference Frames, arXiv:quant-ph/0002031 (J. Phys. A: Math. Gen. **34**, 7103-7109 (2001))

- [2] Salart, D., Baas, A., Branciard, C. et al.: Testing spooky action at a distance, arXiv:0808.3316 (Nature **454**, 861-864 (2008))
 - [3] Cocciano, B., Faetti, S. and Fronzoni, L.: A lower bound for the velocity of quantum communications in the preferred frame, arXiv:1006.2697 (Phys. Lett. A **375**, 379-384 (2011))
 - [4] Yin, J., Cao, Y., Yong, H.-L. et al.: Bounding the speed of ‘spooky action at a distance’, arXiv:1303.0614 (Phys. Rev. Lett. **110**, 260407 (2013))
 - [5] We prefer to speak of two effects of one reduction rather than of two reductions. The effects themselves may take some small time intervals; this is considered negligible here. The reduction is conceived to occur independently of being observed or not [6, Sec. 4]
 - [6] Jabs, A.: Quantum mechanics in terms of realism, arXiv:quant-ph/9606017
 - [7] Gisin, N., Scarani, V., Tittel, W. et al.: Optical tests of quantum nonlocality: from EPR-Bell tests towards experiments with moving observers, Ann. Phys. (Leipzig) **9** (11-12), 831-841 (2000)
 - [8] Tangherlini, F.R.: Galilean-Like Transformation Allowed by General Covariance and Consistent with Special Relativity, <http://dx.doi.org/10.4236/jmp.2014.55033> (Journal of Modern Physics **5**, 230-243 (2014))
 - [9] Pauli, W.: Theory of Relativity (Dover Publications, New York, 1958) p. 16
 - [10] Rindler, W.: Introduction to Special Relativity, 2nd. ed. (Clarendon Press, Oxford, 1991) p. 83
-