

## Journal Pre-proofs

Feeding broccoli and cauliflower to dairy sheep: influence on feed intake, metabolic health status and milk production and composition

A. Martín, F.J. Giráldez, I. Mateos, C. Saro, J. Mateo, S. Andrés, I. Caro, M.J. Ranilla

PII: S1751-7311(25)00113-2  
DOI: <https://doi.org/10.1016/j.animal.2025.101530>  
Reference: ANIMAL 101530


To appear in: *Animal*

Received Date: 10 September 2024  
Revised Date: 16 April 2025  
Accepted Date: 17 April 2025

Please cite this article as: A. Martín, F.J. Giráldez, I. Mateos, C. Saro, J. Mateo, S. Andrés, I. Caro, M.J. Ranilla, Feeding broccoli and cauliflower to dairy sheep: influence on feed intake, metabolic health status and milk production and composition, *Animal* (2025), doi: <https://doi.org/10.1016/j.animal.2025.101530>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier B.V. on behalf of The animal Consortium.



## Feeding broccoli and cauliflower to dairy sheep: influence on feed intake, metabolic health status and milk production and composition

A. Martín <sup>a</sup>, F.J. Giráldez <sup>a</sup>, I. Mateos <sup>a,b</sup>, C. Saro <sup>a,b</sup>, J. Mateo <sup>c</sup>, S. Andrés <sup>a</sup>, I. Caro <sup>d</sup>, M.J. Ranilla <sup>a,b</sup>

<sup>a</sup> *Instituto de Ganadería de Montaña (CSIC- Universidad de León), ctra. León-Vega de infanzones, 24346, Vega de Infanzones, León, Spain*

<sup>b</sup> *Departamento de Producción Animal, Universidad de León, Campus Vegazana s/n, 24071 León, Spain*

<sup>c</sup> *Departamento de Higiene y Tecnología de los Alimentos, Universidad de León, Campus Vegazana s/n, 24071 León, Spain*

<sup>d</sup> *Departamento de Pediatría e Inmunología, Obstetricia y Ginecología, Nutrición y Bromatología, Psiquiatría e Historia de la Ciencia, Universidad de Valladolid, Avda. Ramón y Cajal, 7, 47005 Valladolid, Spain*

Corresponding author: María José Ranilla. Email: mjrang@unileon.es

### Highlights

- Supplementation of a dairy sheep diet with industrial waste.
- Broccoli and cauliflower have no effect on animal performance or metabolic status.
- Broccoli supplementation increased protein content in milk.
- Both diets increased saturated and decreased monounsaturated fatty acids in milk.
- Feeding broccoli or cauliflower did not affect the flavour of the cheese.

### Abstract

The production of broccoli and cauliflower generates large amounts of vegetable by-products that often accumulate in the environment; however, these residues have nutritional value for ruminants. This study evaluated the effects of including industrial broccoli and cauliflower waste in the diet of dairy ewes on feed intake, metabolic health and milk production and composition. Thirty Assaf dairy ewes in mid-lactation were divided into three groups, with 10 animals per group. The control group (**CON**) received

a total mixed ratio (TMR) *ad libitum*, the broccoli group (**BRO**) received the same TMR plus 1.5 kg FM/d of fresh chopped broccoli, and the cauliflower (**CAU**) group received the TMR plus 1.5 kg/d of fresh chopped cauliflower. The trial lasted 42 d, divided into 21 d for adaptation and 21 d of measurements, where feed intake and milk yield were recorded daily. Milk samples were taken once a week. At the end of the experiment, fresh cheese was prepared, using the milk from each feeding group. A sensory analysis was carried out, to detect eventual differences in flavour among treatments. Blood samples were taken to analyse the biochemical profile and acid-base status. The animals of the BRO and the CAU groups ate all the vegetal offered without decreasing the daily intake of TMR or affecting milk production. No differences were observed between groups in the biochemical profile or the blood acid-base status, with the exception of the urea concentration, which was higher in the BRO group compared to the control (55.2, 61.8, 57.2 mg/dL for CON, BRO and CAU;  $P=0.022$ ). Regarding milk composition, the most relevant differences were observed in protein content, being higher in the BRO group ( $P=0.006$ ) than in the CON and CAU groups, and in the fatty acids profile, which showed an increase in the saturated fatty acids content ( $P=0.012$ ) and a decrease in the monounsaturated fatty acids content ( $P=0.005$ ) in the groups supplemented with broccoli and cauliflower. In conclusion, supplementation with 1.5 kg of broccoli or cauliflower in lactating sheep did not affect feed intake, milk production or health status of the animals and had little impact on the quality of milk.

**Keywords:** by-product, ruminant feeding, ewes, agro-industrial wastes, food surplus

## Implications

The use of food not marketable for humans, such as waste and surpluses of broccoli and cauliflower, for animal feeding has economic and environmental relevance. For growers, it offers a way to profit from current losses. Environmentally, feeding ruminants with these by-products would help reduce agro-industrial waste, which could otherwise contribute to environmental pollution. The inclusion of 1.5 kg of broccoli or cauliflower in the diet did not adversely affect the health status or milk production of lactating ewes. These findings suggest that such agricultural by-products can be considered as sustainable feed resources for dairy sheep.

## Introduction

Worldwide broccoli and cauliflower production have considerably increased in last decades, reaching 27 million t in 2020 (FAO, 2020). This increase is likely due to their proven health-promoting effects as human food (Aghajanzpour et al., 2017; Jeffery and Araya, 2009; Owis, 2015), combined with the fact that broccoli and cauliflower are relatively easy to grow, prepare, and are usually competitively priced compared to other legumes. Spain is one of the main producing countries of broccoli and cauliflower, with a production of 746 510 t in 2020 (Conversa et al., 2019; and FAO, 2020). Only between 61 and 80 % of the above-ground biomass of broccoli and cauliflower, respectively, is comestible for humans (Rodríguez-Pallares and Rojas-González, 2022). Added to that, a significant proportion of comestible broccoli and cauliflower that do not meet commercial standards is discarded.

Brassica species appear to have a high nutritive value for ruminants, with high protein content, although low in fiber and relatively high in the calcium-phosphorous ratio

(Wiedenhoeft et al., 1994). Broccoli and cauliflower contain high amounts of secondary plant compounds, such as phenols, glucosinolates, sulforaphane; as well as vitamin C, E and A and a range of essential minerals (Nagraj et al., 2020; Picchi et al., 2020). However, despite the nutritive potential of these vegetables, not many studies have been carried out to evaluate their use for livestock feeding. To the best of our knowledge, no *in vivo* studies have been published yet regarding the use of cauliflower waste in dairy livestock, although *in vitro* studies suggest that it could be used without negative effects on ruminal fermentation (de Evan et al., 2020b).

In contrast, the inclusion of broccoli waste has been evaluated in dairy cows, goats and sheep, with contradictory effects on feed intake, milk yield and composition (Aziz, 2021; Monllor et al., 2020a, 2020b; Yi et al., 2015), which could be related to level of inclusion and the type of broccoli waste (dry, fresh or silage, or with leaves...). To our knowledge, no research has been conducted to study their effects on micelle properties or on sensory attributes of dairy products, as cheese.

This study was therefore aimed to evaluate the influence of supplementing lactating ewes with a limited amount of flowers and stems of broccoli or cauliflower on feed intake and efficiency, metabolic-health status and milk yield and composition.

## Material and methods

### *Experimental design*

The experimental protocols were approved by the Institutional Animal Care and Use Committee of the Universidad de León (approval number ULE\_014\_2016) and the Junta de Castilla y León (Spain), following proceedings described in Spanish and European Union legislation (Royal Decree 53/2013 and Council Directive 2010/63/EU).

Thirty Assaf dairy ewes in the middle of lactation ( $64 \pm 0.6$  d postpartum) were divided into three homogeneous groups (10 ewes per group) based on body weight ( $73.6 \pm 1.72$  kg) and milk yield ( $1980 \pm 102$  g/d). All animals received the same total mixed ration (TMR) *ad libitum*. The control group (**CON**) only received the TMR, the broccoli group (**BRO**) received daily the TMR diet plus 1.5 kg of freshly chopped commercial broccoli florets and the cauliflower group (**CAU**) received daily the TMR diet plus 1.5 kg of freshly chopped commercial cauliflower florets. TMR and BRO or CAU supplements were given in different feeders. The chemical composition of the TMR, the broccoli and the cauliflower is summarized in Table 1.

### *Experimental procedures*

Throughout the trial, the animals were housed in individual pens (2.25 m<sup>2</sup>) allowing visual and tactile contact with other animals. Fresh water was offered *ad libitum*. The trial was divided into two periods: an adaptation (21 d) and a measurement period (21 d). During the adaptation period, the animals received the TMR *ad libitum* and broccoli and cauliflower were progressively introduced. During the experimental period, BRO and CAU sheep were given 1.5 kg of broccoli and cauliflower, respectively, immediately after milking (at 9:00 in the morning). After 2 h, they were offered the TMR *ad libitum*.

Individual feed intake was recorded during the measurement period. Daily, before milking, refusals from each animal were removed and weighed and a sample was taken for chemical analysis. In addition, two 250 g samples of TMR, chopped broccoli and cauliflower were taken, one at the beginning of the trial and one at the end for chemical analysis. The amount of TMR offered was calculated and weighed daily to allow 15% leftovers. The individual intake was calculated as the difference in DM of the offered feeds and the DM of the remains.

Animals were weighed at the beginning of the adaptation period (day 1), to balance the groups, and then on days 21 and 42 (beginning and end of the experimental period). All animals remained in good health during the trial, except one from the cauliflower group which had to be removed from the experiment due to health issues (mastitis) unrelated to experimental diets.

Blood samples were collected to assess the biochemical profile and health status of the animals on day 0 (just before the start of the trial), day 21 (before the experimental period), and day 42 (at the end of the experiment). Two samples were taken per animal, one with heparin for acid-base analysis and the other without anticoagulant for biochemical analysis.

During the experiment, ewes were milked once a day and milk yield was recorded using automatic meters (DeLaval® MM 25 SG, Germany). On days 1, 3, 7, 10, 14, 17, and 21 of the measurement period milk samples were taken from each ewe to analyze chemical composition and somatic cell count. Additionally, a 12-mL milk sample was taken from each ewe on days 1, 7, 14 and 21, frozen, lyophilized and stored until fatty acid (FA) and liposoluble vitamin analysis. On the last day of the experiment, an additional milk sample (40-mL) was taken for casein micelle characterization, and 1 L of milk was also taken from each ewe and used for making fresh cheese for sensory analysis. These samples were immediately refrigerated at 4 °C.

The daily milk yield and the daily DM intake were used to calculate the FCR. The FCR was calculated as the ratio between the average milk production and the average DM intake (milk production (kg) / total DMI) at weekly intervals during the 21-day experimental period (Da Silva, 2021; 2023; 2024).

### **Chemical analysis**

The offered feed and the remains were analysed for DM (ISO 6496:1999), ash (ISO 5984:2002) and CP (ISO 5983:2009). Broccoli florets and the cauliflower samples were dried at 40°C for 24 h. Afterwards, samples were homogenized and frozen (-18°C) until fatty acid (FA) and liposoluble vitamin analysis, following the procedures described below for milk samples. Table 2 shows the FA percentage and liposoluble vitamins in the diet for the CON group (only TMR), the BRO group (TMR plus 1.5 kg of broccoli) and the CAU group (TMR plus 1.5 kg of cauliflower) calculated considering the average ingestion of TMR for each group (Da Silva, 2021)..

Blood samples with heparin were immediately analysed after sampling in a VetStat blood gas and electrolytes analyzer (Idexx, Barcelona, Spain) to assess the acid-base status by determining pH, bicarbonate ( $\text{HCO}_3^-$ ),  $\text{CO}_2$  pressure ( $\text{pCO}_2$ ), anion gap, total  $\text{CO}_2$  ( $\text{tCO}_2$ ),  $\text{Na}^+$ ,  $\text{K}^+$ , and  $\text{Cl}^-$  concentrations.

Blood samples without anticoagulant were allowed to clot in a water bath at 37°C for 30 min and then centrifuged at 3520 x *g* for 16 min at 4°C. The serum was stored at -80°C until it was used to measure biochemical parameters [alanine aminotransferase (ALT), glucose, calcium, urea, creatinine, lactic acid, albumin, gamma-glutamyl transferase (GGT),  $\beta$ -hydroxybutyric acid (BHB) and non-esterified fatty acids (NEFA)] using the clinical chemistry and turbidimetry analyzer BA400 (Biosystems, Barcelona, Spain).

Automated infrared (Miko-Scan 225, A/S Foss Electric, Denmark) was used to determine the content of crude fat, CP, total solids, urea and BHB in milk samples. The lactose content was calculated by subtracting the amount of fat and crude protein from the total solids. For the analysis of somatic cells, a Fossomatic 90 analyzer (Foss Electric, Denmark) was used.

For FA analysis, duplicate samples of 0.15 g lyophilised ewe milk were submitted to *in situ* transesterification (Carrapiso et al., 2000). Gas chromatography was performed using the method of Liu et al. (2007) with minor modifications. Helium was used at 3 mL/min, injector, detector and transfer line temperatures were 200, 300, and 230 °C, respectively. Two  $\mu$ L of extract were injected with a 20:1 split ratio. The oven temperature started at 170 °C (held for 24 min), then increased to 220 °C at 7.5 °C/min, and to 230 °C at 10 °C/min (held for 5 min). Detection, identification and quantification were carried out according to Andrés et al. (2019). Results were expressed as percentage of total FA.

Retinol and tocopherols were determined, in duplicate, in the milk samples. For the analysis of retinol, an aliquot (0.25 g) of lyophilized ewe milk was subjected to a saponification reaction and then solvent extraction with hexane (Mestre Prates et al., 2006). The filtered (0.45  $\mu$ m) hexane solution was analysed by high-performance liquid chromatography following Jin et al. (2015). Tocopherols were extracted, from 0.2 g of lyophilized milk, and analysed as describe by Humada et al. (2014), with the exception that no internal standard was added. The external standard method, with retinol,  $\alpha$ -tocopherol,  $\gamma$ -tocopherol and  $\delta$ -tocopherol standards (Sigma Aldrich Química, S.L., Madrid, Spain) was used for identification and quantification.

Samples for micelle characterization were centrifuged for 20 min at 3000 x *g* to separate the fat content from the milk. An aliquot of 4 mL of centrifuged-skimmed ewe milk (in the bottom layer of the tubes) was diluted with 4 mL of water then filtered through a 0.45  $\mu$ m-pore-size membrane with the help of a syringe and afterwards refrigerated in plugged glass tubes for up to 24 h (Glantz et al., 2010).

Micelle characteristics (particle size, distribution, and surface charge) were measured by dynamic light scattering (DLS) using a Zetasizer Advance - Pro (Red) instrument (Malvern Panalytical Ltd, Malvern, UK) equipped with a 663 nm and 10 mW He-Ne laser operating, at an output power of 148 V and a scattering angle of 173°. The measurements were carried out in triplicate at 25 °C with a refractive index of 1.345 and an equilibrating time of 2 min per sample. The D (0.5) value (median), Z-average, and polydispersity index (PDI) data were collected.

Triangle tests were performed by an untrained 30 member-panel. Analyses were carried out with fresh cheese prepared from the ewe's milk from each experimental group. After milking a mixture of one L of milk per ewe (10 L in total) was sieved and pasteurised at 65°C for 30 min, the milk was then tempered to 36 °C and curdled with commercial microbial rennet in 40 min. The coagulum was cut crossways into c.a. 1 cm<sup>3</sup> cubes, stirred for 30 min, the whey was then drawn off and the curd was manually salted with 1.5 % common salt. The salted curd was placed into 1-kg plastic moulds, was left to drain for 30 min and then stored under refrigeration until the next day. Two triangular tests were carried out with the cheeses in the same session. One to compare CON vs BRO cheeses and the other to compare CON vs CAU cheeses. In each test, three 20-g samples of ground cheese were given to the panellists; two samples came from the milk of one feeding treatment, and the other sample was from the other treatment. Each panellist was asked to identify the sample that did not match the other two. Physical facilities and sample preparation followed the recommendations described by Poste et al. (1991).

### ***Statistical analysis***

LBW changes and micelle properties were analysed using the GLM procedure of SAS (SAS Inst. Inc., Cary, NC) with the three experimental diets (CON, BRO and CAU) as the only source of variation. In all cases, the individual ewe was considered the experimental unit. All other data (feed intake, FCR, LBW, blood analysis, milk production and composition and casein properties) were analysed by repeated measures using the MIXED procedure of SAS (SAS Inst. Inc., Cary, NC). The model included fixed effects of diet, time and their interaction. The effect of animal nested in the diet was used as the error term to test the effect of diet. Significance was declared at  $P < 0.05$  and means were separated using the least significant difference procedure. Results of sensory tests were analysed by statistical significance tables for triangle tests (ISO, 2021; norm 4120).

## **Results**

### ***Feed intake, milk yield and feed efficiency***

Table 3 shows the mean values of feed intake and protein intake and body weight gain during the measurement period (21 days). No significant differences were observed in the dry matter intake (DMI) of TMR between the different experimental groups (3.14, 3.21 and 3.14 kg DM/ewe/d for CON, BRO and CAU;  $P = 1.00$ ). The DMI of broccoli and cauliflower was 0.125 and 0.124 kg/ewe/d, respectively. Thus, the total DMI by the broccoli and cauliflower group was 3.33 and 3.28 kg DM/ewe/d, respectively, being significantly ( $P < 0.05$ ) higher in BRO than in CON group. Protein intake was also higher in BRO than in the other groups (643, 690, 660 g/ewe/d in CON, BRO and CAU;  $P = 0.006$ ).

No differences were observed either in milk yield, weight gain or live weight between experimental groups. Feed conversion ratio (kg DMI/kg milk yield) was unaffected by dietary treatment ( $P > 0.05$ ).

### ***Biochemical blood analysis***

Table 4 shows the biochemical profile and acid-base parameters of the three experimental groups and the values at the three sampling times (days 1, 21 and 42). Regarding biochemical profile, only urea showed significant differences ( $P < 0.05$ ), with higher values in BRO group than in CON group. CAU animals showed intermediate values. Glucose, urea, BHB and NEFA varied between days of sampling ( $P < 0.05$ ), increasing glucose and BHB concentration, and decreasing urea and NEFA concentration through the trial. There was no diet x day interaction in any biochemical parameter ( $P > 0.05$ ).

Dietary treatment did not affect acid-base parameters, although the diet x day interaction was significant ( $P < 0.05$ ) for anion gap and Na concentration. Anion gap decreased with time in CON group ( $P = 0.001$ ), and increased in BRO group, with no changes on CAU group. Na concentration decreased with time in CON group and increased in both BRO and CAU group ( $P = 0.016$ ).

### ***Milk quality trails***

The mean values of milk quality parameters are shown in Table 5. There were no differences ( $P > 0.05$ ) in lactose, total solids, fat, urea and BHB concentrations. However, the protein content was affected ( $P = 0.003$ ) by the diet, being higher in the milk from animals supplemented with broccoli compared to CON group (49.7, 52.3, 49.8 g/ kg for CON, BRO and CAU).

Table 6 shows the FA profile and the liposoluble vitamin content of the ewes' milk as a function of the dietary treatment. The mean values for each group include the results from the milk sampled during all the experimental period (on days 1, 7, 14 and 21). The intake of brassicas by the ewes showed a modest but significant effect ( $P < 0.05$ ) on several FA percentages. Dietary brassica supplementation increased the saturated fatty acids (SFA) content by more than two percent units while decreased that of monounsaturated fatty acids (MUFA). Neither polyunsaturated fatty acids (PUFA) nor total  $n-6$  or  $n-3$  percentages were affected, although the  $n-6/n-3$  ratio decreased in the supplemented groups. Regarding individual FA, significant differences ( $P < 0.05$ ) were observed between CON and both BRO and CAU ewe milk. The supplementation with either of the two brassicas decreased the percentage of  $c9-18:1$  and increased that of  $c9,t11-18:2$ . The supplementation with BRO further increased 14:0 and 16:0 FA percentages and decreased those of iso-17:0 and 18:0 FAs compared to CON group.

Regarding to the casein properties, the experimental diets appeared to affect micelle size and size distribution properties in ewe milk (Table 7). There was a statistical trend ( $P = 0.066$ ) towards higher Z-average size in BRO milk. Moreover, the polydispersity index (indicating the variation in casein micelle sizes) was higher in CAU and in BRO than in CON ( $P < 0.05$  and  $P < 0.10$  respectively). Fig. 1 shows the mean particle size distribution of the micelles in the ewe milk from each dietary treatment. It can be observed a tendency toward a narrower bell-shaped curve for the CON milk and a tendency to larger particle size, i.e. a curve placed more to the right in the abscissa axis, for the BRO group milk.

The triangle test showed that fresh cheese from the BRO or CAU treatments could not be discriminated by the panelists ( $P > 0.05$ ; only a total of 7 or 10 of the 30 panellists were right in their response when compared the BRO or CAU with CON, respectively).



## Discussion

### ***Feed intake, milk yield and feed efficiency***

The mean values of DMI (Table 3) agree with those observed by other authors in sheep of the same breed, in a similar physiological state and consuming diets of similar characteristics (Prieto et al., 2013; Pulido et al., 2012).

The animals showed a high degree of acceptance for the broccoli and cauliflower, as they ate the whole amount offered. Due to the high moisture content of broccoli and cauliflower (92%), their daily DMI was around 125 g/d, accounting for less than 4% of the total DMI. This is probably the reason why supplementation with broccoli or cauliflower did not decrease TMR consumption. Due to the high moisture content of these materials, the water consumption associated with the feed was much higher in the BRO and CAU group compared to the CON ones (283, 1662, 1664 g/ewe/d for CON, BRO and CAU). Some authors, such as Felton and DeVries (2010), observed a linear reduction in DMI as DM content of the ration was reduced. This effect is associated with distension of the rumen and the response of ruminal mechanoreceptors, which send signals to the brain's satiety centres (Allen, 1996). However, in the present study, increased moisture in the ration did not result in lower DMI. This may be due to the animals of the broccoli and cauliflower group compensate for the greater intake of water from the ration with a lower intake of drinking water (INRA, 2019). Monllor et al. (2020b) observed a decrease in feed intake when goats were supplemented with broccoli silage, however it is important to highlight that the proportion of broccoli in the ration was much higher than in our study (40-60 vs 4%).

The milk yield agreed with those reported by Pulido et al. (2012) and Prieto et al. (2013) for Assaf ewes milked once a day. Milk yield was unaffected by dietary treatments, which is in concordance with the small magnitude of the differences observed on feed intake. In fact, there were no differences between experiment groups in live body weight change during the experimental period. These results are in agreement with those of Yi et al. (2015) in dairy cows and Monllor et al. (2020a) in goats, who, despite the higher inclusion of broccoli (40%), also did not observe differences in milk production.

### ***Biochemical blood profile***

It should be noted that all animals (ewes in mid-lactation) showed a positive body weight change, which suggest that they were in positive energy balance. In fact, blood glucose values increased and NEFA decreased throughout the experimental period, suggesting a reduction in glucose oxidation and lipolysis (Lisuzzo et al., 2022), associated with a reduction in the daily milk production and a positive energy balance. It should also be noted that the ewes did not become pregnant after the peak of lactation, so the energy requirements for milk production decreased throughout the experiment.

During negative energy balance, fat mobilization increases the flux of NEFA to the liver, which is partially transformed into ketone bodies, such as BHB (Lisuzzo et al., 2022). On this physiological basis, a reduction in blood BHB would be expected as NEFAs decreased and glucose increased. However, in the present study an opposite evolution

was observed, although it was probably not biologically relevant since the values were below the minimum ( $>0.8$  mmol/L) considered to suggest animals in subclinical ketosis (Araujo et al., 2020).

Although blood urea was higher in the BRO group than in the CON group, values are within the normal range for sheep (21-75 mg/dL; Jackson & Cockcroft, 2002). This parameter is closely related to the ingestion of degradable protein and in this case protein intake (643 and 690 g/ewe/d for CON and BRO) was higher in the BRO group, and *in vitro* studies (de Evan et al., 2020b, 2020a) have determined high rumen degradability of protein in cauliflower (88.5 %) and in broccoli (85.3%), which may lead to increase both production of ammonia in the rumen and blood urea levels. In contrast, Monllor et al. (2020b) observed a lower concentration of urea in the blood in goats fed diets including 25 to 40% of broccoli silage (on DM basis), although in this study feed intake was depressed in supplemented animals, reducing also protein intake. Although in our experiment the broccoli supplement was given fresh and not ensiled, it should be noted that it represented only the 10% of the total DM consumed. Therefore, a negative effect when using higher levels of broccoli cannot be ruled out.

There was also a trend in ALT to decrease in the BRO group (Table 4). ALT is used to assess liver status, with high values consistent with liver damage, while low values are associated with vitamin B6 deficiency (Ramati et al., 2015) and other factors. In broccoli 70-90% of glucosinolates are glucoraphanin, known for its antioxidant properties and association with liver protection. Human studies, such as the one conducted by Satomi et al. (2022a), have demonstrated a reduction in ALT values after a 24-week ingestion of 137.1  $\mu\text{g}$  of glucoraphanin. Notably, broccoli flowers contain approximately 246  $\mu\text{g/g}$  of DM glucoraphanin and could be the reason for the reduction of ALT in BRO group. Nevertheless, all values are within the normal range (Lisuzzo et al., 2022) and the observed differences could have not biological significance.

Finally, a diet x day interaction ( $P=0.005$ ) was detected for the anion gap seeing that values increase over time in BRO and CAU group animals but not in CON. At the end of the experiment (day 42), the values of the three groups remain within the normal range (12-24 mmol/L, Jackson & Cockcroft, 2002). The anionic gap is calculated as the difference between the cations ( $\text{Na}^+$ ,  $\text{K}^+$ ) and the anions ( $\text{Cl}^-$  and  $\text{HCO}_3^-$ ) and the increase in the anion gap may be related to an increase in  $\text{Na}^+$  values. In our case,  $\text{Na}^+$  values increased slightly over time in the groups supplemented with broccoli and cauliflowers. This is in line with the study of Monllor et al. (2020b) in which they found an increase in  $\text{Na}^+$  concentrations in the milk of goats fed 40% broccoli. This increase in  $\text{Na}^+$  milk values was possibly caused by an increase in  $\text{Na}^+$  in the plasma.

### ***Milk production and quality***

The mean values of milk chemical composition (total solids, fat, protein and lactose), as well as of milk production (Table 5), were into the ranges reported for the same breed of sheep in similar conditions (mid-lactation and milked once daily; (Prieto et al., 2013; Pulido et al., 2012)). In our trial, protein intake was higher in BRO ewes compared to CON group and this could explain the higher milk protein content in the animals supplemented with broccoli (Jaime and Purroy, 1995; Pulina et al., 2006).

Regarding the fatty acid profile in milk, the most important changes were an increase in SFA and a decrease in MUFA in the BRO and CAU groups compared to CON (Table 6). Few previous studies were found on the effect of brassica supplementation in dairy ruminants on the FA profile of milk (Monllor et al., 2020a, 2020c; Seguel et al., 2020). Seguel et al. (2020) fed lactating cows with a diet containing turnip or rape at an approximate level of 20% of the DMI reported an increase of SFA and PUFA percentages and a decrease of MUFA levels in supplemented animals respect to control ones. Similar effects were observed by Monllor et al. (2020a) who reported an increase in SFA percentages and a decrease in MUFA and PUFA percentages in the broccoli group versus the control group. This is in partial agreement with our results, since we observed a higher percentage of SFA and lower percentages of MUFA (a few percentage units each) and no effect on PUFA.

Regarding the effect on those milk fatty acids relevant for human health, we observed that brassica feeding decreased values in the n-6/n-3 ratio, being BRO more effective than CAU. This difference is linked to the observed tendency to lower n-3 values in the milk from CON group, although there were no significant differences in the separated values of n-6 and n-3 percentages among treatments. The lower n-6/n-3 ratio in BRO milk is also in concordance with Monllor et al. (2020a) who observed a decrease in both n-3 and n-6 concentration in the milk of the animals that received broccoli.

Looking at casein properties, the experimental diets seemed to influence micelle size and size distribution properties in ewe milk, as illustrated in Table 7 and Fig. 1. It has been reported that micelle diameters may be directly correlated with the content of total protein and negatively with the content of some caseins, such as k-casein (Freitas et al., 2019; Li et al., 2022). Therefore, the greater z-average diameter observed in BRO group could be related to the greater protein content. Unfortunately, casein profile was not evaluated so cannot confirm any effect of dietary treatments on casein profile.

Specific feedstuffs can contain low threshold odour active compounds or its direct metabolic precursors that can be transferred to the blood and then to milk. This is the case of S-methyl-cysteine sulfoxide or glucosinolates, which are present at relatively high amounts in brassica plants (Barry, 2013; Wiedenhoeft and Barton, 1995). In the event that these compounds passed into milk, their flavour could be detected. In this study, despite being feed on brassicas, it appeared that the potential presence odour active compounds did not affect milk flavour. Other studies carried out in lactating cows also observed no effects on flavour characteristics of milk or cheeses when fed different brassica species, such as forage rape (Seguel et al., 2020), typon (a hybrid of a Chinese cabbage and turnip) forage (Wiedenhoeft and Barton, 1995) or forage turnip (Seguel et al., 2020). Likewise, the milk from cows fed with high concentrations of S-methyl-cysteine sulfoxide (Maiga et al., 2011) also was unaffected.

In conclusion, daily supplementation with 1.5 kg of fresh broccoli or cauliflower florets in lactating sheep fed a TMR *ad libitum* did not affect either feed intake, milk production or health status of the animal. Milk quality was also not negatively affected by the inclusion of the vegetables but a slight increase in protein content was observed in broccoli-supplemented ewes.

Remains to be explored the possibility of including these vegetables in the diet of dairy sheep to replace conventional dietary components, through longer-term studies and with higher inclusion rates. Furthermore, it would be worth investigating affordable methods for their conservation and use as feed, since the high-water content of these vegetables poses challenges in terms of storage and environment, but it is also an opportunity to reuse water that would be lost without any benefit.

### **Ethics approval**

The experimental protocols were approved by the Institutional Animal Care and Use Committee of the Universidad de León (approval number ULE\_014\_2016) and the Junta de Castilla y León (Spain), following proceedings described in Spanish and European Union legislation (Royal Decree 53/2013 and Council Directive 2010/63/EU).

### **Data and model availability statement**

None of the data or models are deposited in an official repository. The data presented in this article will be available upon reasonable request to the authors.

### **Declaration of generative AI and AI-assisted technologies in the writing process**

During the preparation of this work the author(s) did not use any AI and AI-assisted technologies.

### **Author ORCIDs**

Alba Martín: 0000-0002-2531-5047; Francisco Javier Giráldez: 0000-0002-0680-1541; Iván Mateos: 0000-0002-9764-0404; Cristina Saro: 0000-0002-1887-4529; Javier Mateo: 0000-0003-4288-9867; Sonia Andrés: 0000-0001-5255-1582; Irma Caro: 0000-0002-1085-7978; María José Ranilla: 0000-0002-4000-7704.

### **Declaration of interest**

None

### **Acknowledgements**

Alba Martín gratefully acknowledges receipt of a pre-doctoral grant (PRE2019-089288) from Ministerio de Ciencia e Innovación (MCIN/AEI/10.13039/501100011033, “El FSE invierte en tu futuro”).

### **Financial support statement**

This work was funded by Ministerio de Ciencia, Innovación y Universidades (ProjectAGL2016-75322-C2-2-R, “FEDER, Una manera de hacer Europa”). Alba Martín gratefully acknowledges receipt of a pre-doctoral grant (PRE2019-089288) from Ministerio de Ciencia e Innovación.

## References

- Aghajanzpour, M., Nazer, M.R., Obeidavi, Z., Akbari, M., Ezati, P., Kor, N.M., 2017. Functional foods and their role in cancer prevention and health promotion: a comprehensive review. *American Journal of Cancer Research* 7, 740.
- Allen, M.S., 1996. Physical constraints on voluntary intake of forages by ruminants. *Journal of Animal Science* 74, 3063–3075. <https://doi.org/10.2527/1996.74123063X>
- Andrés, S., Jaramillo, E., Mateo, J., Caro, I., Carballo, D.E., López, S., Giráldez, F.J., 2019. Grain grinding size of cereals in complete pelleted diets for growing lambs: Effects on animal performance, carcass and meat quality traits. *Meat Science* 157, 107874. <https://doi.org/10.1016/j.meatsci.2019.107874>
- Araujo, C.A.S.C., Minervino, A.H.H., Sousa, R.S., Oliveira, F.L.C., Rodrigues, F.A.M.I., Mori, C.S., Ortolani, E.I. 2020. Validation of handheld  $\beta$ -hydroxybutyrate acid meter to identify hyperketonaemia in ewes. *PeerJ* 8, e8933. <https://doi.org/10.7717/peerj.8933>.
- Aziz, N.M., 2021. Effect of broccoli steam and leaves on sheep performance and milk production. *Annals of the Romanian Society for Cell Biology* 25, 10769–10774.
- Barry, T.N., 2013. The feeding value of forage brassica plants for grazing ruminant livestock. *Animal Feed Science and Technology* 181, 15–25. <https://doi.org/10.1016/j.anifeedsci.2013.01.012>
- Carrapiso, A.I., Timón, L.M., Petró, J.M., Tejeda, J.F., García, C., 2000. In situ transesterification of fatty acids from Iberian pig subcutaneous adipose tissue. *Meat Science* 56, 159–164. [https://doi.org/10.1016/S0309-1740\(00\)00035-8](https://doi.org/10.1016/S0309-1740(00)00035-8)
- Conversa, G., Lazzizzera, C., Bonasia, A., Elia, A., 2019. Growth, N uptake and N critical dilution curve in broccoli cultivars grown under Mediterranean conditions. *Scientia Horticulturae* 244, 109–121. <https://doi.org/10.1016/J.SCIENTA.2018.09.034>
- Da Silva, E. I. C. *Formulação de ração para ovinos*. 1 ed. Belo Jardim: IPA, 2021.
- Da Silva, E. I. C. *Formulação e fabricação de rações para ruminantes*. 1 ed. Belo Jardim: Emanuel Isaque Cordeiro da Silva, 2021.
- Da Silva, E. I. C. 2024. Características do leite ovino e fatores que afetam sua qualidade. *Revista Universitária Brasileira* 2(2), 2-17. <http://dx.doi.org/10.5281/zenodo.13684920>
- Da Silva, E. I. C. 2023. Cálculo de consumo e digestibilidade de alimentos e nutrientes para ruminantes. *Revista Universitária Brasileira* 1(3), 71-88. <http://dx.doi.org/10.5281/zenodo.11657180>
- de Evan, T., Marcos, C.N., Ranilla, M.J., Carro, M.D., 2020a. In vitro and in situ evaluation of broccoli wastes as potential feed for ruminants. *Animals* 10, 1–14. <https://doi.org/10.3390/ani10111989>
- de Evan, T., Vintimilla, A., Molina-Alcaide, E., Ranilla, M., Carro, M., 2020b. Potential of recycling cauliflower and romanesco wastes in ruminant feeding: in vitro studies. *Animals* 10, 1247. <https://doi.org/10.3390/ani10081247>
- De Rancourt, M., Fois, N., Lavín, M.P., Tchakérian, E., Vallerand, F., 2006. Mediterranean

- sheep and goats production: An uncertain future. *Small Ruminant Research* 62, 167–179. <https://doi.org/10.1016/J.SMALLRUMRES.2005.08.012>
- FAO, 2020. FAOSTAT [WWW Document]. Organización de las Naciones Unidas para la alimentación y la agricultura . URL <https://www.fao.org/faostat/es/#home> (accessed 9.25.23).
- Felton, C.A., DeVries, T.J., 2010. Effect of water addition to a total mixed ration on feed temperature, feed intake, sorting behavior, and milk production of dairy cows. *Journal of Dairy Science* 93, 2651–2660. <https://doi.org/10.3168/jds.2009-3009>
- Freitas, D.R. de, Souza, F.N. de, Oliveira, J.S. de, Ferreira, D. dos S., Ladeira, C.V.G., Cerqueira, M.M.O.P., 2019. Association of casein micelle size and enzymatic curd strength and dry matter curd yield. *Ciência Rural* 49, e20180409. <https://doi.org/10.1590/0103-8478cr20180409>
- Glantz, M., Devold, T.G., Vegarud, G.E., Lindmark Månsson, H., Stålhammar, H., Paulsson, M., 2010. Importance of casein micelle size and milk composition for milk gelation. *Journal of Dairy Science* 93, 1444–1451. <https://doi.org/10.3168/JDS.2009-2856>
- Humada, M.J., Sañudo, C., Serrano, E., 2014. Chemical composition, vitamin E content, lipid oxidation, colour and cooking losses in meat from Tudanca bulls finished on semi-extensive or intensive systems and slaughtered at 12 or 14 months. *Meat Science* 96, 908-915. <https://doi.org/10.1016/j.meatsci.2013.10.004>
- INRA, 2019. INRA feeding system for ruminants (2<sup>nd</sup> edition). Wageningen Academic Publishers, Wageningen, The Netherlands.
- ISO, 1999. Animal Feeding Stuffs—Determination of Moisture and Other Volatile Matter Content. International Organization for Standardization (ISO), Geneva, Switzerland.
- ISO, 2002. Animal Feeding Stuffs—Determination of Crude Ash. International Organization for Standardization (ISO), Geneva, Switzerland.
- ISO, 2009. Animal Feeding Stuffs—Determination of Nitrogen Content and Calculation of Crude Protein Content— Part 2: Block Digestion and Steam Distillation Method; International Organization for Standardization (ISO), Geneva, Switzerland.
- Jackson, P.G.G., Cockcroft, P.D., (ed.) 2002. Appendix 3: Laboratory Reference Values: Biochemistry. In *Clinical examination of farm animals*. Blackwell Science, Berlin, Germany, pp. 303-305.
- Jaime, C., Purroy, A., 1995. Level and quality of protein in rations for lactating ewes. *Annales de Zootechnie* 44, 135–142. <https://doi.org/10.1051/animres:19950204>
- Jeffery, E.H., Araya, M., 2009. Physiological effects of broccoli consumption. *Phytochemistry Reviews* 8, 283–298. <https://doi.org/10.1007/S11101-008-9106-4/FIGURES/1>
- Jin, H., Liu, Y., Guo, Z., Yang, F., Wang, J., Li, X., Peng, X., Liang, X., 2015. High-performance liquid chromatography separation of cis - Trans anthocyanin isomers from wild *Lycium ruthenicum* Murr. employing a mixed-mode reversed-phase/strong anion-exchange stationary phase. *Journal of Agricultural and Food Chemistry* 63, 500–508. [https://doi.org/10.1021/JF504525W/SUPPL\\_FILE/JF504525W\\_SI\\_001.PDF](https://doi.org/10.1021/JF504525W/SUPPL_FILE/JF504525W_SI_001.PDF)

- Li, S., Delger, M., Dave, A., Singh, H., Ye, A., 2022. Seasonal variations in the composition and physicochemical characteristics of sheep and goat milks. *Foods* 11, 1737. <https://doi.org/10.3390/foods11121737>
- Lisuzzo, A., Fiore, F., Harvatine, K., Mazzotta, E., Berlanda, M., Spissu, N., Badon, T., Contiero, B., Moscati, L., Fiore, E., 2022. Changes in plasma fatty acids profile in hyperketonemic ewes during early lactation: a preliminary study. *Scientific Reports* 12, 17017 . <https://doi.org/10.1038/s41598-022-21088-5>
- Liu, W., Stepheninbaraj, B., Chen, B., 2007. Analysis and formation of trans fatty acids in hydrogenated soybean oil during heating. *Food Chemistry* 104, 1740–1749. <https://doi.org/10.1016/j.foodchem.2006.10.069>

- Maiga, H.A., Bauer, M.L., Dahlen, C.R., Badaruddin, M., Scholljegerdes, E.J., 2011. Mustard bran in lactating dairy cow diets. *Journal of Dairy Science* 94, 3054–3062. <https://doi.org/10.3168/jds.2010-3971>
- Mestre Prates, J.A., Gonçalves Quaresma, M.A., Branquinho Bessa, R.J., Andrade Fontes, C.M.G., Mateus Alfaia, C.M.P., 2006. Simultaneous HPLC quantification of total cholesterol, tocopherols and  $\beta$ -carotene in Barrosã-PDO veal. *Food Chemistry* 94, 469–477. <https://doi.org/10.1016/j.foodchem.2005.01.021>
- Monllor, P., Muelas, R., Roca, A., Atzori, A.S., Ramón Díaz, J., Sendra, E., Romero, G., 2020a. Long-term feeding of dairy goats with broccoli by-product and artichoke plant silages: milk yield, quality and composition. *Animals* 10, 1670. <https://doi.org/10.3390/ani10091670>
- Monllor, P., Romero, G., Atzori, A.S., Sandoval-Castro, C.A., Ayala-Burgos, A.J., Roca, A., Sendra, E., Díaz, J.R., 2020b. Composition, mineral and fatty acid profiles of milk from goats fed with different proportions of broccoli and artichoke plant by-products. *Foods* 9, 700. <https://doi.org/10.3390/foods9060700>
- Monllor, P., Sandoval-Castro, C.A., Ayala-Burgos, A.J., Roca, A., Romero, G., Ramón Díaz, J., 2020c. Preference study of four alternative silage fodders from the Mediterranean region in Murciano-Granadina goats. *Small Ruminant Research* 192, 106204. <https://doi.org/10.1016/j.smallrumres.2020.106204>
- Nagraj, G.S., Chouksey, A., Jaiswal, S., Jaiswal, A.K., 2020. Broccoli, in: *Nutritional Composition and Antioxidant Properties of Fruits and Vegetables* (ed. Jaiswal, A.K.). Academic Press, London, United Kingdom, pp. 5–17.
- Owis, A.I., 2015. Broccoli; the green beauty: A review . *Journal of Pharmaceutical Sciences and Research* 7, 696–703.
- Picchi, V., Fibiani, M., Lo Scalzo, R., 2020. Cauliflower, in: *Nutritional Composition and Antioxidant Properties of Fruits and Vegetables*. *Nutritional Composition and Antioxidant Properties of Fruits and Vegetables*, (ed. Jaiswal, A.K.). Academic Press, London, United Kingdom, pp. 12-32
- Poste, L.M., Mackie, D.A., Butler, G., Larmond, Elizabeth., 1991. *Laboratory Methods for Sensory Analysis of Food*. Agriculture Canada, Ottawa, Canada.
- Prieto, N., Bodas, R., López-Campos, Ó., Andrés, S., López, S., Giráldez, F.J., 2013. Effect of sunflower oil supplementation and milking frequency reduction on sheep milk production and composition. *Journal of Animal Science* 91, 446–454. <https://doi.org/10.2527/JAS.2012-5187>
- Pulido, E., Giráldez, F.J., Bodas, R., Andrés, S., Prieto, N., 2012. Effect of reduction of milking frequency and supplementation of vitamin E and selenium above requirements on milk yield and composition in Assaf ewes. *Journal of Dairy Science* 95, 3527–3535. <https://doi.org/10.3168/jds.2011-5048>
- Pulina, G., Nudda, A., Battaccone, G., Cannas, A., 2006. Effects of nutrition on the contents of fat, protein, somatic cells, aromatic compounds, and undesirable substances in sheep milk. *Animal Feed Science and Technology* 131, 255–291. <https://doi.org/10.1016/J.ANIFEEDSCI.2006.05.023>



- Ramati, E., Israel, A., Tal Kessler, Petz-Sinuani, N., Sela, B.-A., Idan Goren, Grinfeld, A., Lavi, B., Segal, G., 2015. Low ALT activity amongst patients hospitalized in internal medicine wards is a widespread phenomenon associated with low vitamin B6 levels in their blood. *Harefuah* 154, 89–93, 137.
- Rodríguez-Palleres, X., Rojas-González, F., 2022. Valor nutricional de hojas y tallos de brócoli, apio y betarraga disponibles en un mercado mayorista de Santiago de Chile. *Memorias del Instituto de Investigaciones en Ciencias de la Salud* 20, 97–107. <https://doi.org/10.18004/mem.iics/1812-9528/2022.020.03.97>
- Satomi, S., Takahashi, S., Yoshida, K., Shimizu, S., Inoue, T., Takara, T., Suganuma, H., 2022. Effects of broccoli sprout supplements enriched in glucoraphanin on liver functions in healthy middle-aged adults with high-normal serum hepatic biomarkers: A randomized controlled trial. *Frontiers in Nutrition* 9, 1-10. <https://doi.org/10.3389/fnut.2022.1077271>
- Seguel, G., Keim, J.P., Vargas-Bello-Pérez, E., Geldsetzer-Mendoza, C., Ibáñez, R.A., Alvarado-Gilis, C., 2020. Effect of forage brassicas in dairy cow diets on the fatty acid profile and sensory characteristics of Chanco and Ricotta cheeses. *Journal of Dairy Science* 103, 228–241. <https://doi.org/10.3168/JDS.2019-17167>
- Wiedenhoef, M Hagemann, Barton, B.A., Wiedenhoef, M H, Mills, P., Louis, S., 1994. Management and environment effects on Brassica forage quality. *Agronomy Journal* 86, 227–232. <https://doi.org/10.2134/AGRONJ1994.00021962008600020003X>
- Wiedenhoef, M.H., Barton, B.A., 1995. Taste quality of milk from dairy cows fed forage Brassica cv. Tyfon. *Journal of Sustainable Agriculture* 5, 139–146. [https://doi.org/10.1300/J064v05n03\\_11](https://doi.org/10.1300/J064v05n03_11)
- Yi, X.W., Yang, F., Liu, J.X., Wang, J.K., 2015. Effects of replacement of concentrate mixture by broccoli byproducts on lactating performance in dairy cows. *Asian-Australasian Journal of Animal Sciences* 28, 1449–1453. <https://doi.org/10.5713/AJAS.15.0016>

**Table 1**

Ingredients of the total mixed ration (TMR), and chemical composition of TMR, broccoli and cauliflower used in the diet of lactating ewes.

Ingredients, g/kg	TMR
Alfalfa hay	526
Oat	76
Corn	147

Soy 44/47	134		
Sugar beet pulp	53		
Sugar beet molasses	42		
Calcium carbonate	4		
Salt	5		
Vitamin-mineral mixture	13		
<hr/>			
Chemical composition (g/kg DM)	TMR	Broccoli	Cauliflower
<hr/>			
DM (g/kg FM)	909	83.4	82.5
OM	884	887	889
CP	199	310	209
NDF	237	236	227
ADF	152	144	145
EE	9.01	37.0	22.0
<hr/>			

Abbreviations: TMR=Total mixed ration, OM: Organic matter; EE: Ether extract.

## Table 2

Fatty acid (FA) percentage and liposoluble vitamins in the experimental diets for lactating ewes [CON group (only TMR), BRO group (TMR plus 1.5 kg of broccoli) and CAU group (TMR plus 1.5 kg of cauliflower)]

Item	CON	BRO	CAU
FA (mg/g total FA)			
SFA	11.63	12.07	12.16
14:0	0.83	0.81	0.81
15:0	0.35	0.34	0.35
16:0	0	0.57	0.65
18:0	4.80	4.74	4.76
20:0	1.92	1.90	1.91
22:0	1.47	1.44	1.44
24:0	2.25	2.24	2.24
MUFA	18.45	18.31	18.31
16:1	0.76	0.74	0.74
18:1	16.76	16.57	16.62
PUFA	69.92	69.62	69.53
16:2	0	0.06	0.01
16:3	0	0.02	0.01
18:2	46.08	45.42	45.41
18:3	23.84	24.12	24.10

20:1	0.94	1.01	0.92
18:2/18:3	1.93	1.90	1.90
PUFA/SFA	6.02	5.93	5.92
Liposoluble vitamins (mg/100 g fresh matter)			
$\beta$ -carotene	1.12	1.10	1.10
$\alpha$ -tocopherol	2.81	2.80	2.82
$\gamma$ -tocopherol	0.44	0.44	0.45

The values were estimated considering 1.5 kg daily intake of fresh broccoli or cauliflower with 0.45% lipid content each and a TMR mean daily intake of 3.19 and 3.21 kg (DM) for BRO and CAU groups, respectively, with 9.01% lipid content on a DM basis.

Abbreviations: TMR= total mixed ration, SFA= total saturated fatty acids; MUFA= monounsaturated fatty acids; PUFA= polyunsaturated fatty acids.

**Table 3**

Effect of the different experimental diets [CON group (only TMR), BRO group (TMR plus 1.5 kg of broccoli) and CAU group (TMR plus 1.5 kg of cauliflower)] on feed intake, protein intake (g), live BW (kg), BW gain (kg), milk yield and feed efficiency in lactating ewes.

Parameter	Diet				P value		
	CON	BRO	CAU	SED <sup>1</sup>	Diet	Day	Diet x Day
TMR intake (kg DM/d)	3.14	3.21	3.14	0.071	0.581	0.553	1.000
Total DMI (kg/d)	3.14 <sup>a</sup>	3.33 <sup>b</sup>	3.28 <sup>ab</sup>	0.071	0.032	0.553	1.000

Protein intake (g/d)	643 <sup>a</sup>	690 <sup>b</sup>	660 <sup>a</sup>	13.9	0.006	0.555	1.000
Live BW (kg)	74.6	77.6	80.0	5.36	0.594	0.002	0.927
BW gain (kg)	4.65	4.69	7.03	-	0.544	-	-
Milk yield (L)	1.90	1.82	1.92	0.053	0.147	0.991	1.000
FCR	1.68	1.91	1.78	0.230	0.314	0.067	0.902

Abbreviations: TMR=Total mixed ration, DMI=DM intake, FCR= feed conversion ratio (kg DMI/kg milk)

<sup>1</sup>SED when comparing experimental groups.

Values within a row with different superscripts differ significantly at  $P<0.05$ .

**Table 4**

Effect of the different experimental diets [CON group (only TMR), BRO group (TMR plus 1.5 kg of broccoli) and CAU group (TMR plus 1.5 kg of cauliflower)] and sampling days (day 1, day 21 and day 42) on blood biochemical and acid-base parameters in lactating ewes.

Item	Diet				Day				P value		
	CON	BRO	CAU	SED <sup>1</sup>	1	21	42	SED <sup>2</sup>	Diet	Day	Day x diet
Albumin (g/L)	39.7	40.2	40.7	1.04	40.6	39.9	40.1	1.03	0.618	0.832	0.984
ALT (U/L)	21.6	19.7	22.3	1.18	21.3	21.8	20.4	1.17	0.092	0.498	0.984
Calcium (mg/dl)	10.6	10.8	10.8	0.13	10.6	10.7	10.8	0.13	0.225	0.263	0.994
Creatinine (mg/dL)	0.972	0.962	0.929	0.019	0.969	0.928	0.966	0.019	0.097	0.074	0.979
GGT (U/L)	59.6	58.6	63.3	2.63	63.3	59.0	59.1	2.61	0.183	0.179	0.951
Glucose (mg/dL)	68.8	69.4	69.4	1.09	67.1 <sup>a</sup>	68.9 <sup>a</sup>	71.6 <sup>b</sup>	1.09	0.828	0.001	0.182

Urea (mg/dL)	55.5 <sup>a</sup>	61.8 <sup>b</sup>	57.2 <sup>ab</sup>	2.03	64.0 <sup>b</sup>	56.4 <sup>ab</sup>	54.1 <sup>b</sup>	2.01	0.010	<0.001	0.852
BHB (mmol/L)	0.669	0.649	0.643	0.0478	0.574 <sup>a</sup>	0.658 <sup>b</sup>	0.702 <sup>b</sup>	0.0806	0.841	0.018	0.731
Lactate (µg/dL)	10.9	8.50	8.9	1.33	10.5	8.7	9.2	1.32	0.157	0.381	0.593
NEFA (mmol/L)	0.158	0.131	0.109	0.025	0.239 <sup>b</sup>	0.073 <sup>a</sup>	0.086 <sup>a</sup>	0.025	0.189	0.001	0.795
pH	7.47	7.46	7.48	0.012	7.47	7.47	7.49	0.012	0.647	0.268	0.762
pCO <sub>2</sub> (mmHg)	37.9	36.3	37.3	1.15	37.6 <sup>b</sup>	38.6 <sup>b</sup>	35.3 <sup>a</sup>	1.19	0.367	0.018	0.885
HCO <sub>3</sub> (mmol/L)	25.2	23.7	24.6	0.70	24.3	25.1	24.0	0.70	0.102	0.252	0.639
AnGap (mmol/L)	15.3	16.2	15.3	0.79	16.5 <sup>b</sup>	14.5 <sup>a</sup>	16.0 <sup>ab</sup>	0.79	0.463	0.044	0.001
tCO <sub>2</sub> (mmol/L)	26.3	24.8	25.8	0.73	25.5	26.3	25.1	0.72	0.108	0.224	0.648
Na (mmol/L)	147.5	147.2	147.7	0.38	146.9 <sup>a</sup>	148.0 <sup>b</sup>	147.4 <sup>ab</sup>	0.38	0.406	0.026	0.016
K (mmol/L)	4.68	4.77	4.63	0.107	4.94 <sup>b</sup>	4.59 <sup>a</sup>	4.54 <sup>a</sup>	0.106	0.447	0.006	0.389

Cl (mmol/L)	111.6	111.6	112.5	0.59	111.2 <sup>a</sup>	113.0 <sup>b</sup>	111.9 <sup>ab</sup>	0.89	0.390	0.009	0.089
-------------	-------	-------	-------	------	--------------------	--------------------	---------------------	------	-------	-------	-------

---

Abbreviations: TMR= total mixed ration, ALT = alanine transferase; GGT = gamma glutamyltransferase; BHB =  $\beta$ -hydroxybutyric acid; NEFA = non-esterified fatty acids; pCO<sub>2</sub>: CO<sub>2</sub> pressure; tCO<sub>2</sub>: CO<sub>2</sub> tension; AnGap = Anion gap

<sup>1</sup>SED when comparing experimental groups.

<sup>2</sup>SED when comparing the effect of time.

Values within a row with different superscripts differ significantly at  $P < 0.05$ .



**Table 5**

Effect of the different experimental diets [CON group (only TMR), BRO group (TMR plus 1.5 kg of broccoli) and CAU group (TMR plus 1.5 kg of cauliflower)] on milk quality parameters in lactating ewes.

Item	Diet					P value		
	CON	BRO	CAU	SED <sup>1</sup>	SED <sup>2</sup>	Diet	Day	Diet x day
Total solids (g/kg)	164	165	161	2.2	3.4	0.143	0.439	0.970
Fat (g/kg)	57.1	55.4	54.1	1.70	2.60	0.235	0.241	0.954
Protein (g/kg)	49.7 <sup>a</sup>	52.3 <sup>b</sup>	49.8 <sup>a</sup>	0.85	1.29	0.006	0.850	0.999
Lactose (g/kg)	47.9	47.6	47.4	0.43	0.66	0.556	0.783	0.995
Urea (mg/kg)	552	577	598	15.1	24.1	0.144	0.615	0.999
BHB (mmol/L)	0.025	0.012	0.013	0.009	0.014	0.291	0.356	0.765
Log10 somatic cells	2.00	1.96	1.98	0.251	0.980	0.258	0.299	0.409

Abbreviations: TMR= total mixed ration, BHB =  $\beta$ -hydroxybutyric acid

<sup>1</sup>SED when comparing experimental groups.

<sup>2</sup>SED when comparing the effect of time.

Values within a row with different superscripts differ significantly at  $P < 0.05$ .

**Table 6**

Effect of the different experimental diets [CON group (only TMR), BRO group (TMR plus 1.5 kg of broccoli) and CAU group (TMR plus 1.5 kg of cauliflower)] in lactating ewes on milk FA profile (% FA) and liposoluble vitamin content.

Item	Diet				P value		
	CON	BRO	CAU	SED	Diet	Week	Diet x Week
Fatty acids (%)							
SFA	73.7 <sup>a</sup>	76.0 <sup>b</sup>	75.4 <sup>b</sup>	0.71	0.012	0.458	0.641
8:0	2.34	2.32	2.44	0.073	0.225	<0.001	0.119
10:0	9.92	10.5	10.3	0.31	0.190	0.265	0.633
12:0	6.69	6.95	6.95	0.252	0.500	0.428	0.998
14:0	12.4 <sup>a</sup>	13.16 <sup>b</sup>	12.81 <sup>ab</sup>	0.300	0.038	0.030	0.945
15:0	1.03	1.03	1.07	0.038	0.443	0.409	1.000
16:0	30.1 <sup>a</sup>	32.5 <sup>b</sup>	30.5 <sup>a</sup>	0.85	0.022	0.419	0.934
17:0	0.714	0.648	0.699	0.027	0.053	0.746	0.979
18:0	7.23 <sup>b</sup>	5.68 <sup>a</sup>	6.98 <sup>b</sup>	0.358	<0.001	0.401	0.921
BCFA	2.10	1.94	2.07	0.078	0.114	0.258	0.985
iso-17:0	0.405 <sup>b</sup>	0.366 <sup>a</sup>	0.404 <sup>b</sup>	0.016	0.030	0.307	0.736
anteiso-C17:0	0.731	0.684	0.704	0.040	0.507	0.433	0.998
MUFA	20.41 <sup>b</sup>	19.26 <sup>a</sup>	19.19 <sup>a</sup>	0.611	0.003	0.492	0.680
c7-16:1	0.347	0.346	0.348	0.012	0.992	0.156	0.575
c9-16:1	0.599	0.657	0.597	0.054	0.457	0.614	0.985

t11-18:1	1.48	1.63	1.48	0.117	0.343	0.009	0.742
c9-18:1	15.7 <sup>b</sup>	13.0 <sup>a</sup>	14.0 <sup>a</sup>	0.601	0.001	0.161	0.379
c11-18:1	0.417 <sup>a</sup>	0.518 <sup>b</sup>	0.442 <sup>a</sup>	0.023	<0.001	<0.001	0.461
PUFA	5.86	5.96	5.86	0.171	0.798	<0.001	0.957
18:2n-6	3.30	3.18	3.16	0.095	0.291	0.571	0.984
18:3n-3	0.994	1.07	1.02	0.034	0.103	<0.001	0.921
c9,t11-18:2&	0.358 <sup>a</sup>	0.439 <sup>b</sup>	0.424 <sup>b</sup>	0.027	0.012	<0.001	0.802
n-6	3.76	3.61	3.61	0.107	0.303	0.658	0.981
n-3	1.07	1.15	1.11	0.035	0.095	<0.001	0.932
n-6/n-3	3.57 <sup>c</sup>	3.18 <sup>a</sup>	3.33 <sup>b</sup>	0.058	<0.001	<0.001	0.853
PUFA/SFA	0.08	0.08	0.08	0.003	0.919	<0.001	0.997
Liposoluble vitamins (mg/100g)							
α-tocopherol	106	104	90	13.1	0.442	0.097	0.858
Retinol	130	127	121	7.3	0.497	0.103	0.049

# For brevity only the fatty acids showing percentages higher than 0.4 in any of the treatment were included

Abbreviations: TMR= total mixed ration, FA = fatty acid, SFA= total saturated fatty acids, BCFA= branched-chain fatty acids, MUFA= monounsaturated fatty acids, PUFA= polyunsaturated fatty acids (including conjugated linoleic acids)

& coelutes with t7,c9-18:2.

SED when comparing experimental groups

Values within a row with different superscripts differ significantly at  $P<0.05$ .

**Table 7**

Effect of the different experimental diets [CON group (only TMR), BRO group (TMR plus 1.5 kg of broccoli) and CAU group (TMR plus 1.5 kg of cauliflower)] on micelle size and size distribution properties in ewe milk.

Item	Diet			SED <sup>1</sup>	P value
	CON	BRO	CAU		
Z-Average (nm)	192	203	189	6.1	0.066
Size for the 50 the percentile (nm)	154	154	143	10.7	0.489
Polydispersity index	0.092 <sup>a</sup>	0.105 <sup>b</sup>	0.106 <sup>b</sup>	0.0058	0.040
Z-Potential (mV)	-22.94	-23.38	-23.78	0.932	0.667

Abbreviations: TMR= total mixed ration

<sup>1</sup>SED when comparing experimental groups.

Values within a row with different superscripts differ significantly at  $P < 0.05$ .

### Figure captions

**Fig. 1.** Distribution of micelle by size (frequency in % vs Z-average in nm) in the ewe milk from the control, broccoli and cauliflower groups. Blue: control; orange: broccoli; grey: cauliflower

