

On Fuzzy b-Subimplicative Ideal

Suad Abdulaali Neamah

Department of Mathematics, Faculty of Education for Girls
 University of Kufa, Iraq

E-mail: suada.shabib@uokufa.edu.iq or soshabib@yahoo.com

Abstract: In this paper, we study a new notion of fuzzy subimplicative ideal of a BH-algebra, namely fuzzy subimplicative ideal with respect to an element in BH-algebra is introduced and some related properties are investigated.

Keywords: BH-algebra, fuzzy subimplicative ideal, subimplicative ideal, fuzzy b- subimplicative ideal, Level subset,

1. PRELIMINARIES :

In this section, is devoted to some basic ordinary concepts of BH-algebra, fuzzy ideal, sub-implicative ideal in fuzzy and ordinary, level subset, image and preimage of fuzzy set and homomorphism in BH-algebra, we give some basic concepts about the image of function, the inverse image of a BH-algebra with some remarks in fuzzy senses.

Definition (1.1): [9] A **BH-algebra** is a nonempty set X with a constant 0 and a binary operation $*$ satisfying the following conditions:

- i. $x * x = 0, \forall x \in X.$
- ii. $x * y = 0$ and $y * x = 0$ imply $x = y, \forall x, y \in X.$
- iii. $x * 0 = x, \forall x \in X.$

Remark (1.2): [10] Let X and Y be BH-algebras. A mapping $f: X \rightarrow Y$ is called a **homomorphism** if $f(x*y) = f(x)*f(y), \forall x, y \in X.$ A homomorphism f is called a **monomorphism** (resp., **epimorphism**) if it is injective (resp., surjective). A bijective homomorphism is called an isomorphism. Two BH-algebras X and Y are said to be **isomorphic**, written $X \cong Y,$ if there exists an isomorphism $f: X \rightarrow Y.$ For any homomorphism $f: X \rightarrow Y,$ the set $\{x \in X: f(x)=0'\}$ is called the **kernel** of $f,$ denoted by $\ker(f),$ and the set $\{f(x): x \in X\}$ is called the **image** of $f,$ denoted by $\text{Im}(f).$ Notice that $f(0)=0', \forall$ homomorphism $f.$

Definition (1.3): [1] if $\{A_\alpha, \alpha \in \Lambda\}$ is a family of fuzzy sets in $X,$ then :

$$\bigcap_{i \in I} A_i(x) = \inf \{ A_i(x), i \in I \}, \forall x \in X.$$

$$\bigcup_{i \in I} A_i(x) = \sup \{ A_i(x), i \in I \}, \forall x \in X. \text{ which are also fuzzy sets in } X.$$

Definition (1.4): [2] Let X and Y be any two sets, A be any fuzzy set in X and $f: X \rightarrow Y$ be any function. The set $f^{-1}(y) = \{x \in X \mid f(x) = y\}, \forall y \in Y.$ The fuzzy set B in Y defined by $B(y) = \begin{cases} \sup\{A(x) \mid x \in f^{-1}(y)\}; & \text{if } f^{-1}(y) \neq \emptyset \\ 0 & \text{otherwise} \end{cases}, \forall y \in Y,$ is called the **image** of A under f and is denoted by $f(A).$

Definition (1.5): [2] Let X and Y be any two sets, $f: X \rightarrow Y$ be any function and B be any fuzzy set in $f(A)$. The fuzzy set A in X defined by: $A(x)=B(f(x)), \forall x \in X$ is called the **preimage** of B under f and is denoted by $f^{-1}(B)$.

Definition(1.6): [6] A fuzzy subset A of a BH-algebra X is said to be a **fuzzy ideal** if and only if:

- i. $A(0) \geq A(x), \forall x \in X$.
- ii. $A(x) \geq \min \{A(x*y), A(y)\}, \forall x, y \in X$.

Definition(1.7):[5] A nonempty subset I of a BH-algebra X is called **subimplicative ideal** of X if:

- i. $0 \in I$.
- ii. $((x*(x*y))*(y*x))*z \in I$ and $z \in I$ imply $y*(y*x) \in I, \forall x, y, z \in X$.

Definition (1.8): [5] Let X be a BH-algebra and $b \in X$, a fuzzy subset A of X is called a **sub-implicative ideal with respect to an element b** (or briefly, **b -subimplicative ideal**) of X if it satisfies:

- i. $0 \in I$.
- ii. $((x*(x*y))*(y*x))*z \in I$ and $z \in I$ imply $y*(y*x) \in I, \forall x, y, z \in X$.

Definition(1.9): [5] A fuzzy set A of a BH-algebra X is called a **fuzzy subimplicative ideal** of X if it satisfies:

- i. $A(0) \geq A(x), \forall x \in X$.
- ii. $A(y*(y*x)) \geq \min \{A(((x*(x*y))*(y*x))*z), A(z)\}, \forall x, y, z \in X$.

Proposition(1.10):[5] Let X be a BH-algebra. Then every fuzzy sub-implicative ideal of X is fuzzy ideal of X .

Definition(1.11): [7] Let μ be a fuzzy set in $X, \forall \alpha \in [0, 1]$, the set $\mu_\alpha = \{x \in X, \mu(x) \geq \alpha\}$ is called a **level subset of A** . Note that, μ_α is a subset of X in the ordinary sense.

Remark(1.12):[suad] Let A be a fuzzy subset of a BH-algebra X and $w \in X$. The set $\{x \in X \mid A(w) \leq A(x)\}$ is denoted by $\uparrow A(w)$.

2.

3. THE FUZZY SUBIMPLICATIVE IDEAL WITH RESPECT TO AN ELEMENT OF A BH-ALGEBRA.

We define the concept of a **fuzzy sub-implicative ideal with respect to an element of a BH-algebra**. We discuss some properties of this concept and link it with other types of fuzzy ideal of a BH-algebra.

Definition (2.1): Let X be a BH-algebra and $b \in X$, a fuzzy subset A of X is called a **fuzzy sub-implicative ideal with respect to an element b** (or briefly, **fuzzy b -subimplicative ideal**) of X if it satisfies:

- i. i. $A(0) \geq A(x), \forall x \in X$.
- ii. ii. $A(y*(y*x)) \geq \min \{A(((x*(x*y))*(y*x))*z)*b), A(z)\}, \forall x, y, z \in X$.

Example(2.2): Consider the BH-algebra $X=\{0,1,2,3\}$ with the following operation table:

*	0	1	2
0	0	1	2
1	1	0	2
2	2	2	0

The fuzzy subset A defined by $A(x) = \begin{cases} 0.8 ; & x = 0,1 \\ 0.5 ; & x = 2 \end{cases}$ is a **fuzzy 0-subimplicative ideal** of X.

Theorem (2.3): Let X be a BH-algebra. Then A is a fuzzy subimplicative ideal of X if and only if A is a fuzzy 0-subimplicative ideal of X.

Proof :

Let A be a fuzzy subimplicative ideal of X. Then

i. $A(0) \geq A(x)$, $\forall x \in X$. [By definition (1.9)(i)]

ii. Let x, y, z $\in X$. Then, we have

$$A(y*(y*x)) \geq \min\{A(((x*(x*y))*(y*x))*z), A(z)\} \text{ [By definition(1.5)(ii)]}$$

$$\Rightarrow \min\{A(((x*(x*y))*(y*x))*z*0), A(z)\} = \min\{A(((x*(x*y))*(y*x))*z), A(z)\}$$

[Since X is a BH-algebra; $x*0=x$, $\forall x \in X$.]

$$\Rightarrow A(y*(y*x)) \geq \min\{A(((x*(x*y))*(y*x))*z*0), A(z)\}$$

Therefore, A is a fuzzy 0-subimplicative ideal of X.

Conversely,

Let A be a fuzzy 0-subimplicative ideal of X. Then

i. $A(0) \geq A(x)$, $\forall x \in X$. [By definition (2.1)(i)]

ii. Let x, y, z $\in X$. Then $A(y*(y*x)) \geq \min\{A(((x*(x*y))*(y*x))*z*0), A(z)\}$

[Since A is a fuzzy 0-subimplicative ideal of X. By definition (2.1)(ii)]

$$\Rightarrow \min\{A(((x*(x*y))*(y*x))*z*0), A(z)\} = \min\{A(((x*(x*y))*(y*x))*z), A(z)\}$$

[Since X is a BH-algebra; $x*0=x$, $\forall x \in X$]

$$\Rightarrow A(y*(y*x)) \geq \min\{A(((x*(x*y))*(y*x))*z), A(z)\}.$$

Therefore, A is a fuzzy subimplicative ideal of X. ■

Proposition (2.4): Let X be a BH-algebra, $b \in X$ and A be a fuzzy b-subimplicative ideal of X, such that $A(b)=A(0)$. Then A is a fuzzy ideal of X.

Proof :

Let A be a fuzzy b-subimplicative ideal of X. To prove A is a fuzzy ideal of X.

i. $A(0) \geq A(x)$, $\forall x \in X$. [Since A is a fuzzy b-sub-implicative ideal of X.]

ii. Let x, y, z $\in X$ such that

$$A(x*b) = A((x*0)*b) \text{ [Since X is BH-algebra; } x*0=x]$$

$$= A((x*0)*0)*b) \text{ [Since X is BH-algebra; } x*x=0]$$

$$= A(((x*(x*x))*(x*x))*0)*b \text{ [Since X is BH-algebra; } x*x=0]$$

$$\Rightarrow A(x*(x*x)) \geq \min \{A(((x*(x*x))*(x*x))*0)*b, A(b)\}=A(x*b)$$

[Since A is a fuzzy b-sub-implicative of X. By definition (2.1)(ii)]

$$\text{Now, } A(x*(x*x))=A(x*0)=A(x) \text{ [Since X is BH-algebra ; } x*x=0, x*0=x]$$

$$\Rightarrow A(x) \geq \min \{A(((x*(x*x))*(x*x))*0)*b, A(0)\}=\min \{A(x*b), A(0)\}$$

$$\Rightarrow A(x) \geq \min \{A(x*b), A(b)\}. \text{ [Since } A(b)=A(0)]$$

Therefore, A is a fuzzy ideal of X. ■

Remark(2.5): The following example shows that converse of proposition (2.4) is not correct, $\forall b \in X$.

Example (2.6): Consider the BH-algebra $X= \{0, 1, 2, 3\}$ with the binary operation '*' defined by the following table:

*	0	1	2	3
0	0	0	0	0
1	1	0	0	2
2	2	2	0	1
3	3	3	3	0

The fuzzy subset A defined by $A(x) = \begin{cases} 1 & ; x = 0 \\ 0.5 & ; x = 1,2,3 \end{cases}$ is a fuzzy ideal of X, but A is not a fuzzy 0-subimplicative ideal of X.

Since

if $x=2, y=1, z=0$, then

$$\begin{aligned} A(1*(1*2))&=A(1*0)=A(1)= 0.5 < \min \{A(((2*(2*1))*(1*2))*0)*0, A(0)\} \\ &= \min \{A((2*2))*(1*2), A(0)\} \\ &= \min \{A(0), A(0)\}=A(0)=1. \end{aligned}$$

And A is not a fuzzy 1-subimplicative ideal of X. Since

if $x=2, y=1, z=0$, then

$$\begin{aligned} A(1*(1*2))&=A(1*0)=A(1)= 0.5 < \min \{A(((2*(2*1))*(1*2))*0)*1, A(0)\} \\ &= \min \{A(((2*2))*(1*2))*1, A(0)\} \\ &= \min \{A(0*1), A(0)\}= \min \{A(0), A(0)\}=A(0)=1. \end{aligned}$$

And A is not a fuzzy 2-subimplicative ideal of X. Since

if $x=2, y=1, z=0$, then

$$\begin{aligned} A(1*(1*2))&=A(1*0)=A(1)= 0.5 < \min \{A(((2*(2*1))*(1*2))*0)*2, A(0)\} \\ &= \min \{A(((2*2))*(1*2))*2, A(0)\} \\ &= \min \{A(0*2), A(0)\}= \min \{A(0), A(0)\}=A(0)=1. \end{aligned}$$

And A is not a fuzzy 3-subimplicative ideal of X. Since

if $x=2, y=1, z=0$, then

$$A(1*(1*2))=A(1*0)=A(1)= 0.5 < \min \{A(((2*(2*1))*(1*2))*0)*3, A(0)\}$$

$$= \min \{ A(((2*2))*(1*2))*3, A(0) \}$$

$$= \min \{ A(0*3), A(0) \} = \min \{ A(0), A(0) \} = A(0) = 1.$$

Therefore, A is not a fuzzy b-subimplicative ideal. $\forall b \in X$.

Proposition (2.7): Let X be a BH-algebra. A be a fuzzy subimplicative ideal of X, $b \in X$ such that $A(b) = A(0)$. Then A is a fuzzy b-subimplicative ideal of X.

Proof:

Let A be a fuzzy subimplicative ideal of X. Then

- i. $A(0) \geq A(x), \forall x \in X$ [By definition (1.9)(i)]
- ii. Let $x, y, z \in X$. Then

$$A(y*(y*x)) \geq \min \{ A((((x*(x*y))*(y*x))*z), A(z) \}$$

$$\geq \min \{ \min \{ A((((x*(x*y))*(y*x))*z)*b), A(b) \}, A(z) \}$$

$$= \min \{ A((((x*(x*y))*(y*x))*z)*b), A(z) \}$$

[Since A is a fuzzy ideal of X. By proposition (1.10) and $A(b) = A(0)$]

Therefore, A is fuzzy b-subimplicative ideal of X. ■

Theorem (2.8): Let X be a BH-algebra. Then a fuzzy ideal A of X satisfying the condition:

$$\forall x, y \in X ; A(y*(y*x)) \geq A((x*(x*y))*(y*x)) \quad (b_1)$$

is a fuzzy b-subimplicative ideal of X, where $b \in X$ and $A(b) = A(0)$.

Proof:

Let A be a fuzzy ideal of X. Then, we have

- i. $A(0) \geq A(x), \forall x \in X$. [By definition (1.6)(i)]
- ii. Let $x, y, z \in X$. Then, we have

$$A((x*(x*y))*(y*x)) \geq \min \{ A(((x*(x*y))*(y*x))*z), A(z) \}$$

[Since A is a fuzzy ideal of X. By definition (1.6)(ii)]

$$\geq \min \{ \min \{ A((((x*(x*y))*(y*x))*z)*b), A(b) \}, A(z) \}$$

[Since A is a fuzzy ideal of X.]

$$= \min \{ A((((x*(x*y))*(y*x))*z)*b), A(z) \}$$

[Since $A((((x*(x*y))*(y*x))*z)*b) = \min \{ A((((x*(x*y))*(y*x))*z)*b), A(b) \}$

and $A(b) = A(0)$]

$$\Rightarrow A(y*(y*x)) \geq \min \{ A((((x*(x*y))*(y*x))*z)*b), A(z) \}$$
 [By the condition (b1)]

Then A is a fuzzy b-subimplicative ideal of X, $A(b) = A(0)$. ■

Theorem (2.9): If X is a BH-algebra of X satisfies the condition:

$$\forall x, y \in X ; y*(y*x) = (x*(x*y))*(y*x) \quad (b_2),$$

then every fuzzy ideal of X is a fuzzy b-subimplicative ideal of X, where $b \in X$ and $A(b) = A(0)$.

Proof:

Let A be a fuzzy ideal of X. Then, we have

- i. $A(0) \geq A(x), \forall x \in X$. [By definition (1.6)(i)]
- ii. Let $x, y, z \in X$. Then $A((x*(x*y))*(y*x)) \geq \min \{ A(((x*(x*y))*(y*x))*z), A(z) \}$ [Since A is a fuzzy ideal of X. By definition (1.6)(ii)]

Now, $A(y^*(y^*x))=A((x^*(x^*y))^*(y^*x))$ [By (b2)]

$$\Rightarrow A(y^*(y^*x)) \geq \min \{A(((x^*(x^*y))^*(y^*x))^*z), A(z)\}$$

$\Rightarrow A$ is a fuzzy subimplicative ideal of X . [By definition (1.9)]

Therefore, A is a fuzzy b-subimplicative ideal of X . [By proposition (2.7)]. ■

Theorem (2.10) : Let X be a BH-algebra, $b \in X$ and A be a fuzzy b-subimplicative ideal of X . Then the set X_A is a b-subimplicative ideal of X .

Proof:

Let A be a fuzzy b-subimplicative ideal of X . To prove X_A is a b-subimplicative ideal of X .

i. $A(x) = A(0)$.

If $x=0$, then $0 \in X_A$

ii. Let $x, y, z, b \in X$ such that $((x^*(x^*y))^*(y^*x))^*z \in X_A$ and $z \in X_A$

$$\Rightarrow A(((x^*(x^*y))^*(y^*x))^*z) = A(0) \text{ and } A(z) = A(0)$$

\Rightarrow by definition of fuzzy b-subimplicative ideal of X , we have

$$A(y^*(y^*x)) \geq \min \{A(((x^*(x^*y))^*(y^*x))^*z), A(z)\}$$

$$= \min \{A(0), A(0)\} = A(0)$$

$\Rightarrow A(y^*(y^*x)) \geq A(0)$. But $A(0) \geq A(x)$. [Since A is a fuzzy b-subimplicative ideal of X .]

$$\Rightarrow A(y^*(y^*x)) = A(0)$$

$\Rightarrow y^*(y^*x) \in X_A$. Therefore, X_A is a b-subimplicative ideal of X .

Proposition (2.11): Let $\{A_\alpha | \alpha \in \lambda\}$ be a family of fuzzy b-subimplicative ideals of a BH-algebra X . Then $\bigcap_{\alpha \in \lambda} A_\alpha$ is a fuzzy b-subimplicative ideal of X . $\forall b \in X$

Proof: Let $\{A_\alpha | \alpha \in \lambda\}$ be a family of fuzzy b-subimplicative ideals of X .

i. Let $x \in X$. Then

$$\bigcap_{\alpha \in \lambda} A_\alpha(0) = \inf \{A_\alpha(0) | \alpha \in \lambda\} \geq \inf \{A_\alpha(x) | \alpha \in \lambda\} = \bigcap_{\alpha \in \lambda} A_\alpha(x)$$

[Since A_α is a fuzzy b-subimplicative ideals of X , $\forall \alpha \in \lambda$. By definition (2.1)(i)]

$$\Rightarrow \bigcap_{\alpha \in \lambda} A_\alpha(0) \geq \bigcap_{\alpha \in \lambda} A_\alpha(x)$$

$$\text{ii. Let } x, y, z \in X. \text{ Then, we have } \bigcap_{\alpha \in \lambda} A_\alpha(y^*(y^*x)) = \inf \{A_\alpha(y^*(y^*x)) | \alpha \in \lambda\}$$

$\geq \inf \{\min \{A_\alpha(((x^*(x^*y))^*(y^*x))^*z), A_\alpha(z) | \alpha \in \lambda\}\}$ [Since A_α is a fuzzy b-subimplicative ideals of X , $\forall \alpha \in \lambda$. By definition (2.1)(ii)]

$$= \min \{ \inf \{ A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), A_{\alpha}(z) \mid \alpha \in \lambda \} \}$$

$$= \min \{ \inf \{ A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b) \mid \alpha \in \lambda \}, \inf \{ A_{\alpha}(z) \mid \alpha \in \lambda \} \} \quad = \min \left\{ \bigcap_{\alpha \in \lambda} A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), \right.$$

$$\left. \bigcap_{\alpha \in \lambda} A_{\alpha}(z) \right\}$$

$$\Rightarrow \bigcap_{\alpha \in \lambda} A_{\alpha}(y * (y * x)) \geq \min \left\{ \bigcap_{\alpha \in \lambda} A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), \bigcap_{\alpha \in \lambda} A_{\alpha}(z) \right\}$$

Therefore, $\bigcap_{\alpha \in \lambda} A_{\alpha}$ is a fuzzy b-subimplicative ideal of X. $\forall b \in X$ ■

Proposition (2.12): Let $\{A_{\alpha} \mid \alpha \in \lambda\}$ be a chain of fuzzy b-subimplicative ideals of a BH-algebra X. Then $\bigcup_{\alpha \in \lambda} A_{\alpha}$ is a fuzzy b-subimplicative ideal of X. $\forall b \in X$.

Proof: Let $\{A_{\alpha} \mid \alpha \in \lambda\}$ be a chain of fuzzy b-subimplicative ideal of X.

$$i. \text{ Let } x \in X. \text{ Then } \bigcup_{\alpha \in \lambda} A_{\alpha}(0) = \sup \{ A_{\alpha}(0) \mid \alpha \in \lambda \} \geq \sup \{ A_{\alpha}(x) \mid \alpha \in \lambda \} = \bigcup_{\alpha \in \lambda} A_{\alpha}(x)$$

[Since A_{α} is a fuzzy b-subimplicative ideal of X, $\forall \alpha \in \lambda$. By definition(2.1)(i)]

$$\Rightarrow \bigcup_{\alpha \in \lambda} A_{\alpha}(0) \geq \bigcup_{\alpha \in \lambda} A_{\alpha}(x)$$

$$ii. \text{ Let } x, y, z \in X. \text{ Then, we have } \bigcup_{\alpha \in \lambda} A_{\alpha}(y * (y * x)) = \sup \{ A_{\alpha}(y * (y * x)) \mid \alpha \in \lambda \}$$

$$\geq \sup \{ \min \{ A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), A_{\alpha}(z) \mid \alpha \in \lambda \} \}$$

[Since A_{α} is a fuzzy b-subimplicative ideals of X, $\forall \alpha \in \lambda$. By definition (2.1)(ii)]

$$= \min \{ \sup \{ A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), A_{\alpha}(z) \mid \alpha \in \lambda \} \} \text{ [Since } A_{\alpha} \text{ is a chain]}$$

$$= \min \{ \sup \{ A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b) \mid \alpha \in \lambda \}, \sup \{ A_{\alpha}(z) \mid \alpha \in \lambda \} \}$$

$$= \min \left\{ \bigcup_{\alpha \in \lambda} A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), \bigcup_{i \in \Gamma} A_{\alpha}(z) \right\}$$

$$\Rightarrow \bigcup_{\alpha \in \lambda} A_{\alpha}(y * (y * x)) \geq \min \left\{ \bigcup_{\alpha \in \lambda} A_{\alpha}(((x * (x * y)) * (y * x)) * z) * b), \bigcup_{\alpha \in \lambda} A_{\alpha}(z) \right\}$$

Therefore, $\bigcup_{\alpha \in \lambda} A_{\alpha}$ is a fuzzy b-subimplicative ideal of X. $\forall b \in X$ ■

Proposition(2.13): Let $f: (X, *, 0) \rightarrow (Y, *, 0')$ be a BH-epimorphism. If A is a fuzzy b-subimplicative ideal of X, then $f(A)$ is a fuzzy b-subimplicative ideal of Y. $\forall b \in X$

Proof: Let A be a fuzzy b-subimplicative ideal of X. Then

i. Let $y \in Y$. Then there exists $x \in X$.

$$(f(A))(0') = \sup \{A(x_1) \mid x_1 \in f^{-1}(0')\}$$

$$= A(0) \geq \sup \{A(x) \mid x \in X\} \geq \sup \{A(x_1) \mid x = f^{-1}(y)\} = (f(A))(y)$$

[Since A is a fuzzy b-subimplicative ideal of X. By definition (2.1)(i)]

$$\Rightarrow (f(A))(0') \geq (f(A))(y), \forall y \in Y.$$

iii. Let $y_1, y_2, y_3 \in Y$. Then there exist

$$f(x_1) = y_1, f(x_2) = y_2, f(z) = y_3, f(b) = y_4 \text{ such that } x_1, x_2, z, b \in X$$

$$(f(A))(y_2 * (y_2 * y_1)) = \sup \{A(x_2 * (x_2 * x_1)) \mid x_2 * (x_2 * x_1) \in f^{-1}((y_2 * (y_2 * y_1)))\}$$

$$\geq \min \{ \sup \{A(((x_1 * (x_1 * x_2)) * (x_2 * x_1)) * z) * b), A(z) \mid (((x_1 * (x_1 * x_2)) * (x_2 * x_1)) * z) * b \in$$

$$f^{-1}(((y_1 * (y_1 * y_2)) * (y_2 * y_1)) * y_3) * y_4) \text{ and } z \in f^{-1}(y_3) \}$$

[Since A is a fuzzy b-subimplicative ideal of X. By definition (2.1)(ii)]

$$\geq \min \{ \sup \{A(((x_1 * (x_1 * x_2)) * (x_2 * x_1)) * z) * b) \mid ((x_1 * (x_1 * x_2)) * (x_2 * x_1)) * z * b \in$$

$$f^{-1}(((y_1 * (y_1 * y_2)) * (y_2 * y_1)) * y_3) * y_4) \}, \sup \{A(z) \mid z \in f^{-1}(y_3) \}$$

$$= \min \{ ((f(A))(((f(x_1 * (x_1 * x_2)) * (x_2 * x_1)) * z) * b)), (f(A))(f(z)) \}$$

$$= \min \{ ((f(A))(((f(x_1) * f(x_1 * x_2)) * f(x_2 * x_1)) * f(z)) * f(b)), (f(A))(f(z)) \}$$

[Since f is an epimorphism. By remark (1.2)]

$$= \min \{ ((f(A))(((y_1 * (y_1 * y_2)) * (y_2 * y_1)) * y_3) * y_4), (f(A))(y_3) \}$$

$$\Rightarrow (f(A))(y_2 * (y_2 * y_1)) \geq \min \{ ((f(A))(((y_1 * (y_1 * y_2)) * (y_2 * y_1)) * y_3) * y_4), (f(A))(y_3) \}$$

Therefore, $f(A)$ is a fuzzy b-subimplicative ideal of Y. ■

Proposition (2.14):

Let X be a BH-algebra and A be a fuzzy subset of X. Then A is a fuzzy b-subimplicative ideal of X if and only if $A^{\#}(x) = A(x) + 1 - A(0)$ is a fuzzy b-subimplicative ideal of X, where $b \in X$.

Proof:

Let A be a fuzzy b-subimplicative ideal of X. Then

i. $A^\#(0) = A(0) + 1 - A(0)$

$\Rightarrow A^\#(0) = 1$. Then $A^\#(0) \geq A^\#(x), \forall x \in X$

ii. Let $x, y, z \in X$ and $b \in X$. Then

$A^\#(y^*(y^*x)) = A(y^*(y^*x)) + 1 - A(0)$

$\geq \min \{A(((x^*(x^*y))^*(y^*x))^*z)*b), A(z)\} + 1 - A(0)$

[Since A is a fuzzy b-subimplicative ideal of X . By definition (2.1)(ii)]

$= \min \{A(((x^*(x^*y))^*(y^*x))^*z)*b) + 1 - A(0), A(z) + 1 - A(0)\}$

$\geq \min \{A^\#(((x^*(x^*y))^*(y^*x))^*z)*b), A^\#(z)\}$

$\therefore A^\#(y^*(y^*x)) \geq \min \{A^\#(((x^*(x^*y))^*(y^*x))^*z)*b), A^\#(z)\}$

$\Rightarrow A^\#$ is a fuzzy b-subimplicative ideal of X .

Conversely,

Let $A^\#$ be a fuzzy b-subimplicative ideal of X .

i. Let $x \in X$. Then we have

$A(0) = A^\#(0) - 1 + A(0) \geq A^\#(0) - 1 + A(0) = A(x)$

[Since $A^\#$ be a fuzzy b-subimplicative ideal of X . By definition (2.1)(i)]

$\Rightarrow A(0) \geq A(x), \forall x \in X$.

ii. Let $x, y, z \in X$ and $b \in X$. Then

$A(y^*(y^*x)) = A^\#(y^*(y^*x)) - 1 + A(0) \geq \min \{A^\#(((x^*(x^*y))^*(y^*x))^*z)*b), A^\#(z)\} - 1 + A(0)$

[Since $A^\#$ is a fuzzy b-subimplicative ideal of X . By definition (2.1)(ii)]

$= \min \{A^\#(((x^*(x^*y))^*(y^*x))^*z)*b) - 1 + A(0), A^\#(z) - 1 + A(0)\}$

$\geq \min \{A(((x^*(x^*y))^*(y^*x))^*z)*b), A(z)\}$

$\Rightarrow A(y^*(y^*x)) \geq \min \{A(((x^*(x^*y))^*(y^*x))^*z)*b), A(z)\}$

Then A is a fuzzy b-subimplicative ideal of X . ■

Proposition(2.15) Let X be a BH-algebra and let $w, b \in X$. If A is a fuzzy b-sub-implicative ideal of X , then $\uparrow A(w)$ is a b-subimplicative ideal of X .

Proof:

Let A be a fuzzy b-subimplicative ideal of X . Then

i. $A(0) \geq A(x), \forall x \in X$.

[Since A is a fuzzy b-subimplicative ideal of X . By definition (2.1)(i)]

$\Rightarrow A(0) \geq A(w) \Rightarrow 0 \in \uparrow A(w)$

ii. Let $x, y, z \in X$ such that $((x^*(x^*y))^*(y^*x))^*z)*b \in \uparrow A(w)$ and $z \in \uparrow A(w)$

$\Rightarrow A(w) \leq A(((x^*(x^*y))^*(y^*x))^*z)*b)$ and $A(w) \leq A(z)$

$\Rightarrow A(w) \leq \min \{A(((x^*(x^*y))^*(y^*x))^*z)*b), A(z)\}$

But $A(y^*(y^*x)) \geq \min \{A(((x^*(x^*y))^*(y^*x))^*z)*b), A(z)\}$

[Since A is a fuzzy b-subimplicative ideal of X . By definition (2.1)(ii)]

$\Rightarrow A(w) \leq A(y^*(y^*x))$

$\Rightarrow y^*(y^*x) \in \uparrow A(w)$.

Therefore, $\uparrow A(w)$ is a b-subimplicative ideal of X . ■

Theorem (2.16):

Let X be a BH-algebra, A be a fuzzy ideal of X $A(b) = A(0)$. Then A is a fuzzy b-subimplicative ideal of X if and only if A_α is a b-subimplicative ideal of X, $\forall \alpha \in [0, A(0)]$.

Proof:

Let A be a fuzzy b-subimplicative ideal of X. To prove A_α is a b-subimplicative ideal of X.

i. Let $x \in A_\alpha$. Then $A(x) \geq \alpha$ [By definition (1.11) of A_α]

But $A(0) \geq A(x)$. [Since A is a fuzzy b-subimplicative ideal of X. By definition(2.1)(i)]

$\Rightarrow A(0) \geq \alpha$.

$\Rightarrow 0 \in A_\alpha$.

ii. Let $x, y, z, b \in X$ such that $((x*(x*y))*(y*x))*z*b \in A_\alpha$ and $z \in A_\alpha$.

$\Rightarrow A(((x*(x*y))*(y*x))*z*b) \geq \alpha$ and $A(z) \geq \alpha$ [By definition(1.11) of A_α]

$\min \{ A(((x*(x*y))*(y*x))*z*b), A(z) \} \geq \alpha$

But $A(y*(y*x)) \geq \min \{ A(((x*(x*y))*(y*x))*z*b), A(z) \}$

[Since A is a fuzzy b-subimplicative ideal of X. By definition (2.1)(ii)]

$\Rightarrow A(y*(y*x)) \geq \alpha \Rightarrow y*(y*x) \in A_\alpha$ [By definition (1.11) of A_α]

Therefore, A_α is a b-subimplicative ideal of X, $\forall b \in A_\alpha$.

Conversely,

To prove A is a fuzzy b-subimplicative ideal of X.

i. $0 \in A_\alpha$. [By definition (1.8)(i)]. Then $A(0) \geq \alpha = A(x), \forall x \in X$.

ii. Let $x, y, z \in X$ such that $\alpha = \min \{ A(((x*(x*y))*(y*x))*z*b), A(z) \}$

$\Rightarrow A(((x*(x*y))*(y*x))*z*b) \geq \alpha$ and $A(z) \geq \alpha$

$\Rightarrow ((x*(x*y))*(y*x))*z*b \in A_\alpha$ and $z \in A_\alpha$.

$\Rightarrow y*(y*x) \in A_\alpha$ [Since A_α is a b-subimplicative ideal of X. By definition (1.8)(ii)]

$\Rightarrow A(y*(y*x)) \geq \alpha$

$\Rightarrow A(y*(y*x)) \geq \min \{ A(((x*(x*y))*(y*x))*z*b), A(z) \}$

Therefore, A is a fuzzy b-subimplicative ideal of X, $A(b) = A(0)$. ■

References

1. D. Dubois and H. Prade, "Fuzzy Sets and Systems", Academic Press. INC. (London) LTD, fifth Avenue, New Yurk, 1980.
2. E. M.kim and S.S. Ahn," On Fuzzy n-fold Strong Ideals of BH-algebras", J. Appl. Math. & Informatics. 30,(No.3) – 4,pp. 665 - 676,2012
3. H. H. Abbass and H. A. Dahham,"Some Types of Fuzzy Ideals with Respect to an Element of a BG-algebra", M.S.c thesis, Kufa University, 2012.
4. H. H. Abbass and H. M. A. Saeed, "The Fuzzy Closed BCH-algebra with Respect to an Element of a BH-algebra", M.S.c thesis, Kufa University, 2011.
5. H. H. Abbass and S. A. Neamah. Saeed, "The Fuzzy Implicative Ideal with Respect to an Element of a BH-algebra", M.S.c thesis, Kufa University, 2014.

6. H. H. Abbass, "The Q-Smarandache Closed Ideal and Q-Smarandache Fuzzy Closed Ideal with Respect to an Element of a Q-Smarandache BCH-algebra", Journal kerbala university, vol.10, No.3 scientific, pp.302-312,2012.
7. M. Ganesh, "Introduction to Fuzzy Sets and Fuzzy Logic", fourth printing, 2009.
8. Q. Zhang, E. H. Roh and Y. B. Jun, "On Fuzzy BH-algebras", J. Huanggang, Normal Univ. 21(3), pp. 14–19, 2001.
9. Y. B. Jun, E. H. Roh and H. S. Kim, "On BH-algebras", Scientiae Mathematicae 1(1), pp. 347–354, 1998.
10. Y. B. Jun, H. S. Kim and M. Kondo, " On BH-relations in BH-algebras", Scientiae Mathematicae Japonicae Online, Vol.9, pp. 91–94, 2003.]