



# Introduction to Image Processing via Neutrosophic Techniques

A. A. Salama<sup>1</sup>, Florentin Smarandache<sup>2</sup> and Mohamed Eisa<sup>3</sup>

<sup>1</sup> Department of Mathematics and Computer Science, Faculty of Sciences, Port Said University, 23 December Street, Port Said 42522, Egypt.  
Email: drsalama44@gmail.com

<sup>2</sup> Department of Mathematics, University of New Mexico 705 Gurley Ave. Gallup, NM 87301, USA.  
Email: smarand@unm.edu

<sup>3</sup> Computer Science Department, Port Said University, 42526 Port Said, Egypt.  
Email: mmmeisa@yahoo.com

**Abstract.** This paper is an attempt of proposing the processing approach of neutrosophic technique in image processing. As neutrosophic sets is a suitable tool to cope with imperfectly defined images, the properties, basic operations distance measure, entropy measures, of the neutrosophic sets method are presented here. In this paper we, introduce the distances between neutrosophic sets: the Hamming distance, the normalized Hamming

distance, the Euclidean distance and normalized Euclidean distance. We will extend the concepts of distances to the case of neutrosophic hesitancy degree. Entropy plays an important role in image processing. In our further considerations on entropy for neutrosophic sets the concept of cardinality of a neutrosophic set will also be useful. Possible applications to image processing are touched upon.

**Keywords:** Neutrosophic sets; Hamming distance; Euclidean distance; Normalized Euclidean distance; Image processing.

## 1. Introduction

Since the world is full of indeterminacy, the neutrosophics found their place into contemporary research. Smarandache [9, 10] and Salama et al [ 4, 5, 6, 7, 8, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 45]. Entropy plays an important role in image processing. In this paper we, introduce the distances between neutrosophic sets: the Hamming distance. In this paper we, introduce the distances between neutrosophic sets: the Hamming distance, The normalized Hamming distance, the Euclidean distance and normalized Euclidean distance. We will extend the concepts of distances to the case of neutrosophic hesitancy degree. In our further considerations on entropy for neutrosophic sets the concept of cardinality of a neutrosophic set will also be useful.

## 2. Terminologies

Neutrosophy has laid the foundation for a whole family of new mathematical theories generalizing both their

classical and fuzzy counterparts [1, 2, 3, 11, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 46] such as a neutrosophic set theory. We recollect some relevant basic preliminaries, and in particular, the work of Smarandache in [9, 10] and Salama et al. [4, 5, 6, 7, 8, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 45]. Smarandache introduced the neutrosophic components T, I, F which represent the membership, indeterminacy, and non-membership values respectively, where  $]0^-, 1^+[$  is nonstandard unit interval. Salama et al. introduced the following:

Let  $X$  be a non-empty fixed set. A neutrosophic set  $A$  is an object having the form  $A = \langle \mu_A(x), \sigma_A(x), \nu_A(x) \rangle$  where  $\mu_A(x)$ ,  $\sigma_A(x)$  and  $\nu_A(x)$  which represent the degree of membership function (namely  $\mu_A(x)$ ), the degree of indeterminacy (namely  $\sigma_A(x)$ ), and the degree of non-membership (namely  $\nu_A(x)$ ) respectively of each element  $x \in X$  to the set  $A$  where

$$0^- \leq \mu_A(x), \sigma_A(x), \nu_A(x) \leq 1^+ \text{ and}$$

$$0^- \leq \mu_A(x) + \sigma_A(x) + \nu_A(x) \leq 3^+ .\text{Smarandache}$$

introduced the following: Let T, I, F be real standard or nonstandard subsets of  $]0^-, 1^+[$ , with

$$\text{Sup}_T = t_{\text{sup}}, \text{inf}_T = t_{\text{inf}}$$

$$\text{Sup}_I = i_{\text{sup}}, \text{inf}_I = i_{\text{inf}}$$

$$\text{Sup}_F = f_{\text{sup}}, \text{inf}_F = f_{\text{inf}}$$

$$n\text{-sup} = t_{\text{sup}} + i_{\text{sup}} + f_{\text{sup}}$$

$$n\text{-inf} = t_{\text{inf}} + i_{\text{inf}} + f_{\text{inf}},$$

T, I, F are called neutrosophic components

### 3. Distances Between Neutrosophic Sets

We will now extend the concepts of distances presented in [11] to the case of neutrosophic sets.

#### Definition 3.1

Let  $A = \{(\mu_A(x), \nu_A(x), \gamma_A(x)), x \in X\}$  and

$$B = \{(\mu_B(x), \nu_B(x), \gamma_B(x)), x \in X\}$$

$$X = \{x_1, x_2, x_3, \dots, x_n\}$$

i) The Hamming distance is equal to

$$d_{Ns}(A, B) = \sum_{i=1}^n (|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\gamma_A(x_i) - \gamma_B(x_i)|)$$

ii) The Euclidean distance is equal to

$$e_{Ns}(A, B) = \sqrt{\sum_{i=1}^n ((\mu_A(x_i) - \mu_B(x_i))^2 + (\nu_A(x_i) - \nu_B(x_i))^2 + (\gamma_A(x_i) - \gamma_B(x_i))^2)}$$

iii) The normalized Hamming distance is equal to

$$NH_{Ns}(A, B) = \frac{1}{2n} \sum_{i=1}^n (|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\gamma_A(x_i) - \gamma_B(x_i)|)$$

iv) The normalized Euclidean distance is equal to

$$NE_{Ns}(A, B) = \sqrt{\frac{1}{2n} \sum_{i=1}^n ((\mu_A(x_i) - \mu_B(x_i))^2 + (\nu_A(x_i) - \nu_B(x_i))^2 + (\gamma_A(x_i) - \gamma_B(x_i))^2)}$$

#### Example 3.1

Let us consider for simplicity degenerated neutrosophic sets  $A, B, D, G, F$  in  $X = \{a\}$ . A full description of each neutrosophic set i.e.


$$A = \{(\mu_A(x), \nu_A(x), \gamma_A(x)), a \in X\}, \text{ may be exemplified}$$

$$\text{by } A = \{ \langle 1, 0, 0 \rangle, a \in X \}, B = \{ \langle 0, 1, 0 \rangle, a \in X \},$$

$$D = \{ \langle 0, 0, 1 \rangle, a \in X \}, G = \{ \langle 0.5, 0.5, 0 \rangle, a \in X \},$$

$$E = \{ \langle 0.25, 0.25, 0.05 \rangle, a \in X \} .$$

Let us calculate four distances between the above neutrosophic sets using i), ii), iii) and iv) formulas ,



(Fig.1) A geometrical interpretation of the neutrosophic considered in Example 5.1 .

$$\text{We obtain } e_{Ns}(A, D) = \frac{1}{2}, e_{Ns}(B, D) = \frac{1}{2},$$

$$e_{Ns}(A, B) = \frac{1}{2}, e_{Ns}(A, G) = \frac{1}{2}, e_{Ns}(B, G) = \frac{1}{2},$$

$$e_{Ns}(E, G) = \frac{1}{4}, e_{Ns}(D, G) = \frac{1}{4}, NE_{Ns}(A, B) = 1,$$

$$NE_{Ns}(A, D) = 1, NE_{Ns}(B, D) = 1, NE_{Ns}(A, G) = \frac{1}{2},$$

$$NE_{Ns}(B, G) = \frac{1}{2}, NE_{Ns}(B, D) = \frac{1}{2}, NE_{Ns}(E, G) = \frac{\sqrt{3}}{4}, \text{ a}$$

$$\text{nd } NE_{Ns}(D, G) = \frac{\sqrt{3}}{2},$$

From the above results the triangle ABD (Fig.1) has edges equal to  $\sqrt{2}$  and

$$e_{Ns}(A, D) = e_{Ns}(B, D) = e_{Ns}(A, B) = \frac{1}{2} \text{ and}$$

$$NE_{Ns}(A, B) = NE_{Ns}(A, D) = NE_{Ns}(B, D) =$$

$$2NE_{Ns}(A, G) = 2NE_{Ns}(B, G) = 1, \text{ and } NE_{Ns}(E, G) \text{ is}$$

equal to half of the height of triangle with all edges equal to  $\sqrt{2}$  multiplied by,  $\frac{1}{\sqrt{2}}$  i.e.  $\frac{\sqrt{3}}{4}$  .

#### Example 3.2

Let us consider the following neutrosophic sets A and B in  $X = \{a, b, c, d, e\}$  ,

$$A = \{ \langle 0.5, 0.3, 0.2 \rangle, \langle 0.2, 0.6, 0.2 \rangle, \langle 0.3, 0.2, 0.5 \rangle, \langle 0.2, 0.2, 0.6 \rangle, \langle 1, 0, 0 \rangle \}$$

$$B = \{ \langle 0.2, 0.6, 0.2 \rangle, \langle 0.3, 0.2, 0.5 \rangle, \langle 0.5, 0.2, 0.3 \rangle, \langle 0.9, 0, 0.1 \rangle, \langle 0, 0, 0 \rangle \}$$

.Then

$$d_{Ns}(A, B) = 3, NH_{Ns}(A, B) = 0.43, e_{Ns}(A, B) = 1.49$$

$$\text{and } NE_{Ns}(A, B) = 0.55.$$

#### Remark 3.1

Clearly these distances satisfy the conditions of metric space.

#### Remark 3.2

It is easy to notice that for formulas i), ii), iii) and iv) the following is valid:

$$\text{a) } 0 \leq d_{Ns}(A, B) \leq n$$

$$\text{b) } 0 \leq NH_{Ns}(A, B) \leq 1$$

$$\text{c) } 0 \leq e_{Ns}(A, B) \leq \sqrt{n}$$

$$\text{d) } 0 \leq NE_{Ns}(A, B) \leq 1.$$

This representation of a neutrosophic set (Fig. 2) will be a point of departure for neutrosophic crisp distances, and entropy of neutrosophic sets.




Fig. 2. A three-dimension representation of a neutrosophic set[9, 10].

#### 4. Hesitancy Degree and Cardinality for Neutrosophic Sets

We will now extend the concepts of distances to the case of neutrosophic hesitancy degree. By taking into account the four parameters characterization of neutrosophic sets i.e.  $A = \{ \langle \mu_A(x), \nu_A(x), \gamma_A(x), \pi_A(x) \rangle, x \in X \}$

##### Definition 4.1

Let  $A = \{ (\mu_A(x), \nu_A(x), \gamma_A(x)), x \in X \}$  and  $B = \{ (\mu_B(x), \nu_B(x), \gamma_B(x)), x \in X \}$  on  $X = \{x_1, x_2, x_3, \dots, x_n\}$

For a neutrosophic

set  $A = \{ (\mu_A(x), \nu_A(x), \gamma_A(x)), x \in X \}$  in  $X$ , we call  $\pi_A(x) = 3 - \mu_A(x) - \nu_A(x) - \gamma_A(x)$ , the neutrosophic index of  $x$  in  $A$ . It is a hesitancy degree of  $x$  to  $A$  it is obvious that  $0 \leq \pi_A(x) \leq 3$ .

##### Definition 4.2

Let  $A = \{ (\mu_A(x), \nu_A(x), \gamma_A(x)), x \in X \}$  and  $B = \{ (\mu_B(x), \nu_B(x), \gamma_B(x)), x \in X \}$  in  $X = \{x_1, x_2, x_3, \dots, x_n\}$  then

i) The Hamming distance is equal to

$$d_{Ns}(A, B) = \sum_{i=1}^n (|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\gamma_A(x_i) - \gamma_B(x_i)| + |\pi_A(x_i) - \pi_B(x_i)|)$$

. Taking into account that

$$\pi_A(x_i) = 3 - \mu_A(x_i) - \nu_A(x_i) - \gamma_A(x_i) \quad \text{and}$$

$$\pi_B(x_i) = 3 - \mu_B(x_i) - \nu_B(x_i) - \gamma_B(x_i)$$

we have

$$|\pi_A(x_i) - \pi_B(x_i)| = |3 - \mu_A(x_i) - \nu_A(x_i) - \gamma_A(x_i) - 3 + \mu_B(x_i) + \nu_B(x_i) + \gamma_B(x_i)| \\ \leq |\mu_B(x_i) - \mu_A(x_i)| + |\nu_B(x_i) - \nu_A(x_i)| + |\gamma_B(x_i) - \gamma_A(x_i)|$$

.

ii) The Euclidean distance is equal to

$$e_{Ns}(A, B) = \sqrt{\sum_{i=1}^n ((\mu_A(x_i) - \mu_B(x_i))^2 + (\nu_A(x_i) - \nu_B(x_i))^2 + (\gamma_A(x_i) - \gamma_B(x_i))^2 + (\pi_A(x_i) - \pi_B(x_i))^2)}$$

we have

$$(\pi_A(x_i) - \pi_B(x_i))^2 = \\ (-\mu_A(x_i) - \nu_A(x_i) - \gamma_A(x_i) + \mu_B(x_i) + \nu_B(x_i) + \gamma_B(x_i))^2 \\ = (\mu_B(x_i) - \mu_A(x_i))^2 + (\nu_A(x_i) - \nu_B(x_i))^2 + \\ (\gamma_A(x_i) - \gamma_B(x_i))^2 \\ + 2(\mu_B(x_i) - \mu_A(x_i))(\nu_A(x_i) - \nu_B(x_i)) \\ (\gamma_B(x_i) - \gamma_A(x_i))$$

iii) The normalized Hamming distance is equal to

$$NH_{Ns}(A, B) = \frac{1}{2n} \sum_{i=1}^n (|\mu_A(x_i) - \mu_B(x_i)| + |\nu_A(x_i) - \nu_B(x_i)| + |\gamma_A(x_i) - \gamma_B(x_i)| + |\pi_A(x_i) - \pi_B(x_i)|)$$

iv) The normalized Euclidean distance is equal to

$$NE_{Ns}(A, B) = \sqrt{\frac{1}{2n} \sum_{i=1}^n ((\mu_A(x_i) - \mu_B(x_i))^2 + (\nu_A(x_i) - \nu_B(x_i))^2 + (\gamma_A(x_i) - \gamma_B(x_i))^2 + (\pi_A(x_i) - \pi_B(x_i))^2)}$$

#### 5.2 Remark

It is easy to notice that for formulas i), ii), iii) and the following is valid:

- a)  $0 \leq d_{Ns}(A, B) \leq 2n$
- b)  $0 \leq NH_{Ns}(A, B) \leq 2$
- c)  $0 \leq e_{Ns}(A, B) \leq \sqrt{2n}$
- d)  $0 \leq NE_{Ns}(A, B) \leq \sqrt{2}$ .

#### 5. from Images to Neutrosophic Sets, and Entropy

Given the definitions of the previous section several possible contributions are discussed. Neutrosophic sets may be used to solve some of the problems of data causes problems in the classification of pixels. Hesitancy in images originates from various factors, which in their majority are due to the inherent weaknesses of the acquisition and the imaging mechanisms. Limitations of the acquisition chain, such as the quantization noise, the suppression of the dynamic range, or the nonlinear behavior of the mapping system, affect our certainty on deciding whether a pixel is “gray” or “edgy” and therefore introduce a degree of hesitancy associated with the corresponding pixel. Therefore, hesitancy should encapsulate the aforementioned sources of indeterminacy that characterize digital images. Defining the membership component of the A–NS that describes the brightness of pixels in an image, is a more straightforward task that can be carried out in a similar manner as in traditional fuzzy image processing systems. In the presented heuristic framework, we consider the membership value of a gray level  $g$  to be its normalized

intensity level; that

is  $\mu_A(g) = \frac{g}{L-1}$  where  $g \in \{0, \dots, L-1\}$ . It should be

mentioned that any other method for calculating  $\mu_A(g)$  can also be applied.

In the image is  $A$  being  $(x, y)$  the coordinates of each pixel and the  $g(x, y)$  be the gray level of the pixel  $(x, y)$  implies  $0 \leq g(x, y) \leq L-1$ . Each image pixel is associated with four numerical values:

- A value representing the membership  $\mu_A(x)$ , obtained by means of membership function associated with the set that represents the expert's knowledge of the image.
- A value representing the indeterminacy  $\nu_A(x)$ , obtained by means of the indeterminacy function associated with the set that represents the ignorance of the expert's decision.
- A value representing the non-membership  $\gamma_A(x)$ , obtained by means of the non-membership function associated with the set that represents the ignorance of the expert's decision.
- A value representing the hesitation measure  $\pi_A(x)$ , obtained by means of the  $\pi_A(x) = 3 - \mu_A(x) - \nu_A(x) - \gamma_A(x)$ .

Let an image  $A$  of size  $M \times N$  pixels having  $L$  gray levels ranging between 0 and  $L-1$ . The image in the neutrosophic domain is considered as an array of neutrosophic singletons. Here, each element denoted the degree of the membership, indeterminacy and non-membership according to a pixel with respect to an image considered. An image  $A$  in neutrosophic set is  $A = \{ \langle \mu_A(g_{ij}), \nu_A(g_{ij}), \gamma_A(g_{ij}) \rangle, g_{ij} \in \{0, \dots, L-1\} \}$

where  $\mu_A(g_{ij}), \nu_A(g_{ij}), \gamma_A(g_{ij})$  denote the degrees of membership indeterminacy and non-membership of the  $(i, j)$ -th pixel to the set  $A$  associated with an image

property  $\mu_A(g) = \frac{g - g_{\min}}{g - g_{\max}}$  where  $g_{\min}$  and  $g_{\max}$  are

the minimum and the maximum gray levels of the image. Entropy plays an important role in image processing. In our further considerations on entropy for neutrosophic sets the concept of cardinality of a neutrosophic set will also be useful

### Definition 5.1

Let  $A = \langle (\mu_A(x), \nu_A(x), \gamma_A(x)), x \in X \rangle$  a neutrosophic set in  $X$ , first, we define two cardinalities of a neutrosophic set

- The least (sure) cardinality of  $A$  is equal to so is called sigma-count, and is called here the

$$\min \sum cont(A) = \sum_{i=1} \mu_A(x_i) + \sum_{i=1} \nu_A(x_i)$$

- The biggest cardinality of  $A$ , which is possible due to  $\pi_A(x)$  is equal to

$$\max \sum cont(A) = \sum_{i=1} (\mu_A(x_i) + \pi_A(x_i)) + \sum_{i=1} \nu_A(x_i) + \pi_A(x_i)$$

and, clearly for  $A^c$  we have

$$\min \sum cont(A^c) = \sum_{i=1} \gamma_A(x_i) + \sum_{i=1} \nu_A(x_i),$$

$$\max \sum cont(A^c) = \sum_{i=1} (\gamma_A(x_i) + \pi_A(x_i)) + \sum_{i=1} \nu_A(x_i) + \pi_A(x_i)$$

. Then the cardinality of neutrosophic set is defined as the interval

$$Card(A) = [\min \sum Cont(A), \max \sum Cont(A)]$$

### Definition 5.2

An entropy on  $NS(X)$  is a real-valued functional  $E: NS(X) \rightarrow [0,1]$ , satisfying the following axiomatic requirements:

E<sub>1</sub>:  $E(A) = 0$  iff  $A$  is a neutrosophic crisp set; that is

$$\mu_A(x_i) = 0 \text{ or } \mu_A(x_i) = 1 \text{ for all } x_i \in X.$$

E<sub>2</sub>:  $E(A) = 1$  iff  $\mu_A(x_i) = \nu_A(x_i) = \gamma_A(x_i)$  for all  $x_i \in X$ . that is  $A = A^c$ .

E<sub>3</sub>:  $E(A) \leq E(B)$  if  $A$  refine  $B$ ; i.e.  $A \leq B$ .

E<sub>4</sub>:  $E(A) = E(A^c)$

Where a neutrosophic entropy measure be define as

$$E(A) = \frac{1}{n} \sum_{i=1}^n \frac{\max Count(A_i \cap A_i^c)}{\max Count(A_i \cup A_i^c)} \text{ where}$$

$n = Cardinal(X)$  and  $A_i$  denotes the single-element

$A$ -NS corresponding to the  $i^{\text{th}}$  element of the universe  $X$  and is described as

$$A_i = \{ (\mu_A(x_i), \nu_A(x_i), \gamma_A(x_i)), x_i \in X \}.$$

In other words,  $A_i$  is the  $i^{\text{th}}$  "component" of  $A$ .

Moreover,  $\max Count(A)$  denotes the biggest cardinality of  $A$  and is given by :

$$\max \sum cont(A) = \sum_{i=1} (\mu_A(x_i) + \pi_A(x_i)) + \sum_{i=1} \nu_A(x_i) + \pi_A(x_i)$$

### Conclusion

Some of the properties of the neutrosophic sets, Distance measures, Hesitancy Degree, Cardinality and Entropy measures are briefed in this paper. These measures can be used effectively in image processing and pattern recognition. The future work will cover the application of these measures.

## References

- [1] K. Atanassov, intuitionistic fuzzy sets, in V. Sgurev, ed., VII ITKRS Session, Sofia (June 1983 central Sci. and Techn. Library, Bulg. Academy of Sciences (1984).
- [2] K. Atanassov, intuitionistic fuzzy sets, *Fuzzy Sets and Systems* 2087-96, (1986).
- [3] K. Atanassov, Review and new result on intuitionistic fuzzy sets, preprint IM-MFAIS-1-88, Sofia, (1988).
- [4] I. M. Hanafy, A. A. Salama and K. Mahfouz, "Correlation Coefficient of Neutrosophic Sets by Centroid Method" *International Journal of Probability and Statistics*, 2(1),(2013)pp 9-12.
- [5] I. M. Hanafy, A. A. Salama and K. Mahfouz, "Correlation of neutrosophic Data," *International Refereed Journal of Engineering and Science (IRJES)*, Vol.(1), Issue 2, (2012)pp.39-43.
- [6] I. M. Hanafy, A.A. Salama and K. M. Mahfouz, "Neutrosophic Classical Events and Its Probability," *International Journal of Mathematics and Computer Applications Research (IJMCAR)*, Vol.(3), Issue1,(2013) pp171-178.
- [7] A.A. Salama and S.A. Alblowi, "Neutrosophic Set and Neutrosophic Topological Spaces," *ISOR J. Mathematics (IOSR-JM)*, Vol. (3). Issue (4), (2012)pp 31-35.
- [8] A. A. Salama and S.A. Alblowi, "Generalized Neutrosophic Set and Generalized Neutrosophic Topological Spaces," *Journal Computer Science and Engineering*, Vol.(2) No. (7), (2012)pp98-102.
- [9] F.Smarandache, "Neutrosophic set a generalization of the intuitionistic fuzzy sets", *Proceedings of the third conference of the European Society for fuzzy logic and Technology, EUSFLAT, Septemper Zittau Geamany Univ. of Applied Sciences at Zittau Goerlit 2*, (2003)pp141-146.
- [10] Florentin Smarandache, "Neutrosophy and Neutrosophic Logic," *First International Conference on Neutrosophy, Neutrosophic Logic, Set, Probability and Statistics University of New Mexico, Gallup, NM 87301, USA(2002)*.
- [11] Eulalia Szmids and Janus Koeprzyk, "Distance between intuitionistic fuzzy sets," *Fuzzy Sets and Systems* 114, (2000)pp 505-518.
- [12] A. A. Salama, "Neutrosophic Crisp Points & Neutrosophic Crisp Ideals," *Neutrosophic Sets and Systems*, Vol.1, No. 1,( 2013)pp 50-54.
- [13] A. A. Salama and F. Smarandache, "Filters via Neutrosophic Crisp Sets," *Neutrosophic Sets and Systems*, Vol.1, No. 1, (2013)pp34-38.
- [14] A. A. Salama, "The Concept of Neutrosophic Set and Basic Properties of Neutrosophic Set Operations," *WASET 2012 PARIS, FRANC, International University of Science, Engineering and Technology*, (2012).
- [15] A. A. Salama, Florentin Smarandache, and Valeri Kroumov, "Neutrosophic Crisp Sets & Neutrosophic Crisp Topological Spaces," *Neutrosophic Sets and Systems*, Vol. (2), (2014) pp25-30.
- [16] A. A. Salama, F. Smarandache and S. A. Alblowi, "The Characteristic Function of a neutrosophic Set," *Neutrosophic Sets and Systems*, Vol.3, (2014)pp14-18.
- [17] A. A. Salama, Mohamed Eisa and M. M. Abdelmoghny, "Neutrosophic Relations Database," *International Journal of Information Science and Intelligent System*, 3(2),(2014)pp33-46
- [18] A. A. Salama, Florentin Smarandache and Valeri Kroumov, "Neutrosophic Closed Set and Neutrosophic Continuous Functions," *Neutrosophic Sets and Systems*, Vol.(4), (2014) pp4-8.
- [19] A. A. Salama, Florentin Smarandache and S. A. Alblowi, "New Neutrosophic Crisp Topological Concepts," *Neutrosophic Sets and Systems*, Vol(4), (2014)pp50-54.
- [20] A. A. Salama, Said Broumi and Florentin Smarandache, "Neutrosophic Crisp Open Set and Neutrosophic Crisp Continuity via Neutrosophic Crisp Ideals," *I.J. Information Engineering and Electronic Business*,(3), (2014)pp1-8.
- [21] A. A. Salama, Haitham A. El-Ghareeb, Ayman M. Manie and M. M. Lotfy, "Utilizing Neutrosophic Set in Social Network Analysis e-Learning Systems," *International Journal of Information Science and Intelligent System*, Vol.(3), No.4, (2014).
- [22] A. A. Salama, Haithem A. El-Ghareeb, Ayman M. Manie and Florentin Smarandache, "Introduction to Develop Some Software Programes for dealing with Neutrosophic Sets," *Neutrosophic Sets and Systems*, Vol.(3),(2014) pp51-52.
- [23] A. A. Salama, Mohamed Abdelfattah, Mohamed Eisa, "Distances, Hesitancy Degree

- and Flexible Querying via Neutrosophic Sets, International Journal of Computer Application (IJCA) , Issue4, Vol.(3): May 2014.
- [24] A. A. Salama, Said Broumi and Florentin Smarandache, Some Types of Neutrosophic Crisp Sets and Neutrosophic Crisp Relations , I.J. Information Engineering and Electronic Business, (2014) ,(Accepted).
- [25] A. A. Salama and Said Broumi, Roughness of neutrosophic sets, Elixir International Journal (2014), (Accepted).
- [26] A. A. Salama and Florentin Smarandache, Filters via Neutrosophic Crisp Sets, Neutrosophic Sets and Systems, Vol. (1), ( 2013)pp34-37.
- [27] A. A. Salama, Mohamed Abdelfattah and Mohamed Eisa, A Novel Model for Implementing Security over Mobile Ad-hoc Networks using Intuitionistic Fuzzy Function, International Journal of Emerging Technologies in Computational and Applied Sciences (IJETCAS), Vol.(7), No.(1), (2014)pp01-07.
- [28] A. A. Salama, Mohamed Abdelfattah and S. A. Alblowi, Some Intuitionistic Topological Notions of Intuitionistic Region, Possible Application to GIS Topological Rules, International Journal of Enhanced Research in Management & Computer Applications, Vol.(3), No.(6), (2014)pp1-13.
- [29] M. E. Abd El-Monsef, A.A. Nasef, A. A. Salama, Extensions of fuzzy ideals, Bull. Calcutta Math. Soc. 92, No.3, 181-188 (2000).
- [30] M.E. Abd El-Monsef, A.A. Nasef, and A.A. Salama, Some fuzzy topological operators via fuzzy ideals, Chaos Solitons Fractals, Vol.(12), No.13, (2001)pp 2509-2515
- [31] M. E. Abd El-Monsef, A. A. Nasef and A. A. Salama, Fuzzy L-open sets and fuzzy L-continuous functions"Analele Universitatii de Vest din Timisoara Seria Matematica-Informatica, Vol.( 40), No.2, (2002)pp3-13
- [32] I. M. Hanafy and A.A. Salama, A unified framework including types of fuzzy compactness, Conference Topology and Analysis in Applications Durban, 12-16 July, School of Mathematical Sciences, UKZN, ( 2004)
- [33] A.A. Salama, Fuzzy Hausdorff spaces and fuzzy irresolute functions via fuzzy ideals, V Italian-Spanish Conference on General Topology and its Applications June 21-23, Almeria, Spain, (2004)
- [34] M.E. Abdel Monsef, A. Kozae, A. A. Salama and H. Elagamy, Fuzzy Ideals and Bigranule Computing, 20<sup>th</sup> Conference of Topology and its Applications Port Said, Univ., Egypt, (2007).
- [35] A.A. Salama, Intuitionistic Fuzzy Ideals Theory and Intuitionistic Fuzzy Local Functions, CTAC'08 the 14<sup>th</sup> Biennial Computational Techniques and Applications Conference 13–16 July Australian National University, Canberra, ACT, Australia(2008).
- [36] A.A. Salama, Fuzzy Bitopological Spaces Via Fuzzy Ideals, Blast, August 6-10, University of Denver, Denver, CO, USA, (2008).
- [37] A.A. Salama, A New Form of Fuzzy Compact spaces and Related Topics via Fuzzy Idealization, Journal of fuzzy System and Mathematics Vol.(24), No.2, (2010) pp33-39.
- [38] A.A. Salama and A. Hassan, On Fuzzy Regression Model, the Egyptian Journal for commercial Studies, Vol.(34), No.4.(2010) pp305-319.
- [39] A.A. Salama and S.A. Alblowi, Neutrosophic Set Theory and Neutrosophic Topological Ideal Spaces, The First International Conference on Mathematics and Statistics (ICMS'10) to be held at the American University, (2010).
- [40] A. A. Salama, A New Form of Fuzzy Hausdorff Space and Related Topics via Fuzzy Idealization, IOSR Journal of Mathematics (IOSR-JM), Vol.(3), Issue 5 (2012)pp 01-04 .
- [41] M.E. Abd El-Monsef, A. M. Kozae, A.A. Salama and H. Elagamy, Fuzzy Bitopological Ideals Theory, IOSR Journal of Computer Engineering (IOSRJCE), Vol.(6) ,Issue 4, (2012) pp 01-05.
- [42] I.M. Hanafy, A.A. Salama , M. Abdelfattah and Y. Wazery, " Security in Mant Based on Pki using Fuzzy Function" IOSR Journal of Computer Engineering, Vol.(6), Issue 3,(2012) pp 54-60.
- [43] M. E. Abd El-Monsef, A.Kozae, A.A. Salama, and H. M. Elagamy, Fuzzy Pairwise L-Open Sets and Fuzzy Pairwise L-Continuous Functions, International Journal of Theoretical and Mathematical Physics, Vol.3, No.2, March 2013, pp 69-72.
- [44] I.M. Hanafy, A.A. Salama, M. Abdelfattah and Y. M. Wazery, AIS MODEL FOR BOTNET DETECTION IN MANET USING FUZZY FUNCTION, International Journal of Computer Networking, Wireless and Mobile Communications (IJCNWMC), Vol. 3, Issue 1, Mar (2013)pp95-102 .
- [45] I. M. Hanafy, A. A. Salama, O. M. Khaled and K. M. Mahfouz Correlation of Neutrosophic Sets in Probability Spaces, JAMSI, Vol.10, No.(1), (2014) pp45-52.
- [46] L.A. Zadeh, Fuzzy Sets. In form and control 8 (1965) 338 – 353.

Received: July 29, 2014. Accepted: August 19, 2014.