

# A true and falsifiable statement $\Psi$ with the predicate $\mathcal{K}$ of the current mathematical knowledge, where $\Psi$ strengthens a theorem and does not express what is currently unproved in mathematics

Apoloniusz Tyszk

## Abstract

The theorem of Royer and Case states that there exists a limit-computable function  $\beta_1 : \mathbb{N} \rightarrow \mathbb{N}$  which eventually dominates every computable function  $\delta_1 : \mathbb{N} \rightarrow \mathbb{N}$ . We present an alternative proof of this theorem.  $\mathcal{K}$  denotes both the knowledge predicate satisfied by every currently known theorem and the set of all currently known theorems. The set  $\mathcal{K}$  is time-dependent and publicly available. Any theorem of any mathematician from past or present forever belongs to  $\mathcal{K}$ . We prove: (1) there exists a limit-computable function  $f : \mathbb{N} \rightarrow \mathbb{N}$  of unknown computability which eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation, (2) statement (1) strengthens a theorem. We present both constructive and non-constructive proof of (1). Statement (1) claims that there exists a function  $f : \mathbb{N} \rightarrow \mathbb{N}$  such that ( $f$  is computable in the limit)  $\wedge$  ( $\neg\mathcal{K}(f$  is computable))  $\wedge$  ( $\neg\mathcal{K}(f$  is uncomputable))  $\wedge$  ( $f$  eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation). Since Martin Davis' conjecture on single-fold Diophantine representations disproves statement (1), statement (1) has all properties from the title of the article.

**Key words and phrases:** eventual domination, limit-computable function, predicate  $\mathcal{K}$  of the current mathematical knowledge, single-fold Diophantine representation, time-dependent truth in mathematics with the predicate  $\mathcal{K}$ .

## 1 The goal of the article

$\mathcal{K}$  denotes both the knowledge predicate satisfied by every currently known theorem and the set of all currently known theorems. The set  $\mathcal{K}$  is time-dependent and publicly available. Any theorem of any mathematician from past or present forever belongs to  $\mathcal{K}$ . We formulate statements of the form:

there exists a mathematical object  $\mathcal{X}$  such that

$\mathcal{X}$  satisfies a mathematical condition  $\mathcal{C}$  and

it is unknown whether or not  $\mathcal{X}$  satisfies a mathematical condition  $\mathcal{D}$ .

We present a statement  $\Psi$  of the above form which has all properties from the title of the article.

## 2 Statements with the predicate $\mathcal{K}$ which do not have all properties from the title of the article

Let  $\mathcal{T}$  denote the set of twin primes.

**Proposition 1.** *The statement*

$$(\neg\mathcal{K}(\text{card}(\mathcal{T}) = \omega)) \wedge (\neg\mathcal{K}(\text{card}(\mathcal{T}) < \omega))$$

*is true, falsifiable, and expresses what is currently unproved in mathematics.*

**Statement 1.** *There exists a non-zero integer  $n$  such that*

$$(\neg\mathcal{K}(n < 0)) \wedge (\neg\mathcal{K}(n > 0)) \tag{1}$$

*Proof.* It holds for

$$n = \begin{cases} -1, & \text{if Continuum Hypothesis holds} \\ 1, & \text{otherwise} \end{cases}$$

□

**Proposition 2.** *Statement 1 holds forever.*

*Proof.* Since *Continuum Hypothesis* is independent from *ZFC*, conjunction (1) holds forever for the above  $n$ . □

**Proposition 3.** *Statement 1 does not express what is currently unproved in mathematics.*


### 3 Classical computability theory

For  $n \in \mathbb{N}$ , let

$$E_n = \{1 = x_k, x_i + x_j = x_k, x_i \cdot x_j = x_k : i, j, k \in \{0, \dots, n\}\}$$

**Theorem 1.** ([4, p. 118]). *There exists a limit-computable function  $\beta_1 : \mathbb{N} \rightarrow \mathbb{N}$  which eventually dominates every computable function  $\delta_1 : \mathbb{N} \rightarrow \mathbb{N}$ .*

We present an alternative proof of Theorem 1. For every  $n \in \mathbb{N}$ , we define  $\beta_1(n)$  as the smallest  $b \in \mathbb{N}$  such that if a system of equations  $S \subseteq E_n$  has a solution in  $\mathbb{N}^{n+1}$ , then this solution belongs to  $\{0, \dots, b\}^{n+1}$ . The function  $\beta_1 : \mathbb{N} \rightarrow \mathbb{N}$  is computable in the limit and eventually dominates every computable function  $\delta_1 : \mathbb{N} \rightarrow \mathbb{N}$ , see [5]. Flowchart 1 describes a semi-algorithm which computes  $\beta_1(n)$  in the limit.



**Flowchart 1**

A semi-algorithm which computes  $\beta_1(n)$  in the limit

**Conjecture 1.** ([1, pp. 341–342], [2, p. 42], [3, p. 745]). Every listable set  $\mathcal{X} \subseteq \mathbb{N}^k$  ( $k \in \mathbb{N} \setminus \{0\}$ ) has a single-fold Diophantine representation.

Let  $\Phi$  denote the following statement: *the function  $\mathbb{N} \ni n \rightarrow 2^n \in \mathbb{N}$  eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation.* For  $n \in \mathbb{N}$ , let

$$g(n) = \begin{cases} 2^n, & \text{if } \Phi \text{ holds} \\ \beta_1(n), & \text{otherwise} \end{cases}$$

The function  $g : \mathbb{N} \rightarrow \mathbb{N}$  is computable if and only if  $\Phi$  holds. Currently,

$$(\neg\mathcal{K}(\Phi)) \wedge (\neg\mathcal{K}(\neg\Phi)) \wedge (\neg\mathcal{K}(g \text{ is computable})) \wedge (\neg\mathcal{K}(g \text{ is uncomputable}))$$


**Lemma 1.** *The function  $g$  is computable in the limit and eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation.*

*Proof.* It follows from Theorem 1. □

For every  $n \in \mathbb{N}$ , we define  $\beta(n)$  as the smallest  $b \in \mathbb{N}$  such that if a system of equations  $S \subseteq E_n$  has a unique solution in  $\mathbb{N}^{n+1}$ , then this solution belongs to  $\{0, \dots, b\}^{n+1}$ .

**Theorem 2.** *The function  $\beta : \mathbb{N} \rightarrow \mathbb{N}$  is computable in the limit and eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation.*

*Proof.* This is proved in [5]. The term "dominated" in the title of [5] means "eventually dominated". Flowchart 2 describes a semi-algorithm which computes  $\beta(n)$  in the limit.



**Flowchart 2**

A semi-algorithm which computes  $\beta(n)$  in the limit □

#### 4 A statement described by the title of the article

**Statement 2.** *There exists a limit-computable function  $f : \mathbb{N} \rightarrow \mathbb{N}$  of unknown computability which eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation.*

*Proof.* Statement 2 follows constructively from Theorem 2 by taking  $f = \beta$  and the following conjunction:

$$(\neg\mathcal{K}(\beta \text{ is computable})) \wedge (\neg\mathcal{K}(\beta \text{ is uncomputable}))$$

Statement 2 follows non-constructively from Lemma 1 by taking  $f = g$  and the following conjunction:

$$(\neg\mathcal{K}(g \text{ is computable})) \wedge (\neg\mathcal{K}(g \text{ is uncomputable}))$$

□

**Proposition 4.** *Statement 2 has all properties from the title of the article.*

*Proof.* Statement 2 claims that there exists a function  $f : \mathbb{N} \rightarrow \mathbb{N}$  such that

$$(f \text{ is computable in the limit}) \wedge (\neg\mathcal{K}(f \text{ is computable})) \wedge (\neg\mathcal{K}(f \text{ is uncomputable})) \wedge$$

( $f$  eventually dominates every function  $\delta : \mathbb{N} \rightarrow \mathbb{N}$  with a single-fold Diophantine representation)

Conjecture 1 disproves Statement 2. Statement 2 without the epistemic condition is a theorem. □

Since the function  $\beta_1$  in Theorem 1 is not computable, Statement 2 does not follow from Theorem 1. Ignoring the epistemic condition in Statement 2, Statement 2 follows from Theorem 1 by taking  $f = \beta_1$ .

## References

- [1] M. Davis, Yu. Matiyasevich, J. Robinson, *Hilbert's tenth problem, Diophantine equations: positive aspects of a negative solution*; in: Mathematical developments arising from Hilbert problems (ed. F. E. Browder), Proc. Sympos. Pure Math., vol. 28, Part 2, Amer. Math. Soc., Providence, RI, 1976, 323–378, <http://doi.org/10.1090/pspum/028.2>; reprinted in: The collected works of Julia Robinson (ed. S. Feferman), Amer. Math. Soc., Providence, RI, 1996, 269–324.
- [2] Yu. Matiyasevich, *Hilbert's tenth problem: what was done and what is to be done*, in: Proceedings of the Workshop on Hilbert's tenth problem: relations with arithmetic and algebraic geometry (Ghent, 1999), Contemp. Math. 270, Amer. Math. Soc., Providence, RI, 2000, 1–47, <http://doi.org/10.1090/conm/270>.
- [3] Yu. Matiyasevich, *Towards finite-fold Diophantine representations*, J. Math. Sci. (N. Y.) vol. 171, no. 6, 2010, 745–752, <http://doi.org/10.1007%2Fs10958-010-0179-4>.
- [4] J. S. Royer and J. Case, *Subrecursive Programming Systems: Complexity and Succinctness*, Birkhäuser, Boston, 1994.
- [5] A. Tyszka, *All functions  $g : \mathbb{N} \rightarrow \mathbb{N}$  which have a single-fold Diophantine representation are dominated by a limit-computable function  $f : \mathbb{N} \setminus \{0\} \rightarrow \mathbb{N}$  which is implemented in MuPAD and whose computability is an open problem*, in: Computation, cryptography, and network security (eds. N. J. Daras, M. Th. Rassias), Springer, Cham, 2015, 577–590, [http://doi.org/10.1007/978-3-319-18275-9\\_24](http://doi.org/10.1007/978-3-319-18275-9_24).

Apoloniusz Tyszka  
Technical Faculty  
Hugo Kołłątaj University  
Balicka 116B, 30-149 Kraków, Poland  
E-mail: [rtyszka@cyf-kr.edu.pl](mailto:rtyszka@cyf-kr.edu.pl)