

On true and falsifiable statements Ψ with the predicate \mathcal{K} of the current mathematical knowledge, where Ψ does not express what is proved or unproved in mathematics without \mathcal{K}

Apoloniusz Tyszka

ABSTRACT. The theorem of Royer and Case states that there exists a limit-computable function $\beta_1 : \mathbb{N} \rightarrow \mathbb{N}$ which eventually dominates every computable function $\delta_1 : \mathbb{N} \rightarrow \mathbb{N}$. We present an alternative proof of this theorem. \mathcal{K} denotes both the knowledge predicate satisfied by every currently known theorem and the finite set of all currently known theorems. The set \mathcal{K} is time-dependent and publicly available. Any theorem of any mathematician from past or present forever belongs to \mathcal{K} . We prove: (1) there exists a limit-computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ of unknown computability which eventually dominates every function $\delta : \mathbb{N} \rightarrow \mathbb{N}$ with a single-fold Diophantine representation. We present both constructive and non-constructive proof of (1). Statement (1) claims that there exists a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that $(f \text{ is computable in the limit}) \wedge (\neg \mathcal{K}(f \text{ is computable})) \wedge (\neg \mathcal{K}(f \text{ is uncomputable})) \wedge (f \text{ eventually dominates every function } \delta : \mathbb{N} \rightarrow \mathbb{N} \text{ with a single-fold Diophantine representation})$. Since Martin Davis' conjecture on single-fold Diophantine representations disproves Statement (1), Statement (1) justifies the title of the article. Let Λ denote the statement: *there exists a positive integer b such that it is unknown whether or not the equation $x! + b = y^2$ has infinitely many solutions in positive integers x and y* . We prove that Λ justifies the title of the article.

Key words and phrases: Diophantine equation $x! + b = y^2$, eventual domination, limit-computable function, predicate \mathcal{K} of the current mathematical knowledge, single-fold Diophantine representation, time-dependent truth in mathematics with the predicate \mathcal{K} .

2020 Mathematics Subject Classification: 03F65.

\mathcal{K} denotes both the knowledge predicate satisfied by every currently known theorem and the finite set of all currently known theorems. The set \mathcal{K} is time-dependent and publicly available. Any theorem of any mathematician from past or present forever belongs to \mathcal{K} .

1. STATEMENTS WITH THE PREDICATE \mathcal{K} WHICH DO NOT JUSTIFY THE TITLE OF THE ARTICLE

Let \mathcal{T} denote the set of twin primes.

Proposition 1. *The true statement*

$$(\neg \mathcal{K}(\text{card}(\mathcal{T}) = \omega)) \wedge (\neg \mathcal{K}(\text{card}(\mathcal{T}) < \omega))$$

does not justify the title of the article.

Statement 1. *There exists a non-zero integer n such that*

$$(\neg\mathcal{K}(n < 0)) \wedge (\neg\mathcal{K}(n > 0))$$

Proof. Let

$$n = \begin{cases} -1, & \text{if Continuum Hypothesis holds} \\ 1, & \text{otherwise} \end{cases}$$

□

Proposition 2. *Statement 1 does not justify the title of the article.*

Proof. Since *Continuum Hypothesis* is independent from *ZFC*, Statement 1 holds forever. □


2. CLASSICAL COMPUTABILITY THEORY

For $n \in \mathbb{N}$, let

$$E_n = \{1 = x_k, x_i + x_j = x_k, x_i \cdot x_j = x_k : i, j, k \in \{0, \dots, n\}\}$$

Theorem 1. ([5, p. 118]). *There exists a limit-computable function $\beta_1 : \mathbb{N} \rightarrow \mathbb{N}$ which eventually dominates every computable function $\delta_1 : \mathbb{N} \rightarrow \mathbb{N}$.*

We present an alternative proof of Theorem 1. For every $n \in \mathbb{N}$, we define $\beta_1(n)$ as the smallest $b \in \mathbb{N}$ such that if a system of equations $S \subseteq E_n$ has a solution in \mathbb{N}^{n+1} , then this solution belongs to $\{0, \dots, b\}^{n+1}$. The function $\beta_1 : \mathbb{N} \rightarrow \mathbb{N}$ is computable in the limit and eventually dominates every computable function $\delta_1 : \mathbb{N} \rightarrow \mathbb{N}$, see [6]. Flowchart 1 describes a semi-algorithm which computes $\beta_1(n)$ in the limit.



Flowchart 1

A semi-algorithm which computes $\beta_1(n)$ in the limit

Conjecture 1. ([1, pp. 341–342], [3, p. 42], [4, p. 745]). *Every listable set $\mathcal{X} \subseteq \mathbb{N}^k$ ($k \in \mathbb{N} \setminus \{0\}$) has a single-fold Diophantine representation.*

Let Φ denote the following statement: *the function $\mathbb{N} \ni n \rightarrow 2^n \in \mathbb{N}$ eventually dominates every function $\delta : \mathbb{N} \rightarrow \mathbb{N}$ with a single-fold Diophantine representation.* For $n \in \mathbb{N}$, let

$$g(n) = \begin{cases} 2^n, & \text{if } \Phi \text{ holds} \\ \beta_1(n), & \text{otherwise} \end{cases}$$

The function $g : \mathbb{N} \rightarrow \mathbb{N}$ is computable if and only if Φ holds. Currently,

$$(\neg\mathcal{K}(\Phi)) \wedge (\neg\mathcal{K}(\neg\Phi)) \wedge (\neg\mathcal{K}(g \text{ is computable})) \wedge (\neg\mathcal{K}(g \text{ is uncomputable}))$$


Lemma 1. *The function g is computable in the limit and eventually dominates every function $\delta : \mathbb{N} \rightarrow \mathbb{N}$ with a single-fold Diophantine representation.*

Proof. It follows from Theorem 1. □

For every $n \in \mathbb{N}$, we define $\beta(n)$ as the smallest $b \in \mathbb{N}$ such that if a system of equations $S \subseteq E_n$ has a unique solution in \mathbb{N}^{n+1} , then this solution belongs to $\{0, \dots, b\}^{n+1}$.

Theorem 2. *The function $\beta : \mathbb{N} \rightarrow \mathbb{N}$ is computable in the limit and eventually dominates every function $\delta : \mathbb{N} \rightarrow \mathbb{N}$ with a single-fold Diophantine representation.*

Proof. This is proved in [6]. The term "dominated" in the title of [6] means "eventually dominated". Flowchart 2 describes a semi-algorithm which computes $\beta(n)$ in the limit.



Flowchart 2

A semi-algorithm which computes $\beta(n)$ in the limit

□

3. STATEMENTS WITH THE PREDICATE \mathcal{K} WHICH JUSTIFY THE TITLE OF THE ARTICLE

Statement 2. *There exists a limit-computable function $f : \mathbb{N} \rightarrow \mathbb{N}$ of unknown computability which eventually dominates every function $\delta : \mathbb{N} \rightarrow \mathbb{N}$ with a single-fold Diophantine representation.*

Proof. Statement 2 follows constructively from Theorem 2 by taking $f = \beta$ and the following conjunction:

$$(\neg\mathcal{K}(\beta \text{ is computable})) \wedge (\neg\mathcal{K}(\beta \text{ is uncomputable}))$$

Statement 2 follows non-constructively from Lemma 1 by taking $f = g$ and the following conjunction:

$$(\neg\mathcal{K}(g \text{ is computable})) \wedge (\neg\mathcal{K}(g \text{ is uncomputable}))$$

□

Proposition 3. *Statement 2 justifies the title of the article.*

Proof. Statement 2 claims that there exists a function $f : \mathbb{N} \rightarrow \mathbb{N}$ such that

$$(f \text{ is computable in the limit}) \wedge (\neg\mathcal{K}(f \text{ is computable})) \wedge$$

$$(\neg\mathcal{K}(f \text{ is uncomputable})) \wedge (f \text{ eventually dominates every function } \delta : \mathbb{N} \rightarrow \mathbb{N} \text{ with a single-fold Diophantine representation})$$

Conjecture 1 disproves Statement 2. □

Since the function β_1 in Theorem 1 is not computable, Statement 2 does not follow from Theorem 1. Ignoring the epistemic condition in Statement 2, Statement 2 follows from Theorem 1 by taking $f = \beta_1$.

Statement 3. *There exists a positive integer b such that it is unknown whether or not the equation $x! + b = y^2$ has infinitely many solutions in positive integers x and y .*

Proof. It holds for $b = 1$, see [2]. □

Proposition 4. *Statement 3 justifies the title of the article.*

Proof. Statement 3 claims that there exists a positive integer b such that

$$\begin{aligned} &\neg\mathcal{K}(\text{the equation } x! + b = y^2 \text{ has infinitely} \\ &\text{many solutions in positive integers } x \text{ and } y) \wedge \\ &\neg\mathcal{K}(\text{the equation } x! + b = y^2 \text{ has at most finitely} \\ &\text{many solutions in positive integers } x \text{ and } y) \end{aligned}$$

A weak form of Szpiro's conjecture implies that the equation $x! + b = y^2$ has at most finitely many solutions in positive integers x and y when $b \in \mathbb{N} \setminus \{0\}$, see [2]. □

In [8], the author showed that the predicate \mathcal{K} non-trivially extends constructive mathematics, see also [7], [9], [10].

REFERENCES

- [1] M. Davis, Yu. Matiyasevich, J. Robinson, *Hilbert's tenth problem, Diophantine equations: positive aspects of a negative solution*; in: Mathematical developments arising from Hilbert problems (ed. F. E. Browder), Proc. Sympos. Pure Math., vol. 28, Part 2, Amer. Math. Soc., Providence, RI, 1976, 323–378, <http://doi.org/10.1090/pspum/028.2>; reprinted in: The collected works of Julia Robinson (ed. S. Feferman), Amer. Math. Soc., Providence, RI, 1996, 269–324.
- [2] A. Dąbrowski, *On the Diophantine equation $x! + A = y^2$* , Nieuw Arch. Wisk. IV. Ser. 14 (1996), no. 3, 321–324.
- [3] Yu. Matiyasevich, *Hilbert's tenth problem: what was done and what is to be done*, in: Proceedings of the Workshop on Hilbert's tenth problem: relations with arithmetic and algebraic geometry (Ghent, 1999), Contemp. Math. 270, Amer. Math. Soc., Providence, RI, 2000, 1–47, <http://doi.org/10.1090/conm/270>.
- [4] Yu. Matiyasevich, *Towards finite-fold Diophantine representations*, J. Math. Sci. (N. Y.) vol. 171, no. 6, 2010, 745–752, <http://doi.org/10.1007%2Fs10958-010-0179-4>.
- [5] J. S. Royer and J. Case, *Subrecursive Programming Systems: Complexity and Succinctness*, Birkhäuser, Boston, 1994.
- [6] A. Tyszka, *All functions $g : \mathbb{N} \rightarrow \mathbb{N}$ which have a single-fold Diophantine representation are dominated by a limit-computable function $f : \mathbb{N} \setminus \{0\} \rightarrow \mathbb{N}$ which is implemented in MuPAD and whose computability is an open problem*, in: Computation, cryptography, and network security (eds. N. J. Daras, M. Th. Rassias), Springer, Cham, 2015, 577–590, http://doi.org/10.1007/978-3-319-18275-9_24.
- [7] A. Tyszka, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that refer to the current knowledge on \mathcal{X}* , Journal of Applied Computer Science & Mathematics 16 (2022), no. 2, 31–35, <http://doi.org/10.4316/JACSM.202202005>.
- [8] A. Tyszka, *In constructive and informal mathematics, in contradistinction to any empirical science, the predicate of the current knowledge in the subject is necessary*, Asian Research Journal of Mathematics 19 (2023), no. 12, 69–79, <http://doi.org/10.9734/arjom/2023/v19i12773>.
- [9] A. Tyszka, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$* , Pi Mu Epsilon J. 15 (2023), no. 8, 493–504.
- [10] A. Tyszka, *Statements and open problems on decidable sets $\mathcal{X} \subseteq \mathbb{N}$ that contain informal notions and refer to the current knowledge on \mathcal{X}* , Creat. Math. Inform. 32 (2023), no. 2, 247–253, http://semnul.com/creative-mathematics/wp-content/uploads/2023/07/creative_2023_32_2_247_253.pdf.

Apoloniusz Tyszka
Technical Faculty
Hugo Kołłątaj University
Balicka 116B, 30-149 Kraków, Poland
E-mail: rtyszka@cyf-kr.edu.pl