

# Theorems and open problems that concern decidable sets $\mathcal{X} \subseteq \mathbb{N}$ and cannot be formulated in the formal language of classical mathematics as they refer to the current knowledge on $\mathcal{X}$

AGNIESZKA KOZDĘBA, APOLONIUSZ TYSZKA

**ABSTRACT.** Edmund Landau's conjecture states that the set  $\mathcal{P}_{n^2+1}$  of primes of the form  $n^2 + 1$  is infinite. Landau's conjecture implies the following unproven statement  $\Phi$ :  $\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, (((24!)!)!)]$ . We heuristically justify the statement  $\Phi$ . This justification does not yield the finiteness/infiniteness of  $\mathcal{P}_{n^2+1}$ . We present a new heuristic argument for the infiniteness of  $\mathcal{P}_{n^2+1}$ , which is not based on the statement  $\Phi$ . The distinction between algorithms whose existence is provable in *ZFC* and constructively defined algorithms which are currently known inspires theorems and open problems that concern decidable sets  $\mathcal{X} \subseteq \mathbb{N}$  and cannot be formulated in the formal language of classical mathematics as they refer to the current knowledge on  $\mathcal{X}$ .

**2020 Mathematics Subject Classification:** 03F55, 11A41.

**Key words and phrases:** algorithms whose existence is provable in *ZFC*, conjecturally infinite sets  $\mathcal{X} \subseteq \mathbb{N}$ , constructively defined integer  $n$  satisfies  $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$ , current knowledge on a set  $\mathcal{X} \subseteq \mathbb{N}$ , known elements of a set  $\mathcal{X} \subseteq \mathbb{N}$  whose infiniteness is false or unproven, mathematical definitions, theorems and open problems with epistemic and informal notions, physical limits of computation, primes of the form  $n^2 + 1$ ,  $\mathcal{X}$  is decidable by a constructively defined algorithm which is currently known.

## 1. INTRODUCTION

Sections 2–4 contain purely mathematical results, which we summarize now shortly starting from the results of sections 2 and 3. Edmund Landau's conjecture states that the set  $\mathcal{P}_{n^2+1}$  of primes of the form  $n^2 + 1$  is infinite, see [14]–[16]. Landau's conjecture implies the following unproven statement  $\Phi$ :  $\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, (((24!)!)!)]$ . Let  $f(1) = 2$ ,  $f(2) = 4$ , and let  $f(n + 1) = f(n)!$  for every integer  $n \geq 2$ . Let  $B$  denote the system of equations:

$$\{x_i! = x_k : i, k \in \{1, \dots, 9\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, 9\}\}$$

We write some system  $\mathcal{U} \subseteq B$  of 9 equations which has exactly two solutions in positive integers  $x_1, \dots, x_9$ , namely  $(1, \dots, 1)$  and  $(f(1), \dots, f(9))$ . No known system  $\mathcal{S} \subseteq B$  with a finite number of solutions in positive integers  $x_1, \dots, x_9$  has

a solution  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  satisfying  $\max(x_1, \dots, x_9) > f(9)$ . For every known system  $\mathcal{S} \subseteq B$ , if the finiteness/infiniteness of the set

$$\{(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9 : (x_1, \dots, x_9) \text{ solves } \mathcal{S}\}$$

is unknown, then the statement

$$\exists x_1, \dots, x_9 \in \mathbb{N} \setminus \{0\} ((x_1, \dots, x_9) \text{ solves } \mathcal{S}) \wedge (\max(x_1, \dots, x_9) > f(9))$$

remains unproven.

We write some system  $\mathcal{A} \subseteq B$  of 8 equations. Let  $\Lambda$  denote the statement: *if the system  $\mathcal{A}$  has at most finitely many solutions in positive integers  $x_1, \dots, x_9$ , then each such solution  $(x_1, \dots, x_9)$  satisfies  $x_1, \dots, x_9 \leq f(9)$* . The statement  $\Lambda$  is equivalent to the statement  $\Phi$ . It heuristically justifies the statement  $\Phi$ . This justification does not yield the finiteness/infiniteness of  $\mathcal{P}_{n^2+1}$ .

In section 4, we present a new heuristic argument for the infiniteness of  $\mathcal{P}_{n^2+1}$ , which is not based on the statement  $\Phi$ .

Theorems and open problems in sections 5–8 involve epistemic and informal notions and justify the next sentence. The distinction between algorithms whose existence is provable in *ZFC* and constructively defined algorithms which are currently known inspires theorems and open problems that concern decidable sets  $\mathcal{X} \subseteq \mathbb{N}$  and cannot be formulated in the formal language of classical mathematics as they refer to the current knowledge on  $\mathcal{X}$ .

## 2. NUMBER-THEORETIC STATEMENTS $\Psi_n$

Let  $f(1) = 2$ ,  $f(2) = 4$ , and let  $f(n+1) = f(n)!$  for every integer  $n \geq 2$ . Let  $\mathcal{U}_1$  denote the system of equations which consists of the equation  $x_1! = x_1$ . For an integer  $n \geq 2$ , let  $\mathcal{U}_n$  denote the following system of equations:

$$\begin{cases} x_1! = x_1 \\ x_1 \cdot x_1 = x_2 \\ \forall i \in \{2, \dots, n-1\} x_i! = x_{i+1} \end{cases}$$

**Lemma 1.** *For every positive integer  $n$ , the system  $\mathcal{U}_n$  has exactly two solutions in positive integers  $x_1, \dots, x_n$ , namely  $(1, \dots, 1)$  and  $(f(1), \dots, f(n))$ .*

Let  $B_n$  denote the following system of equations:

$$\{x_i! = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$$

For every positive integer  $n$ , no known system  $\mathcal{S} \subseteq B_n$  with a finite number of solutions in positive integers  $x_1, \dots, x_n$  has a solution  $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$  satisfying  $\max(x_1, \dots, x_n) > f(n)$ . For every positive integer  $n$  and for every known system  $\mathcal{S} \subseteq B_n$ , if the finiteness/infiniteness of the set

$$\{(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n : (x_1, \dots, x_n) \text{ solves } \mathcal{S}\}$$

is unknown, then the statement

$$\exists x_1, \dots, x_n \in \mathbb{N} \setminus \{0\} ((x_1, \dots, x_n) \text{ solves } \mathcal{S}) \wedge (\max(x_1, \dots, x_n) > f(n))$$

remains unproven.

For a positive integer  $n$ , let  $\Psi_n$  denote the following statement: *if a system  $S \subseteq B_n$  has at most finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq f(n)$* . The statement  $\Psi_n$  says that for subsystems of  $B_n$  with a finite number of solutions, the largest known solution is indeed the largest possible. The statements  $\Psi_1$  and  $\Psi_2$  hold trivially. There is no reason to assume the validity of the statement  $\forall n \in \mathbb{N} \setminus \{0\} \Psi_n$ .

**Theorem 1.** *For every statement  $\Psi_n$ , the bound  $f(n)$  cannot be decreased.*

*Proof.* It follows from Lemma 1 because  $\mathcal{U}_n \subseteq B_n$ . □

**Theorem 2.** *For every integer  $n \geq 2$ , the statement  $\Psi_{n+1}$  implies the statement  $\Psi_n$ .*

*Proof.* If a system  $S \subseteq B_n$  has at most finitely many solutions in positive integers  $x_1, \dots, x_n$ , then for every integer  $i \in \{1, \dots, n\}$  the system  $S \cup \{x_i! = x_{n+1}\}$  has at most finitely many solutions in positive integers  $x_1, \dots, x_{n+1}$ . The statement  $\Psi_{n+1}$  implies that  $x_i! = x_{n+1} \leq f(n+1) = f(n)!$ . Hence,  $x_i \leq f(n)$ . □

**Theorem 3.** *Every statement  $\Psi_n$  is true with an unknown integer bound that depends on  $n$ .*

*Proof.* For every positive integer  $n$ , the system  $B_n$  has a finite number of subsystems. □

3. A SPECIAL CASE OF THE STATEMENT  $\Psi_9$  APPLIES TO EDMUND LANDAU'S CONJECTURE THAT THE SET  $\mathcal{P}_{n^2+1}$  OF PRIMES OF THE FORM  $n^2 + 1$  IS INFINITE


**Lemma 2.** *For every positive integers  $x$  and  $y$ ,  $x! \cdot y = y!$  if and only if*

$$(x + 1 = y) \vee (x = y = 1)$$

Let  $\mathcal{A}$  denote the following system of equations:

$$\left\{ \begin{array}{l} x_2! = x_3 \\ x_3! = x_4 \\ x_5! = x_6 \\ x_8! = x_9 \\ x_1 \cdot x_1 = x_2 \\ x_3 \cdot x_5 = x_6 \\ x_4 \cdot x_8 = x_9 \\ x_5 \cdot x_7 = x_8 \end{array} \right.$$

Lemma 2 and the diagram in Figure 1 explain the construction of the system  $\mathcal{A}$ .



**Fig. 1** Construction of the system  $\mathcal{A}$

**Lemma 3.** (Wilson's theorem, [4, p. 89]). For every integer  $x \geq 2$ ,  $x$  is prime if and only if  $x$  divides  $(x-1)! + 1$ .

**Lemma 4.** For every integer  $x_1 \geq 2$ , the system  $\mathcal{A}$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1^2 + 1$  is prime. In this case, the integers  $x_2, \dots, x_9$  are uniquely determined by the following equalities:

$$\begin{aligned}
 x_2 &= x_1^2 \\
 x_3 &= (x_1^2)! \\
 x_4 &= ((x_1^2)!)! \\
 x_5 &= x_1^2 + 1 \\
 x_6 &= (x_1^2 + 1)! \\
 x_7 &= \frac{(x_1^2)! + 1}{x_1^2 + 1} \\
 x_8 &= (x_1^2)! + 1 \\
 x_9 &= ((x_1^2)! + 1)!
 \end{aligned}$$

*Proof.* By Lemma 2, for every integer  $x_1 \geq 2$ , the system  $\mathcal{A}$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1^2 + 1$  divides  $(x_1^2)! + 1$ . Hence, the claim of Lemma 4 follows from Lemma 3.  $\square$

**Lemma 5.** There are only finitely many tuples  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$ , which solve the system  $\mathcal{A}$  and satisfy  $x_1 = 1$ . It is true as every such tuple  $(x_1, \dots, x_9)$  satisfies  $x_1, \dots, x_9 \in \{1, 2\}$ .

*Proof.* The equality  $x_1 = 1$  implies that  $x_2 = x_1 \cdot x_1 = 1$ . Hence,  $x_3 = x_2! = 1$ . Therefore,  $x_4 = x_3! = 1$ . The equalities  $x_5! = x_6$  and  $x_5 = 1 \cdot x_5 = x_3 \cdot x_5 = x_6$  imply that  $x_5, x_6 \in \{1, 2\}$ . The equalities  $x_8! = x_9$  and  $x_8 = 1 \cdot x_8 = x_4 \cdot x_8 = x_9$  imply that  $x_8, x_9 \in \{1, 2\}$ . The equality  $x_5 \cdot x_7 = x_8$  implies that  $x_7 = \frac{x_8}{x_5} \in \left\{\frac{1}{1}, \frac{1}{2}, \frac{2}{1}, \frac{2}{2}\right\} \cap (\mathbb{N} \setminus \{0\}) = \{1, 2\}$ .  $\square$

**Conjecture 1.** *The statement  $\Psi_9$  is true when is restricted to the system  $\mathcal{A}$ .*

Edmund Landau's conjecture states that the set  $\mathcal{P}_{n^2+1}$  of primes of the form  $n^2 + 1$  is infinite, see [14]–[16].

**Theorem 4.** *Conjecture 1 proves the following implication: if there exists an integer  $x_1 \geq 2$  such that  $x_1^2 + 1$  is prime and greater than  $f(7)$ , then the set  $\mathcal{P}_{n^2+1}$  is infinite.*

*Proof.* Suppose that the antecedent holds. By Lemma 4, there exists a unique tuple  $(x_2, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^8$  such that the tuple  $(x_1, x_2, \dots, x_9)$  solves the system  $\mathcal{A}$ . Since  $x_1^2 + 1 > f(7)$ , we obtain that  $x_1^2 \geq f(7)$ . Hence,  $(x_1^2)! \geq f(7)! = f(8)$ . Consequently,

$$x_9 = ((x_1^2)! + 1)! \geq (f(8) + 1)! > f(8)! = f(9)$$

Conjecture 1 and the inequality  $x_9 > f(9)$  imply that the system  $\mathcal{A}$  has infinitely many solutions  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$ . According to Lemmas 4 and 5, the set  $\mathcal{P}_{n^2+1}$  is infinite.  $\square$

Landau's conjecture implies the following unproven statement  $\Phi$ :

$$\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, (((24!)!)!)]$$

Theorem 5 heuristically justifies the statement  $\Phi$ . This justification does not yield the finiteness/infiniteness of  $\mathcal{P}_{n^2+1}$ .

**Theorem 5.** *Conjecture 1 implies the statement  $\Phi$ .*

*Proof.* It follows from Theorem 4 and the equality  $f(7) = (((24!)!)!)!$ .  $\square$

**Theorem 6.** *The statement  $\Phi$  implies Conjecture 1.*

*Proof.* By Lemmas 4 and 5, if positive integers  $x_1, \dots, x_9$  solve the system  $\mathcal{A}$ , then

$$(x_1 \geq 2) \wedge (x_5 = x_1^2 + 1) \wedge (x_5 \text{ is prime})$$

or  $x_1, \dots, x_9 \in \{1, 2\}$ . In the first case, Lemma 4 and the statement  $\Phi$  imply that the inequality  $x_5 \leq (((24!)!)!)! = f(7)$  holds when the system  $\mathcal{A}$  has at most finitely many solutions in positive integers  $x_1, \dots, x_9$ . Hence,  $x_2 = x_5 - 1 < f(7)$  and  $x_3 = x_2! < f(7)! = f(8)$ . Continuing this reasoning in the same manner, we can show that every  $x_i$  does not exceed  $f(9)$ .  $\square$

#### 4. A NEW HEURISTIC ARGUMENT FOR THE INFINITENESS OF $\mathcal{P}_{n^2+1}$

The system  $\mathcal{A}$  contains four factorials and four multiplications. Let  $\mathcal{F}$  denote the family of all systems  $\mathcal{S} \subseteq \mathcal{B}_9$  which contain at most four factorials and at most four multiplications.

Among known systems  $\mathcal{S} \in \mathcal{F}$ , the following system  $C$

$$\left\{ \begin{array}{l} x_1! = x_2 \\ x_2 \cdot x_9 = x_1 \\ x_2 \cdot x_2 = x_3 \\ x_3 \cdot x_3 = x_4 \\ x_4 \cdot x_4 = x_5 \\ x_5! = x_6 \\ x_6! = x_7 \\ x_7! = x_8 \end{array} \right.$$

attains the greatest solution in positive integers  $x_1, \dots, x_9$  and has at most finitely many solutions in  $(\mathbb{N} \setminus \{0\})^9$ . Only the tuples  $(1, \dots, 1)$  and  $(2, 2, 4, 16, 256, 256!, (256!)!, ((256!)!)!, 1)$  solve  $C$  and belong to  $(\mathbb{N} \setminus \{0\})^9$ .

For every known system  $\mathcal{S} \in \mathcal{F}$ , if the finiteness of the set

$$\{(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9 : (x_1, \dots, x_9) \text{ solves } \mathcal{S}\}$$

is unproven and conjectured, then the statement

$$\exists x_1, \dots, x_9 \in \mathbb{N} \setminus \{0\} ((x_1, \dots, x_9) \text{ solves } \mathcal{S}) \wedge (\max(x_1, \dots, x_9) > ((256!)!)!$$

remains unproven.

Let  $\Gamma$  denote the statement: *if the system  $\mathcal{A}$  has at most finitely many solutions in positive integers  $x_1, \dots, x_9$ , then each such solution  $(x_1, \dots, x_9)$  satisfies  $x_1, \dots, x_9 \leq ((256!)!)!$ . The number  $46^{512} + 1$  is prime ([9]) and greater than  $256!$ , see also [12, p. 239] for the primality of  $150^{2048} + 1$ . Hence, the statement  $\Gamma$  is equivalent to the infiniteness of  $\mathcal{P}_{n^2+1}$ . It heuristically justifies the infiniteness of  $\mathcal{P}_{n^2+1}$  in a sophisticated way.*

## 5. THE DISTINCTION BETWEEN ALGORITHMS WHOSE EXISTENCE IS PROVABLE IN ZFC AND CONSTRUCTIVELY DEFINED ALGORITHMS WHICH ARE CURRENTLY KNOWN

Algorithms always terminate. Semi-algorithms may not terminate. Examples 1–4 and the proof of Statement 1 explain the distinction between *existing algorithms* (i.e. algorithms whose existence is provable in ZFC) and *known algorithms* (i.e. algorithms whose definition is constructive and currently known). A definition of an integer  $n$  is called *constructive*, if it provides a known algorithm with no input that returns  $n$ . Definition 1 applies to sets  $\mathcal{X} \subseteq \mathbb{N}$  whose infiniteness is false or unproven.

**Definition 1.** *We say that a non-negative integer  $k$  is a known element of  $\mathcal{X}$ , if  $k \in \mathcal{X}$  and we know an algebraic expression that defines  $k$  and consists of the following signs: 1 (one), + (addition), - (subtraction),  $\cdot$  (multiplication),  $\wedge$  (exponentiation with exponent in  $\mathbb{N}$ ), ! (factorial of a non-negative integer), ( (left parenthesis), ) (right parenthesis).*

The set of known elements of  $\mathcal{X}$  is finite and time-dependent, so cannot be defined in the formal language of classical mathematics. Let  $t$  denote the largest twin prime that is smaller than  $(((((9!)!)!)!)!)!$ . The number  $t$  is an unknown element of the set of twin primes.

**Definition 2.** *Conditions (1)–(5) concern sets  $\mathcal{X} \subseteq \mathbb{N}$ .*

(1) *A known algorithm with no input returns an integer  $n$  satisfying  $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$ .*

(2) *A known algorithm for every  $k \in \mathbb{N}$  decides whether or not  $k \in \mathcal{X}$ .*

(3) *No known algorithm with no input returns the logical value of the statement  $\text{card}(\mathcal{X}) = \omega$ .*

(4) *There are many elements of  $\mathcal{X}$  and it is conjectured, though so far unproven, that  $\mathcal{X}$  is infinite.*

(5)  *$\mathcal{X}$  is naturally defined. The infiniteness of  $\mathcal{X}$  is false or unproven.  $\mathcal{X}$  has the simplest definition among known sets  $\mathcal{Y} \subseteq \mathbb{N}$  with the same set of known elements.*

Condition (3) implies that no known proof shows the finiteness/infiniteness of  $\mathcal{X}$ . No known set  $\mathcal{X} \subseteq \mathbb{N}$  satisfies Conditions (1)–(4) and is widely known in number theory or naturally defined, where this term has only informal meaning.

Let  $[\cdot]$  denote the integer part function. Let  $\beta = (((24!)!)!)!$ .

**Lemma 6.**  $\log_2(\log_2(\log_2(\log_2(\log_2(\log_2(\log_2(\beta))))))) \approx 1.42298$ .

*Proof.* We ask Wolfram Alpha at <http://wolframalpha.com>. □

**Example 1.** *The set  $\mathcal{X} = \mathcal{P}_{n^2+1}$  satisfies Condition (3).*

**Example 2.** *The set  $\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } [\frac{\beta}{\pi}] \text{ is odd} \\ \emptyset, & \text{otherwise} \end{cases}$  does not satisfy Condition (3) because we know an algorithm with no input that computes  $[\frac{\beta}{\pi}]$ . The set of known elements of  $\mathcal{X}$  is empty. Hence, Condition (5) fails for  $\mathcal{X}$ .*

**Example 3.** ([1], [11], [13, p. 9]). *The function*

$$\mathbb{N} \ni n \xrightarrow{h} \begin{cases} 1, & \text{if the decimal expansion of } \pi \text{ contains } n \text{ consecutive zeros} \\ 0, & \text{otherwise} \end{cases}$$

*is computable because  $h = \mathbb{N} \times \{1\}$  or there exists  $k \in \mathbb{N}$  such that*

$$h = (\{0, \dots, k\} \times \{1\}) \cup (\{k+1, k+2, k+3, \dots\} \times \{0\})$$

*No known algorithm computes the function  $h$ .*

**Example 4.** *The set*

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if the continuum hypothesis holds} \\ \emptyset, & \text{otherwise} \end{cases}$$

*is decidable. This  $\mathcal{X}$  satisfies Conditions (1) and (3) and does not satisfy Conditions (2), (4), and (5). These facts will hold forever.*

**Statement 1.** *Condition (1) remains unproven for  $\mathcal{X} = \mathcal{P}_{n^2+1}$ .*

*Proof.* For every set  $\mathcal{X} \subseteq \mathbb{N}$ , there exists an algorithm  $\text{Alg}(\mathcal{X})$  with no input that returns

$$n = \begin{cases} 0, & \text{if } \text{card}(\mathcal{X}) \in \{0, \omega\} \\ \max(\mathcal{X}), & \text{otherwise} \end{cases}$$

This  $n$  satisfies the implication in Condition (1), but the algorithm  $\text{Alg}(\mathcal{P}_{n^2+1})$  is unknown because its definition is ineffective.  $\square$

Proving the statement  $\Phi$  will disprove Statement 1. Statement 1 cannot be formulated in the formal language of classical mathematics because it refers to the current mathematical knowledge. The same is true for Open Problems 1–5 and Statements 2–5.

**Definition 3.** We say that an integer  $n$  is a threshold number of a set  $\mathcal{X} \subseteq \mathbb{N}$ , if  $\text{card}(\mathcal{X}) < \omega \Rightarrow \mathcal{X} \subseteq (-\infty, n]$ .

If a set  $\mathcal{X} \subseteq \mathbb{N}$  is empty or infinite, then any integer  $n$  is a threshold number of  $\mathcal{X}$ . If a set  $\mathcal{X} \subseteq \mathbb{N}$  is non-empty and finite, then the all threshold numbers of  $\mathcal{X}$  form the set  $[\max(\mathcal{X}), \infty) \cap \mathbb{N}$ .

## 6. THE PHYSICAL LIMITS OF COMPUTATION INSPIRE OPEN PROBLEM 1

**Statement 2.** *The set*

$$\mathcal{X} = \{k \in \mathbb{N} : (10^6 < k) \Rightarrow (f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$$

*satisfies Conditions (1)–(4). Condition (5) fails for  $\mathcal{X}$ .*

*Proof.* Condition (4) holds as  $\mathcal{X} \supseteq \{0, \dots, 10^6\}$  and the set  $\mathcal{P}_{n^2+1}$  is conjecturally infinite. By Lemma 6, due to known physics we are not able to confirm by a direct computation that some element of  $\mathcal{P}_{n^2+1}$  is greater than  $f(10^6) > f(7) = \beta$ , see [7]. Thus Condition (3) holds. Condition (2) holds trivially. Since the set

$$\{k \in \mathbb{N} : (10^6 < k) \wedge (f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$$

is empty or infinite, the integer  $10^6$  is a threshold number of  $\mathcal{X}$ . Thus  $\mathcal{X}$  satisfies Condition (1). Condition (5) fails for  $\mathcal{X}$  as the set of known elements of  $\mathcal{X}$  equals  $\{0, \dots, 10^6\}$ .  $\square$

Statement 3 provides a stronger example. To formulate Statement 3 and its proof, we need some lemmas.

For a non-negative integer  $n$ , let  $\theta(n)$  denote the largest integer divisor of  $10^{10^{10}}$  smaller than  $n$ . For a non-negative integer  $n$ , let  $\theta_1(n)$  denote the largest integer divisor of  $10^{10}$  smaller than  $n$ .

**Lemma 7.** *For every integer  $j > 10^{10^{10}}$ ,  $\theta(j) = 10^{10^{10}}$ .*

**Lemma 8.** *For every integer  $j \in (6553600, 7812500]$ ,  $\theta(j) = 6553600$ .*



*Proof.* 6553600 equals  $2^{18} \cdot 5^2$  and divides  $10^{10^{10}}$ .  $7812500 < 2^{24}$ .  $7812500 < 5^{10}$ . We need to prove that every integer  $j \in (6553600, 7812500)$  does not divide  $10^{10^{10}}$ . It holds as the set

$$\{2^u \cdot 5^v : (u \in \{0, \dots, 23\}) \wedge (v \in \{0, \dots, 9\})\}$$

contains 6553600 and 7812500 as consecutive elements.  $\square$

**Lemma 9.** *The number  $6553600^2 + 1$  is prime.*

*Proof.* The following PARI/GP ([8]) command

`isprime(6553600^2+1, {flag=2})`

returns 1. This command performs the APRCL primality test, the best deterministic primality test algorithm ([17, p. 226]). It rigorously shows that the number  $6553600^2 + 1$  is prime.  $\square$

In the next lemmas, the execution of the command `isprime(n, {flag=2})` proves the primality of  $n$ .

**Lemma 10.** *The number  $10142101504^2 + 1$  is prime.  $10142101504 > 10^{10}$ .*

**Lemma 11.** *The function*

$$\mathbb{N} \ni n \xrightarrow{\kappa} \underbrace{\text{the\_exponent\_of\_2\_in\_the\_prime\_factorization\_of\_}n+1}_{\kappa} \in \mathbb{N}$$

*takes every non-negative integer value infinitely often.*

Before Open Problem 1,  $\mathcal{X}$  denotes the set  $\{n \in \mathbb{N} : (\theta(n) + \kappa(n))^2 + 1 \text{ is prime}\}$ .

**Lemma 12.** *The set  $\mathcal{X}$  satisfies  $\text{card}(\mathcal{X}) \geq 629450$ .*

*Proof.* By Lemmas 8 and 9, for every even integer  $j \in (6553600, 7812500]$ , the number  $(\theta(j) + \kappa(j))^2 + 1 = (6553600 + 0)^2 + 1$  is prime. Hence,

$$\{2k : k \in \mathbb{N}\} \cap (6553600, 7812500] \subseteq \mathcal{X}$$

Consequently,

$$\begin{aligned} \text{card}(\mathcal{X}) &\geq \text{card}(\{2k : k \in \mathbb{N}\} \cap (6553600, 7812500]) = \\ &= \frac{7812500 - 6553600}{2} = 629450 \end{aligned}$$

$\square$

**Lemma 13.**  $10242 \in \mathcal{X}$ .  $10242 \notin \mathcal{X}_1 = \{n \in \mathbb{N} : (\theta_1(n) + \kappa(n))^2 + 1 \text{ is prime}\}$ .

*Proof.* The number  $10240 = 2^{11} \cdot 5$  divides  $10^{10^{10}}$ . Hence,  $\theta(10242) = 10240$ . The number  $(\theta(10242) + \kappa(10242))^2 + 1 = (10240 + 0)^2 + 1$  is prime. The set

$$\{2^u \cdot 5^v : (u \in \{0, \dots, 10\}) \wedge (v \in \{0, \dots, 10\})\}$$

contains 10000 and 12500 as consecutive elements. Hence,  $\theta_1(10242) = 10000$ . The number  $(\theta_1(10242) + \kappa(10242))^2 + 1 = (10000 + 0)^2 + 1 = 17 \cdot 5882353$  is composite.  $\square$

**Statement 3.** *The set  $X$  satisfies Conditions (1)–(5) except the requirement that  $X$  is naturally defined.*

*Proof.* Condition (2) holds trivially. Let  $\delta$  denote  $10^{10^{10}}$ . By Lemmas 7 and 11, Condition (1) holds for  $n = \delta$ . Since the statement  $\mathcal{P}_{n^2+1} \cap (\delta^2 + 1, \infty) \neq \emptyset$  remains unproven, Condition (3) holds. Lemma 12 and the implication

$$\mathcal{P}_{n^2+1} \cap (\delta^2 + 1, \infty) \neq \emptyset \implies \text{card}(X) = \omega$$

show that Condition (4) holds. By Lemma 10, the set  $X_1$  is infinite. Since Definition 1 applies to sets  $X \subseteq \mathbb{N}$  whose infiniteness is false or unproven, Condition (5) holds except the requirement that  $X$  is naturally defined.  $\square$

The set  $X$  satisfies Condition (5) except the requirement that  $X$  is naturally defined. It is true because  $X_1$  is infinite and Definition 1 applies only to sets  $X \subseteq \mathbb{N}$  whose infiniteness is false or unproven. Ignoring this restriction,  $X$  still satisfies the same identical condition due to Lemma 13.


**Open Problem 1.** *Is there a set  $X \subseteq \mathbb{N}$  which satisfies Conditions (1)–(5)?*

The answers to Open Problems 1–5 may change in time as they depend on the current mathematical knowledge. These answers are currently negative.

**Theorem 7.** *No set  $X \subseteq \mathbb{N}$  will satisfy Conditions (1)–(4) forever, if for every algorithm with no input, at some future day, a computer will be able to execute this algorithm in 1 second or less.*

*Proof.* The proof goes by contradiction. We fix an integer  $n$  that satisfies Condition (1). Since Conditions (1)–(3) will hold forever, the semi-algorithm in Figure 2 never terminates and sequentially prints the following sentences:

(T)  $n + 1 \notin X, n + 2 \notin X, n + 3 \notin X, \dots$



**Fig. 2** Semi-algorithm that terminates if and only if  $X$  is infinite

The sentences from the sequence (T) and our assumption imply that for every integer  $m > n$  computed by a known algorithm, at some future day, a computer will be able to confirm in 1 second or less that  $(n, m] \cap \mathcal{X} = \emptyset$ . Thus, at some future day, numerical evidence will support the conjecture that the set  $\mathcal{X}$  is finite, contrary to the conjecture in Condition (4).  $\square$

The physical limits of computation ([7]) disprove the assumption of Theorem 7.

**Statement 4.** *Conditions (2)–(5) hold for  $\mathcal{X} = \mathcal{P}_{n^2+1}$ . The statement  $\Phi$  implies that Condition (1) holds for  $\mathcal{X} = \mathcal{P}_{n^2+1}$ .*

*Proof.* The set  $\mathcal{P}_{n^2+1}$  is conjecturally infinite. There are 2199894223892 primes of the form  $n^2 + 1$  in the interval  $[2, 10^{28})$ , see [15]. These two facts imply Condition (4). By Lemma 6, due to known physics we are not able to confirm by a direct computation that some element of  $\mathcal{P}_{n^2+1}$  is greater than  $f(7) = (((24!)!)!) = \beta$ , see [7]. Thus Condition (3) holds. Conditions (2) and (5) hold trivially. The statement  $\Phi$  implies that Condition (1) holds for  $\mathcal{X} = \mathcal{P}_{n^2+1}$  with  $n = (((24!)!)!)!$ .  $\square$

Proving Landau's conjecture will disprove Statement 4.

**Conjecture 2.** *(Conditions (1)–(5) hold for  $\mathcal{X} = \mathcal{P}_{n^2+1}$ )  $\wedge$   $\Phi$ .*

Conjecture 2 implies that every known proof of the statement  $\Phi$  does not yield the finiteness/infiniteness of  $\mathcal{P}_{n^2+1}$ .

#### 7. SATISFIABLE CONJUNCTIONS WHICH CONSIST OF CONDITIONS (1)–(5) AND THEIR NEGATIONS

The set  $\mathcal{X} = \mathcal{P}_{n^2+1}$  satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})$$

The set  $\mathcal{X} = \{0, \dots, 10^6\} \cup \mathcal{P}_{n^2+1}$  satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge (\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

The set  $\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } (f(9^8), f(9^9)) \cap \mathcal{P}_{n^2+1} \neq \emptyset \\ \{0, \dots, 10^6\}, & \text{otherwise} \end{cases}$  satisfies the conjunction

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

**Open Problem 2.** *Is there a set  $\mathcal{X} \subseteq \mathbb{N}$  that satisfies the conjunction*

$$(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})?$$

The numbers  $2^{2^k} + 1$  are prime for  $k \in \{0, 1, 2, 3, 4\}$ . It is open whether or not there are infinitely many primes of the form  $2^{2^k} + 1$ , see [6, p. 158] and [12, p. 74]. It is open whether or not there are infinitely many composite numbers of the form  $2^{2^k} + 1$ , see [6, p. 159] and [12, p. 74]. Most mathematicians believe that  $2^{2^k} + 1$  is composite for every integer  $k \geq 5$ , see [5, p. 23].

The set

$$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } (f(9^8), f(9^9)) \cap \mathcal{P}_{n^2+1} \neq \emptyset \\ \{0, \dots, 10^6\} \cup \\ \{n \in \mathbb{N} : n \text{ is the sixth prime number of the form } 2^{2^k} + 1\}, & \text{otherwise} \end{cases}$$

satisfies the conjunction

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \neg(\text{Condition 5})$$

**Open Problem 3.** *Is there a set  $\mathcal{X} \subseteq \mathbb{N}$  that satisfies the conjunction*

$$\neg(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \neg(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge (\text{Condition 5})?$$

It is possible, although very doubtful, that at some future day, the set  $\mathcal{X} = \mathcal{P}_{n^2+1}$  will solve Open Problem 2. The same is true for Open Problem 3. It is possible, although very doubtful, that at some future day, the set  $\mathcal{X} = \{k \in \mathbb{N} : 2^{2^k} + 1 \text{ is composite}\}$  will solve Open Problem 1. The same is true for Open Problems 2 and 3.

The following table shows satisfiable conjunctions of the form

$$\#(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \#(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \#(\text{Condition 5})$$

where # denotes the negation  $\neg$  or the absence of any symbol.

	(Condition 2) $\wedge$ (Condition 3) $\wedge$ (Condition 4)	(Condition 2) $\wedge$ $\neg$ (Condition 3) $\wedge$ (Condition 4)
(Condition 1) $\wedge$ (Condition 5)	Open Problem 1 (conjecturally solved with $\mathcal{X} = \mathcal{P}_{n^2+1}$ )	Open Problem 2
(Condition 1) $\wedge$ $\neg$ (Condition 5)	$\mathcal{X} = \{k \in \mathbb{N} : (10^6 < k) \Rightarrow$ $(f(10^6), f(k)) \cap \mathcal{P}_{n^2+1} \neq \emptyset\}$	$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } (f(9^8), f(9^9)) \cap \mathcal{P}_{n^2+1} \neq \emptyset \\ \{0, \dots, 10^6\}, & \text{otherwise} \end{cases}$
$\neg$ (Condition 1) $\wedge$ (Condition 5)	$\mathcal{X} = \mathcal{P}_{n^2+1}$	Open Problem 3
$\neg$ (Condition 1) $\wedge$ $\neg$ (Condition 5)	$\mathcal{X} = \{0, \dots, 10^6\} \cup \mathcal{P}_{n^2+1}$	$\mathcal{X} = \begin{cases} \mathbb{N}, & \text{if } (f(9^8), f(9^9)) \cap \mathcal{P}_{n^2+1} \neq \emptyset \\ \{0, \dots, 10^6\} \cup \{n \in \mathbb{N} : n \text{ is} \\ \text{the sixth prime number of} \\ \text{the form } 2^{2^k} + 1\}, & \text{otherwise} \end{cases}$

## 8. PREVIOUSLY KNOWN RESULTS WHICH CORRESPOND TO THE RESULTS OF SECTIONS 5–7

Statements 1–4 and Open Problems 1–3 cannot be formulated in the formal language of classical mathematics. Previously known statements of this type, such as Statement 5, express the current knowledge on particular elements of  $\mathbb{N}$ , which are known to us according to Definition 1. Previously known open problems of this type, such as Open Problems 4 and 5, ask about constructive existence of special elements of  $\mathbb{N}$ .

**Statement 5.** ([2], [3], [6, p. 209], [10]). *The numbers  $2^{2^{22}} + 1$  and  $2^{2^{24}} + 1$  are composite. The known integer divisors of  $2^{2^{22}} + 1$  form the set  $\{-2^{2^{22}} - 1, -1, 1, 2^{2^{22}} + 1\}$ . The known integer divisors of  $2^{2^{24}} + 1$  form the set  $\{-2^{2^{24}} - 1, -1, 1, 2^{2^{24}} + 1\}$ .*

**Open Problem 4.** *Is there a known threshold number of  $\mathcal{P}_{n^2+1}$ ?*

**Open Problem 5.** *Is there a known threshold number of the set of twin primes?*

## 9. SUMMARY

Let  $f(1) = 2$ ,  $f(2) = 4$ , and let  $f(n + 1) = f(n)!$  for every integer  $n \geq 2$ . Edmund Landau's conjecture states that the set  $\mathcal{P}_{n^2+1}$  of primes of the form  $n^2 + 1$  is infinite. Landau's conjecture implies the following unproven statement  $\Phi$ :  $\text{card}(\mathcal{P}_{n^2+1}) < \omega \Rightarrow \mathcal{P}_{n^2+1} \subseteq [2, (((24!)!)!)]$ . Let  $B$  denote the system of equations:  $\{x_i! = x_k : i, k \in \{1, \dots, 9\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, 9\}\}$ . We write some system  $\mathcal{U} \subseteq B$  of 9 equations which has exactly two solutions in positive integers  $x_1, \dots, x_9$ , namely  $(1, \dots, 1)$  and  $(f(1), \dots, f(9))$ . No known system  $\mathcal{S} \subseteq B$  with a finite number of solutions in positive integers  $x_1, \dots, x_9$  has a solution  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  satisfying  $\max(x_1, \dots, x_9) > f(9)$ . For every known system  $\mathcal{S} \subseteq B$ , if the finiteness/infiniteness of the set  $\{(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9 : (x_1, \dots, x_9) \text{ solves } \mathcal{S}\}$  is unknown, then the statement  $\exists x_1, \dots, x_9 \in \mathbb{N} \setminus \{0\} ((x_1, \dots, x_9) \text{ solves } \mathcal{S}) \wedge (\max(x_1, \dots, x_9) > f(9))$  remains unproven. We write some system  $\mathcal{A} \subseteq B$  of 8 equations. Let  $\Lambda$  denote the statement: *if the system  $\mathcal{A}$  has at most finitely many solutions in positive integers  $x_1, \dots, x_9$ , then each such solution  $(x_1, \dots, x_9)$  satisfies  $x_1, \dots, x_9 \leq f(9)$ .* The statement  $\Lambda$  is equivalent to the statement  $\Phi$ . It heuristically justifies the statement  $\Phi$ . This justification does not yield the finiteness/infiniteness of  $\mathcal{P}_{n^2+1}$ . We present a new heuristic argument for the infiniteness of  $\mathcal{P}_{n^2+1}$ , which is not based on the statement  $\Phi$ . Algorithms always terminate. The next theorems and open problems justify the title of the article and involve epistemic and informal notions. We explain the distinction between existing algorithms (i.e. algorithms whose existence is provable in ZFC) and known algorithms (i.e. algorithms whose definition is constructive and currently known). Assuming that the infiniteness of a set  $X \subseteq \mathbb{N}$  is false or unproven, we define which elements of  $X$  are classified as known. No known set  $X \subseteq \mathbb{N}$  satisfies Conditions (1)–(4) and is widely known in number theory or naturally defined, where this term has only informal meaning. (1) *A known algorithm with no input returns an integer  $n$  satisfying  $\text{card}(X) < \omega \Rightarrow X \subseteq (-\infty, n]$ .* (2) *A known algorithm for every  $k \in \mathbb{N}$  decides whether or not  $k \in X$ .* (3) *No known algorithm with no input returns the logical value of the statement  $\text{card}(X) = \omega$ .* (4) *There are many elements of  $X$  and it is conjectured, though so far unproven, that  $X$  is infinite.*

(5)  $\mathcal{X}$  is naturally defined. The infiniteness of  $\mathcal{X}$  is false or unproven.  $\mathcal{X}$  has the simplest definition among known sets  $\mathcal{Y} \subseteq \mathbb{N}$  with the same set of known elements. Conditions (2)–(5) hold for  $\mathcal{X} = \mathcal{P}_{n^2+1}$ . The statement  $\Phi$  implies the conjunction of Conditions (1)–(5) for  $\mathcal{X} = \mathcal{P}_{n^2+1}$ . We define a set  $\mathcal{X} \subseteq \mathbb{N}$  which satisfies Conditions (1)–(5) except the requirement that  $\mathcal{X}$  is naturally defined. We present a table that shows satisfiable conjunctions of the form  $\#(\text{Condition 1}) \wedge (\text{Condition 2}) \wedge \#(\text{Condition 3}) \wedge (\text{Condition 4}) \wedge \#(\text{Condition 5})$ , where  $\#$  denotes the negation  $\neg$  or the absence of any symbol. No set  $\mathcal{X} \subseteq \mathbb{N}$  will satisfy Conditions (1)–(4) forever, if for every algorithm with no input, at some future day, a computer will be able to execute this algorithm in 1 second or less. The physical limits of computation disprove this assumption.

**Acknowledgement.** An earlier version of this article was presented at Logic Colloquium 2021, <http://lc2021.pl>. Agnieszka Kozdęba prepared two diagrams. Apoloniusz Tyszką wrote the article.

#### REFERENCES

- [1] J. Case, M. Ralston, *Beyond Rogers' non-constructively computable function*, in: *The nature of computation*, Lecture Notes in Comput. Sci., 7921, 45–54, Springer, Heidelberg, 2013, [http://link.springer.com/chapter/10.1007/978-3-642-39053-1\\_6](http://link.springer.com/chapter/10.1007/978-3-642-39053-1_6).
- [2] R. Crandall, J. Doenias, C. Norrie, J. Young, *The twenty-second Fermat number is composite*, Math. Comp. 64 (1995), 863–868.
- [3] R. Crandall, E. Mayer, J. Papadopoulos, *The twenty-fourth Fermat number is composite*, Math. Comp. 72 (2003), 1555–1572.
- [4] M. Erickson, A. Vazzana, D. Garth, *Introduction to number theory*, 2nd ed., CRC Press, Boca Raton, FL, 2016.
- [5] J.-M. De Koninck, F. Luca, *Analytic number theory: Exploring the anatomy of integers*, American Mathematical Society, Providence, RI, 2012.
- [6] M. Křížek, F. Luca, L. Somer, *17 lectures on Fermat numbers: from number theory to geometry*, Springer, New York, 2001.
- [7] S. Lloyd, *Ultimate physical limits to computation*, Nature 406 (2000), 1047–1054, <http://doi.org/10.1038/35023282>.
- [8] PARI/GP online documentation, [http://pari.math.u-bordeaux.fr/dochtml/html/Arithmetic\\_functions.html](http://pari.math.u-bordeaux.fr/dochtml/html/Arithmetic_functions.html).
- [9] X. M. Pi, *Searching for generalized Fermat primes (Chinese)*, J. Math. (Wuhan) 18 (1998), no. 3, 276–280.
- [10] *Proth Search Page*, <http://www.prothsearch.com/fermat.html#Complete>.
- [11] R. Reitzig, *How can it be decidable whether  $\pi$  has some sequence of digits?*, <http://cs.stackexchange.com/questions/367/how-can-it-be-decidable-whether-pi-has-some-sequence-of-digits>.
- [12] P. Ribenboim, *The little book of bigger primes*, 2nd ed., Springer, New York, 2004.

- 
- [13] H. Rogers, Jr., *Theory of recursive functions and effective computability*, 2nd ed., MIT Press, Cambridge, MA, 1987.
- [14] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, A002496, *Primes of the form  $n^2 + 1$* , <http://oeis.org/A002496>.
- [15] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, A083844, *Number of primes of the form  $x^2 + 1 < 10^n$* , <http://oeis.org/A083844>.
- [16] Wolfram MathWorld, *Landau's Problems*, <http://mathworld.wolfram.com/LandausProblems.html>.
- [17] S. Y. Yan, *Number theory for computing*, 2nd ed., Springer, Berlin, 2002.

Agnieszka Kozdęba  
Faculty of Environmental Engineering and Land Surveying  
Hugo Kołłątaj University  
Balicka 253C, 30-198 Kraków, Poland  
&  
Institute of Mathematics  
Jagiellonian University  
Łojasiewicza 6, 30-348 Kraków, Poland  
E-mail address: [agnieszka.kozdeba@gmail.com](mailto:agnieszka.kozdeba@gmail.com)

Apoloniusz Tyszka  
Technical Faculty  
Hugo Kołłątaj University  
Balicka 116B, 30-149 Kraków, Poland  
E-mail address: [rttyszka@cyf-kr.edu.pl](mailto:rttyszka@cyf-kr.edu.pl)