

# On *ZFC*-formulae $\varphi(x)$ for which we know a non-negative integer $n$ such that $\max(\{x \in \mathbb{N} : \varphi(x)\}) \leq n$ if the set $\{x \in \mathbb{N} : \varphi(x)\}$ is finite

Apoloniusz Tyszk

## Abstract

Let  $\Gamma(k)$  denote  $(k-1)!$ , and let  $\Gamma_n(k)$  denote  $(k-1)!$ , where  $n \in \{3, \dots, 16\}$  and  $k \in \{2\} \cup [2^{2^{n-3}} + 1, \infty) \cap \mathbb{N}$ . For an integer  $n \in \{3, \dots, 16\}$ , let  $\Sigma_n$  denote the following statement: if a system of equations  $\mathcal{S} \subseteq \{\Gamma_n(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$  with  $\Gamma$  instead of  $\Gamma_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then every tuple  $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$  that solves the original system  $\mathcal{S}$  satisfies  $x_1, \dots, x_n \leq 2^{2^{n-2}}$ . Our hypothesis claims that the statements  $\Sigma_3, \dots, \Sigma_{16}$  are true. The statement  $\Sigma_6$  proves the following implication: if the equation  $x(x+1) = y!$  has only finitely many solutions in positive integers  $x$  and  $y$ , then each such solution  $(x, y)$  belongs to the set  $\{(1, 2), (2, 3)\}$ . The statement  $\Sigma_6$  proves the following implication: if the equation  $x! + 1 = y^2$  has only finitely many solutions in positive integers  $x$  and  $y$ , then each such solution  $(x, y)$  belongs to the set  $\{(4, 5), (5, 11), (7, 71)\}$ . The statement  $\Sigma_9$  implies the infinitude of primes of the form  $n^2 + 1$ . The statement  $\Sigma_9$  implies that any prime of the form  $n! + 1$  with  $n \geq 2^{2^{9-3}}$  proves the infinitude of primes of the form  $n! + 1$ . The statement  $\Sigma_{14}$  implies the infinitude of twin primes. The statement  $\Sigma_{16}$  implies the infinitude of Sophie Germain primes.

**Keywords:** Brocard's problem, Brocard-Ramanujan equation  $x! + 1 = y^2$ , composite Fermat numbers, Erdős' equation  $x(x+1) = y!$ , prime numbers of the form  $n^2 + 1$ , prime numbers of the form  $n! + 1$ , Sophie Germain primes, twin primes.

**2010 Mathematics Subject Classification:** 03B30, 11A41.

## 1 Introduction and basic lemmas

The phrase “we know a non-negative integer  $n$ ” in the title means that we know an algorithm which returns  $n$ . The title of the article cannot be formalized in *ZFC* because the phrase “we know a non-negative integer  $n$ ” refers to currently known non-negative integers  $n$  with some property. A formally stated title may look like this: *On ZFC-formulae  $\varphi(x)$  for which there exists a non-negative integer  $n$  such that ZFC proves that*

$$\text{card}(\{x \in \mathbb{N} : \varphi(x)\}) < \infty \implies \max(\{x \in \mathbb{N} : \varphi(x)\}) \leq n$$

Unfortunately, this formulation admits formulae  $\varphi(x)$  without any known non-negative integer  $n$  such that *ZFC* proves the above implication.

**Lemma 1.** *For every positive integers  $x$  and  $y$ ,  $x! \cdot y = y!$  if and only if*

$$(x + 1 = y) \vee (x = y = 1)$$

Let  $\Gamma(k)$  denote  $(k-1)!$ .

**Lemma 2.** *For every positive integers  $x$  and  $y$ ,  $x \cdot \Gamma(x) = \Gamma(y)$  if and only if*

$$(x + 1 = y) \vee (x = y = 1)$$

**Lemma 3.** *For every non-negative integers  $b$  and  $c$ ,  $b + 1 = c$  if and only if  $2^{2^b} \cdot 2^{2^b} = 2^{2^c}$ .*

**Lemma 4.** *(Wilson's theorem, [7, p. 89]). For every positive integer  $x$ ,  $x$  divides  $(x-1)! + 1$  if and only if  $x = 1$  or  $x$  is prime.*

## 2 Subsets of $\mathbb{N}$ and their threshold numbers

We say that a non-negative integer  $m$  is a threshold number of a set  $X \subseteq \mathbb{N}$ , if  $X$  is infinite if and only if  $X$  contains an element greater than  $m$ , cf. [22] and [23]. If a set  $X \subseteq \mathbb{N}$  is empty or infinite, then any non-negative integer  $m$  is a threshold number of  $X$ . If a set  $X \subseteq \mathbb{N}$  is non-empty and finite, then the all threshold numbers of  $X$  form the set  $\{\max(X), \max(X) + 1, \max(X) + 2, \dots\}$ .

It is conjectured that the set of prime numbers of the form  $n^2 + 1$  is infinite, see [12, pp. 37–38]. It is conjectured that the set of prime numbers of the form  $n! + 1$  is infinite, see [3, p. 443]. A twin prime is a prime number that differs from another prime number by 2. The twin prime conjecture states that the set of twin primes is infinite, see [12, p. 39]. It is conjectured that the set of composite numbers of the form  $2^{2^n} + 1$  is infinite, see [8, p. 23] and [9, pp. 158–159]. A prime  $p$  is said to be a Sophie Germain prime if both  $p$  and  $2p + 1$  are prime, see [20]. It is conjectured that the set of Sophie Germain primes is infinite, see [15, p. 330]. For each of these sets, we do not know any threshold number.

The following statement:

for every non-negative integer  $n$  there exist

$$\text{prime numbers } p \text{ and } q \text{ such that } p + 2 = q \text{ and } p \in \left[10^n, 10^n + 1\right] \quad (1)$$

is a  $\Pi_1$  statement which strengthens the twin prime conjecture, see [4, p. 43]. C. H. Bennett claims that most mathematical conjectures can be settled indirectly by proving stronger  $\Pi_1$  statements, see [1]. Statement (1) is equivalent to the non-halting of a Turing machine. If a set  $X \subseteq \mathbb{N}$  is computable and we know a threshold number of  $X$ , then the infinity of  $X$  is equivalent to the halting of a Turing machine.

The height of a rational number  $\frac{p}{q}$  is denoted by  $H\left(\frac{p}{q}\right)$  and equals  $\max(|p|, |q|)$  provided  $\frac{p}{q}$  is written in lowest terms. The height of a rational tuple  $(x_1, \dots, x_n)$  is denoted by  $H(x_1, \dots, x_n)$  and equals  $\max(H(x_1), \dots, H(x_n))$ .

**Lemma 5.** *The equation  $x^5 - x = y^2 - y$  has only finitely many rational solutions, see [11, p. 212]. The known rational solutions are  $(x, y) = (-1, 0), (-1, 1), (0, 0), (0, 1), (1, 0), (1, 1), (2, -5), (2, 6), (3, -15), (3, 16), (30, -4929), (30, 4930), \left(\frac{1}{4}, \frac{15}{32}\right), \left(\frac{1}{4}, \frac{17}{32}\right), \left(-\frac{15}{16}, -\frac{185}{1024}\right), \left(-\frac{15}{16}, \frac{1209}{1024}\right)$ , and the existence of other solutions is an open question, see [16, pp. 223–224].*

**Corollary 1.** *The set  $\mathcal{T} = \{n \in \mathbb{N} : \text{the equation } x^5 - x = y^2 - y \text{ has a rational solution of height } n\}$  is finite. We know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{T}$ . We do not know any algorithm which returns a threshold number of  $\mathcal{T}$ .*

Let  $\mathcal{D}$  denote the following system of equations:

$$\begin{cases} x^2 + y^2 = s^2 \\ x^2 + z^2 = t^2 \\ y^2 + z^2 = u^2 \\ x^2 + y^2 + z^2 = v^2 \end{cases}$$

Let  $\mathcal{F} = \{z \in \mathbb{N} : \text{the system } \mathcal{D} \text{ has a solution } (x, y, z, s, t, u, v) \in (\mathbb{N} \setminus \{0\})^7 \text{ with } x \leq y \leq z\}$ . A perfect cuboid is a cuboid having integer side lengths, integer face diagonals, and an integer space diagonal.

**Lemma 6.** ([19]). *No perfect cuboids are known.*

**Corollary 2.** *The set  $\mathcal{F}$  is empty or infinite. We know an algorithm which for every  $z \in \mathbb{N}$  decides whether or not  $z \in \mathcal{F}$ . Every non-negative integer  $z$  is a threshold number of  $\mathcal{F}$ .*

Let

$$\mathcal{H} = \begin{cases} \mathbb{N}, & \text{if } \sin\left(999999\right) < 0 \\ \mathbb{N} \cap \left[0, \sin\left(999999\right) \cdot 999999\right) & \text{otherwise} \end{cases}$$

We do not know whether or not the set  $\mathcal{H}$  is finite.

**Proposition 1.** *The number  $999999$  is a threshold number of  $\mathcal{H}$ . We know an algorithm which decides the equality  $\mathcal{H} = \mathbb{N}$ . If  $\mathcal{H} \neq \mathbb{N}$ , then the set  $\mathcal{H}$  consists of all integers from 0 to a non-negative integer which can be computed by a known algorithm. We know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{H}$ .*

### 3 A Diophantine equation whose non-solvability expresses the consistency of ZFC

Gödel's second incompleteness theorem and the Davis-Putnam-Robinson-Matiyasevich theorem imply the following theorem.

**Theorem 1.** *([5, p. 35]). There exists a polynomial  $D(x_1, \dots, x_m)$  with integer coefficients such that if ZFC is arithmetically consistent, then the sentences "The equation  $D(x_1, \dots, x_m) = 0$  is solvable in non-negative integers" and "The equation  $D(x_1, \dots, x_m) = 0$  is not solvable in non-negative integers" are not provable in ZFC.*

Let  $\mathcal{Y}$  denote the set of all non-negative integers  $k$  such that the equation  $D(x_1, \dots, x_m) = 0$  has no solutions in  $\{0, \dots, k\}^m$ . Since the set  $\{0, \dots, k\}^m$  is finite, we know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{Y}$ . Let  $\mathcal{E}$  denote the set of all non-negative integers  $k$  such that the equation  $D(x_1, \dots, x_m) = 0$  has a solution in  $\{0, \dots, k\}^m$ . Since the set  $\{0, \dots, k\}^m$  is finite, we know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{E}$ . Theorem 1 implies Theorems 2 and 3.

**Theorem 2.** *For every  $n \in \mathbb{N}$ , ZFC proves that  $n \in \mathcal{Y}$ . If ZFC is arithmetically consistent, then the sentences " $\mathcal{Y}$  is finite" and " $\mathcal{Y}$  is infinite" are not provable in ZFC. If ZFC is arithmetically consistent, then for every  $n \in \mathbb{N}$  the sentences " $n$  is a threshold number of  $\mathcal{Y}$ " and " $n$  is not a threshold number of  $\mathcal{Y}$ " are not provable in ZFC.*


**Theorem 3.** *The set  $\mathcal{E}$  is empty or infinite. In both cases, every non-negative integer  $n$  is a threshold number of  $\mathcal{E}$ . If ZFC is arithmetically consistent, then the sentences " $\mathcal{E}$  is empty", " $\mathcal{E}$  is not empty", " $\mathcal{E}$  is finite", and " $\mathcal{E}$  is infinite" are not provable in ZFC.*

### 4 Hypothetical statements $\Psi_3, \dots, \Psi_{16}$

For an integer  $n \geq 3$ , let  $\mathcal{U}_n$  denote the following system of equations:

$$\begin{cases} \forall i \in \{1, \dots, n-1\} \setminus \{2\} x_i! = x_{i+1} \\ x_1 \cdot x_2 = x_3 \\ x_2 \cdot x_2 = x_3 \end{cases}$$

The diagram in Figure 1 illustrates the construction of the system  $\mathcal{U}_n$ .



**Fig. 1** Construction of the system  $\mathcal{U}_n$

Let  $g(3) = 4$ , and let  $g(n + 1) = g(n)!$  for every integer  $n \geq 3$ .

**Lemma 7.** For every integer  $n \geq 3$ , the system  $\mathcal{U}_n$  has exactly two solutions in positive integers, namely  $(1, \dots, 1)$  and  $(2, 2, g(3), \dots, g(n))$ .

Let

$$B_n = \{x_i! = x_k : (i, k \in \{1, \dots, n\}) \wedge (i \neq k)\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$$

For an integer  $n \geq 3$ , let  $\Psi_n$  denote the following statement: if a system of equations  $\mathcal{S} \subseteq B_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq g(n)$ . The statement  $\Psi_n$  says that for subsystems of  $B_n$  the largest known solution is indeed the largest possible.

**Hypothesis 1.** The statements  $\Psi_3, \dots, \Psi_{16}$  are true.

**Proposition 2.** Every statement  $\Psi_n$  is true with an unknown integer bound that depends on  $n$ .

*Proof.* For every positive integer  $n$ , the system  $B_n$  has a finite number of subsystems. □

**Proposition 3.** For every statement  $\Psi_n$ , the bound  $g(n)$  cannot be decreased.


*Proof.* It follows from Lemma 7 because  $\mathcal{U}_n \subseteq B_n$ . □

## 5 The Brocard-Ramanujan equation $x! + 1 = y^2$

Let  $\mathcal{A}$  denote the following system of equations:

$$\begin{cases} x_1! = x_2 \\ x_2! = x_3 \\ x_5! = x_6 \\ x_4 \cdot x_4 = x_5 \\ x_3 \cdot x_5 = x_6 \end{cases}$$

Lemma 1 and the diagram in Figure 2 explain the construction of the system  $\mathcal{A}$ .



**Fig. 2** Construction of the system  $\mathcal{A}$

**Lemma 8.** For every  $x_1, x_4 \in \mathbb{N} \setminus \{0, 1\}$ , the system  $\mathcal{A}$  is solvable in positive integers  $x_2, x_3, x_5, x_6$  if and only if  $x_1! + 1 = x_4^2$ . In this case, the integers  $x_2, x_3, x_5, x_6$  are uniquely determined by the following equalities:

$$\begin{aligned} x_2 &= x_1! \\ x_3 &= (x_1!)! \\ x_5 &= x_1! + 1 \\ x_6 &= (x_1! + 1)! \end{aligned}$$

*Proof.* It follows from Lemma 1. □

It is conjectured that  $x! + 1$  is a perfect square only for  $x \in \{4, 5, 7\}$ , see [18, p. 297]. A weak form of Szpiro's conjecture implies that there are only finitely many solutions to the equation  $x! + 1 = y^2$ , see [13].

**Theorem 4.** If the equation  $x_1! + 1 = x_4^2$  has only finitely many solutions in positive integers, then the statement  $\Psi_6$  guarantees that each such solution  $(x_1, x_4)$  belongs to the set  $\{(4, 5), (5, 11), (7, 71)\}$ .


*Proof.* Suppose that the antecedent holds. Let positive integers  $x_1$  and  $x_4$  satisfy  $x_1! + 1 = x_4^2$ . Then,  $x_1, x_4 \in \mathbb{N} \setminus \{0, 1\}$ . By Lemma 8, the system  $\mathcal{A}$  is solvable in positive integers  $x_2, x_3, x_5, x_6$ . Since  $\mathcal{A} \subseteq B_6$ , the statement  $\Psi_6$  implies that  $x_6 = (x_1! + 1)! \leq g(6) = g(5)!$ . Hence,  $x_1! + 1 \leq g(5) = g(4)!$ . Consequently,  $x_1 < g(4) = 24$ . If  $x_1 \in \{1, \dots, 23\}$ , then  $x_1! + 1$  is a perfect square only for  $x_1 \in \{4, 5, 7\}$ . □

## 6 Are there infinitely many prime numbers of the form $n^2 + 1$ ?

Edmund Landau's conjecture states that there are infinitely many primes of the form  $n^2 + 1$ , see [12, pp. 37–38]. Let  $\mathcal{B}$  denote the following system of equations:

$$\left\{ \begin{array}{l} x_2! = x_3 \\ x_3! = x_4 \\ x_5! = x_6 \\ x_8! = x_9 \\ x_1 \cdot x_1 = x_2 \\ x_3 \cdot x_5 = x_6 \\ x_4 \cdot x_8 = x_9 \\ x_5 \cdot x_7 = x_8 \end{array} \right.$$

Lemma 1 and the diagram in Figure 3 explain the construction of the system  $\mathcal{B}$ .



**Fig. 3** Construction of the system  $\mathcal{B}$

**Lemma 9.** For every integer  $x_1 \geq 2$ , the system  $\mathcal{B}$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1^2 + 1$  is prime. In this case, the integers  $x_2, \dots, x_9$  are uniquely determined by the following equalities:

$$\begin{aligned}
 x_2 &= x_1^2 \\
 x_3 &= (x_1^2)! \\
 x_4 &= ((x_1^2)!) \\
 x_5 &= x_1^2 + 1 \\
 x_6 &= (x_1^2 + 1)! \\
 x_7 &= \frac{(x_1^2)! + 1}{x_1^2 + 1} \\
 x_8 &= (x_1^2)! + 1 \\
 x_9 &= ((x_1^2)! + 1)!
 \end{aligned}$$

*Proof.* By Lemma 1, for every integer  $x_1 \geq 2$ , the system  $\mathcal{B}$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1^2 + 1$  divides  $(x_1^2)! + 1$ . Hence, the claim of Lemma 9 follows from Lemma 4.  $\square$

**Lemma 10.** There are only finitely many tuples  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  which solve the system  $\mathcal{B}$  and satisfy  $x_1 = 1$ .

*Proof.* If a tuple  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  solves the system  $\mathcal{B}$  and  $x_1 = 1$ , then  $x_1, \dots, x_9 \leq 2$ . Indeed,  $x_1 = 1$  implies that  $x_2 = x_1^2 = 1$ . Hence, for example,  $x_3 = x_2! = 1$ . Therefore,  $x_8 = x_3 + 1 = 2$  or  $x_8 = 1$ . Consequently,  $x_9 = x_8! \leq 2$ .  $\square$

**Theorem 5.** The statement  $\Psi_9$  proves the following implication: if there exists an integer  $x_1 \geq 2$  such that  $x_1^2 + 1$  is prime and greater than  $g(7)$ , then there are infinitely many primes of the form  $n^2 + 1$ .

*Proof.* Suppose that the antecedent holds. By Lemma 9, there exists a unique tuple  $(x_2, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^8$  such that the tuple  $(x_1, x_2, \dots, x_9)$  solves the system  $\mathcal{B}$ . Since  $x_1^2 + 1 > g(7)$ , we obtain that  $x_1^2 \geq g(7)$ . Hence,  $(x_1^2)! \geq g(7)! = g(8)$ . Consequently,

$$x_9 = ((x_1^2)! + 1)! \geq (g(8) + 1)! > g(8)! = g(9)$$

Since  $\mathcal{B} \subseteq \mathcal{B}_9$ , the statement  $\Psi_9$  and the inequality  $x_9 > g(9)$  imply that the system  $\mathcal{B}$  has infinitely many solutions  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$ . According to Lemmas 9 and 10, there are infinitely many primes of the form  $n^2 + 1$ .  $\square$

**Corollary 3.** Let  $X_9$  denote the set of primes of the form  $n^2 + 1$ . The statement  $\Psi_9$  implies that we know an algorithm such that it returns a threshold number of  $X_9$ , and this number equals  $\max(X_9)$ , if  $X_9$  is finite. Assuming the statement  $\Psi_9$ , a single query to an oracle for the halting problem decides the infinity of  $X_9$ . Assuming the statement  $\Psi_9$ , the infinity of  $X_9$  is decidable in the limit.

*Proof.* We consider an algorithm which computes  $\max(X_9 \cap [1, g(7)])$ . □

## 7 Are there infinitely many prime numbers of the form $n! + 1$ ?

It is conjectured that there are infinitely many primes of the form  $n! + 1$ , see [3, p. 443].

**Theorem 6.** (cf. Theorem 10). The statement  $\Psi_9$  proves the following implication: if there exists an integer  $x_1 \geq g(6)$  such that  $x_1! + 1$  is prime, then there are infinitely many primes of the form  $n! + 1$ .


*Proof.* We leave the analogous proof to the reader. □

## 8 The twin prime conjecture

A twin prime is a prime number that differs from another prime number by 2. The twin prime conjecture states that there are infinitely many twin primes, see [12, p. 39]. Let  $C$  denote the following system of equations:

$$\left\{ \begin{array}{l} x_1! = x_2 \\ x_2! = x_3 \\ x_4! = x_5 \\ x_6! = x_7 \\ x_7! = x_8 \\ x_9! = x_{10} \\ x_{12}! = x_{13} \\ x_{15}! = x_{16} \\ x_2 \cdot x_4 = x_5 \\ x_5 \cdot x_6 = x_7 \\ x_7 \cdot x_9 = x_{10} \\ x_4 \cdot x_{11} = x_{12} \\ x_3 \cdot x_{12} = x_{13} \\ x_9 \cdot x_{14} = x_{15} \\ x_8 \cdot x_{15} = x_{16} \end{array} \right.$$

Lemma 1 and the diagram in Figure 4 explain the construction of the system  $C$ .



**Fig. 4** Construction of the system  $C$

**Lemma 11.** *For every  $x_4, x_9 \in \mathbb{N} \setminus \{0, 1, 2\}$ , the system  $C$  is solvable in positive integers  $x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}$  if and only if  $x_4$  and  $x_9$  are prime and  $x_4 + 2 = x_9$ . In this case, the integers  $x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}$  are uniquely determined by the following equalities:*

$$\begin{aligned}
x_1 &= x_4 - 1 \\
x_2 &= (x_4 - 1)! \\
x_3 &= ((x_4 - 1)!)! \\
x_5 &= x_4! \\
x_6 &= x_9 - 1 \\
x_7 &= (x_9 - 1)! \\
x_8 &= ((x_9 - 1)!)! \\
x_{10} &= x_9! \\
x_{11} &= \frac{(x_4 - 1)! + 1}{x_4} \\
x_{12} &= (x_4 - 1)! + 1 \\
x_{13} &= ((x_4 - 1)! + 1)! \\
x_{14} &= \frac{(x_9 - 1)! + 1}{x_9} \\
x_{15} &= (x_9 - 1)! + 1 \\
x_{16} &= ((x_9 - 1)! + 1)!
\end{aligned}$$

*Proof.* By Lemma 1, for every  $x_4, x_9 \in \mathbb{N} \setminus \{0, 1, 2\}$ , the system  $C$  is solvable in positive integers  $x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}$  if and only if

$$(x_4 + 2 = x_9) \wedge (x_4 | ((x_4 - 1)! + 1)) \wedge (x_9 | ((x_9 - 1)! + 1))$$

Hence, the claim of Lemma 11 follows from Lemma 4.  $\square$

**Lemma 12.** *There are only finitely many tuples  $(x_1, \dots, x_{16}) \in (\mathbb{N} \setminus \{0\})^{16}$  which solve the system  $C$  and satisfy  $(x_4 \in \{1, 2\}) \vee (x_9 \in \{1, 2\})$ .*



*Proof.* If a tuple  $(x_1, \dots, x_{16}) \in (\mathbb{N} \setminus \{0\})^{16}$  solves the system  $C$  and  $(x_4 \in \{1, 2\}) \vee (x_9 \in \{1, 2\})$ , then  $x_1, \dots, x_{16} \leq 7!$ . Indeed, for example, if  $x_4 = 2$  then  $x_6 = x_4 + 1 = 3$ . Hence,  $x_7 = x_6! = 6$ . Therefore,  $x_{15} = x_7 + 1 = 7$ . Consequently,  $x_{16} = x_{15}! = 7!$ .  $\square$

**Theorem 7.** *The statement  $\Psi_{16}$  proves the following implication: if there exists a twin prime greater than  $g(14)$ , then there are infinitely many twin primes.*

*Proof.* Suppose that the antecedent holds. Then, there exist prime numbers  $x_4$  and  $x_9$  such that  $x_9 = x_4 + 2 > g(14)$ . Hence,  $x_4, x_9 \in \mathbb{N} \setminus \{0, 1, 2\}$ . By Lemma 11, there exists a unique tuple  $(x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}) \in (\mathbb{N} \setminus \{0\})^{14}$  such that the tuple  $(x_1, \dots, x_{16})$  solves the system  $C$ . Since  $x_9 > g(14)$ , we obtain that  $x_9 - 1 \geq g(14)$ . Therefore,  $(x_9 - 1)! \geq g(14)! = g(15)$ . Hence,  $(x_9 - 1)! + 1 > g(15)$ . Consequently,

$$x_{16} = ((x_9 - 1)! + 1)! > g(15)! = g(16)$$

Since  $C \subseteq B_{16}$ , the statement  $\Psi_{16}$  and the inequality  $x_{16} > g(16)$  imply that the system  $C$  has infinitely many solutions in positive integers  $x_1, \dots, x_{16}$ . According to Lemmas 11 and 12, there are infinitely many twin primes.  $\square$

**Corollary 4.** (cf. [6]). *Let  $\mathcal{X}_{16}$  denote the set of twin primes. The statement  $\Psi_{16}$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{X}_{16}$ , and this number equals  $\max(\mathcal{X}_{16})$ , if  $\mathcal{X}_{16}$  is finite. Assuming the statement  $\Psi_{16}$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{X}_{16}$ . Assuming the statement  $\Psi_{16}$ , the infinity of  $\mathcal{X}_{16}$  is decidable in the limit.*


*Proof.* We consider an algorithm which computes  $\max(\mathcal{X}_{16} \cap [1, g(14)])$ .  $\square$

## 9 Hypothetical statements $\Delta_5, \dots, \Delta_{14}$ and their consequences

Let  $\lambda(5) = \Gamma(25)$ , and let  $\lambda(n + 1) = \Gamma(\lambda(n))$  for every integer  $n \geq 5$ . For an integer  $n \geq 5$ , let  $\mathcal{J}_n$  denote the following system of equations:

$$\left\{ \begin{array}{l} \forall i \in \{1, \dots, n-1\} \setminus \{3\} \Gamma(x_i) = x_{i+1} \\ x_1 \cdot x_1 = x_4 \\ x_2 \cdot x_3 = x_5 \end{array} \right.$$

Lemma 2 and the diagram in Figure 5 explain the construction of the system  $\mathcal{J}_n$ .



**Fig. 5** Construction of the system  $\mathcal{J}_n$

For every integer  $n \geq 5$ , the system  $\mathcal{J}_n$  has exactly two solutions in positive integers, namely  $(1, \dots, 1)$  and  $(5, 24, 23!, 25, \lambda(5), \dots, \lambda(n))$ . For an integer  $n \geq 5$ , let  $\Delta_n$  denote the following statement: *if a system of equations  $\mathcal{S} \subseteq \{\Gamma(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq \lambda(n)$ .*

**Hypothesis 2.** *The statements  $\Delta_5, \dots, \Delta_{14}$  are true.*

Lemmas 2 and 4 imply that the statements  $\Delta_n$  have similar consequences as the statements  $\Psi_n$ .

**Theorem 8.** *The statement  $\Delta_6$  implies that any prime number  $p \geq 25$  proves the infinitude of primes.*

*Proof.* It follows from Lemmas 2 and 4. We leave the details to the reader.  $\square$

## 10 Hypothetical statements $\Sigma_3, \dots, \Sigma_{16}$ and their consequences

Let  $\Gamma_n(k)$  denote  $(k-1)!$ , where  $n \in \{3, \dots, 16\}$  and  $k \in \{2\} \cup [2^{2^{n-3}} + 1, \infty) \cap \mathbb{N}$ . For an integer  $n \in \{3, \dots, 16\}$ , let

$$Q_n = \{\Gamma_n(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$$

For an integer  $n \in \{3, \dots, 16\}$ , let  $P_n$  denote the following system of equations:

$$\begin{cases} x_1 \cdot x_1 = x_1 \\ \Gamma_n(x_2) = x_1 \\ \forall i \in \{2, \dots, n-1\} x_i \cdot x_i = x_{i+1} \end{cases}$$


**Lemma 13.** *For every integer  $n \in \{3, \dots, 16\}$ ,  $P_n \subseteq Q_n$  and the system  $P_n$  with  $\Gamma$  instead of  $\Gamma_n$  has exactly one solution in positive integers  $x_1, \dots, x_n$ , namely  $(1, 2^{2^0}, 2^{2^1}, 2^{2^2}, \dots, 2^{2^{n-2}})$ .*

For an integer  $n \in \{3, \dots, 16\}$ , let  $\Sigma_n$  denote the following statement: *if a system of equations  $\mathcal{S} \subseteq Q_n$  with  $\Gamma$  instead of  $\Gamma_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then every tuple  $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$  that solves the original system  $\mathcal{S}$  satisfies  $x_1, \dots, x_n \leq 2^{2^{n-2}}$ .*

**Hypothesis 3.** *The statements  $\Sigma_3, \dots, \Sigma_{16}$  are true.*

**Lemma 14.** *(cf. Lemma 2). For every integer  $n \in \{4, \dots, 16\}$  and for every positive integers  $x$  and  $y$ ,  $x \cdot \Gamma_n(x) = \Gamma_n(y)$  if and only if  $(x+1 = y) \wedge (x \geq 2^{2^{n-3}} + 1)$ .*

Let  $\mathcal{Z}_9 \subseteq \mathcal{Q}_9$  be the system of equations in Figure 6.



**Fig. 6** Construction of the system  $\mathcal{Z}_9$

**Lemma 15.** For every positive integer  $x_1$ , the system  $\mathcal{Z}_9$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1 > 2^{2^{9-4}}$  and  $x_1^2 + 1$  is prime. In this case, positive integers  $x_2, \dots, x_9$  are uniquely determined by  $x_1$ . For every positive integer  $n$ , at most finitely many tuples  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  begin with  $n$  and solve the system  $\mathcal{Z}_9$  with  $\Gamma$  instead of  $\Gamma_9$ .

*Proof.* It follows from Lemmas 2, 4, and 14. □

**Lemma 16.** ([17]). The number  $(13!)^2 + 1 = 38775788043632640001$  is prime.

**Lemma 17.**  $\left( (13!)^2 \geq 2^{2^{9-3}} + 1 = 18446744073709551617 \right) \wedge \left( \Gamma_9((13!)^2) > 2^{2^{9-2}} \right)$ .

**Theorem 9.** The statement  $\Sigma_9$  implies the infinitude of primes of the form  $n^2 + 1$ .

*Proof.* It follows from Lemmas 15–17. □


**Theorem 10.** (cf. Theorem 6). The statement  $\Sigma_9$  implies that any prime of the form  $n! + 1$  with  $n \geq 2^{2^{9-3}}$  proves the infinitude of primes of the form  $n! + 1$ .

*Proof.* We leave the proof to the reader. □

**Corollary 5.** Let  $\mathcal{Y}_9$  denote the set of primes of the form  $n! + 1$ . The statement  $\Sigma_9$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{Y}_9$ , and this number equals  $\max(\mathcal{Y}_9)$ , if  $\mathcal{Y}_9$  is finite. Assuming the statement  $\Sigma_9$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{Y}_9$ . Assuming the statement  $\Sigma_9$ , the infinity of  $\mathcal{Y}_9$  is decidable in the limit.

*Proof.* We consider an algorithm which computes  $\max(\mathcal{Y}_9 \cap [1, (2^{2^{9-3}} - 1)! + 1])$ . □

Let  $\mathcal{Z}_{14} \subseteq \mathcal{Q}_{14}$  be the system of equations in Figure 7.



**Fig. 7** Construction of the system  $\mathcal{Z}_{14}$

**Lemma 18.** *For every positive integer  $x_1$ , the system  $\mathcal{Z}_{14}$  is solvable in positive integers  $x_2, \dots, x_{14}$  if and only if  $x_1$  and  $x_1 + 2$  are prime and  $x_1 \geq 2^{2^{14-3}} + 1$ . In this case, positive integers  $x_2, \dots, x_{14}$  are uniquely determined by  $x_1$ . For every positive integer  $n$ , at most finitely many tuples  $(x_1, \dots, x_{14}) \in (\mathbb{N} \setminus \{0\})^{14}$  begin with  $n$  and solve the system  $\mathcal{Z}_{14}$  with  $\Gamma$  instead of  $\Gamma_{14}$ .*

*Proof.* It follows from Lemmas 2, 4, and 14. □


**Lemma 19.** ([21, p. 87]). *The numbers  $459 \cdot 2^{8529} - 1$  and  $459 \cdot 2^{8529} + 1$  are prime (Harvey Dubner).*

**Lemma 20.**  $459 \cdot 2^{8529} - 1 > 2^{2^{14-2}} = 2^{4096}$ .

**Theorem 11.** *The statement  $\Sigma_{14}$  implies the infinitude of twin primes.*

*Proof.* It follows from Lemmas 18–20. □

A prime  $p$  is said to be a Sophie Germain prime if both  $p$  and  $2p + 1$  are prime, see [20]. It is conjectured that there are infinitely many Sophie Germain primes, see [15, p. 330]. Let  $\mathcal{Z}_{16} \subseteq \mathcal{Q}_{16}$  be the system of equations in Figure 8.



**Fig. 8** Construction of the system  $\mathcal{Z}_{16}$

**Lemma 21.** For every positive integer  $x_1$ , the system  $\mathcal{Z}_{16}$  is solvable in positive integers  $x_2, \dots, x_{16}$  if and only if  $x_1$  is a Sophie Germain prime and  $x_1 \geq 2^{2^{16-3}} + 1$ . In this case, positive integers  $x_2, \dots, x_{16}$  are uniquely determined by  $x_1$ . For every positive integer  $n$ , at most finitely many tuples  $(x_1, \dots, x_{16}) \in (\mathbb{N} \setminus \{0\})^{16}$  begin with  $n$  and solve the system  $\mathcal{Z}_{16}$  with  $\Gamma$  instead of  $\Gamma_{16}$ .

*Proof.* It follows from Lemmas 2, 4, and 14. □

**Lemma 22.** ([15, p. 330]).  $8069496435 \cdot 10^{5072} - 1$  is a Sophie Germain prime (Harvey Dubner).

**Lemma 23.**  $8069496435 \cdot 10^{5072} - 1 > 2^{2^{16-2}}$ .

**Theorem 12.** The statement  $\Sigma_{16}$  implies the infinitude of Sophie Germain primes.

*Proof.* It follows from Lemmas 21–23. □

**Theorem 13.** The statement  $\Sigma_6$  proves the following implication: if the equation  $x(x+1) = y!$  has only finitely many solutions in positive integers  $x$  and  $y$ , then each such solution  $(x, y)$  belongs to the set  $\{(1, 2), (2, 3)\}$ .

*Proof.* We leave the proof to the reader. □

The question of solving the equation  $x(x+1) = y!$  was posed by P. Erdős, see [2]. F. Luca proved that the *abc* conjecture implies that the equation  $x(x+1) = y!$  has only finitely many solutions in positive integers, see [10].

**Theorem 14.** The statement  $\Sigma_6$  proves the following implication: if the equation  $x! + 1 = y^2$  has only finitely many solutions in positive integers  $x$  and  $y$ , then each such solution  $(x, y)$  belongs to the set  $\{(4, 5), (5, 11), (7, 71)\}$ .

*Proof.* We leave the proof to the reader. □

## 11 Hypothetical statements $\Omega_3, \dots, \Omega_{16}$ and their consequences

For an integer  $n \in \{3, \dots, 16\}$ , let  $\Omega_n$  denote the following statement: *if a system of equations  $\mathcal{S} \subseteq \{\Gamma(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$  has a solution in integers  $x_1, \dots, x_n$  greater than  $2^{n-2}$ , then  $\mathcal{S}$  has infinitely many solutions in positive integers  $x_1, \dots, x_n$ .* For every  $n \in \{3, \dots, 16\}$ , the statement  $\Sigma_n$  implies the statement  $\Omega_n$ .

**Lemma 24.** *The number  $(65!)^2 + 1$  is prime and  $65! > 2^{2^9-2}$ .*

*Proof.* The following PARI/GP ([14]) command

```
(04:04) gp > isprime((65!)^2+1, {flag=2})
%1 = 1
```

is shown together with its output. This command performs the APRCL primality test, the best deterministic primality test algorithm ([21, p. 226]). It rigorously shows that the number  $(65!)^2 + 1$  is prime.  $\square$

**Lemma 25.** *If positive integers  $x_1, \dots, x_9$  solve the system  $\mathcal{Z}_9$  and  $x_1 > 2^{2^9-2}$ , then  $x_1 = \min(x_1, \dots, x_9)$ .*

**Theorem 15.** *The statement  $\Omega_9$  implies the infinitude of primes of the form  $n^2 + 1$ .*

*Proof.* It follows from Lemmas 15 and 24–25.  $\square$

**Lemma 26.** *If positive integers  $x_1, \dots, x_{14}$  solve the system  $\mathcal{Z}_{14}$  and  $x_1 > 2^{2^{14}-2}$ , then  $x_1 = \min(x_1, \dots, x_{14})$ .*

**Theorem 16.** *The statement  $\Omega_{14}$  implies the infinitude of twin primes.*

*Proof.* It follows from Lemmas 18–20 and 26.  $\square$

## 12 Are there infinitely many composite Fermat numbers?

Integers of the form  $2^{2^n} + 1$  are called Fermat numbers. Primes of the form  $2^{2^n} + 1$  are called Fermat primes, as Fermat conjectured that every integer of the form  $2^{2^n} + 1$  is prime, see [9, p. 1]. Fermat correctly remarked that  $2^{2^0} + 1 = 3$ ,  $2^{2^1} + 1 = 5$ ,  $2^{2^2} + 1 = 17$ ,  $2^{2^3} + 1 = 257$ , and  $2^{2^4} + 1 = 65537$  are all prime, see [9, p. 1].

**Open Problem.** ([9, p. 159]). *Are there infinitely many composite numbers of the form  $2^{2^n} + 1$ ?*

Most mathematicians believe that  $2^{2^n} + 1$  is composite for every integer  $n \geq 5$ , see [8, p. 23]. Let

$$H_n = \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\} \cup \{2^{2^{x_i}} = x_k : i, k \in \{1, \dots, n\}\}$$

Let  $h(1) = 1$ , and let  $h(n+1) = 2^{2^{h(n)}}$  for every positive integer  $n$ .

**Lemma 27.** *The following subsystem of  $H_n$*

$$\begin{cases} x_1 \cdot x_1 = x_1 \\ \forall i \in \{1, \dots, n-1\} 2^{2^{x_i}} = x_{i+1} \end{cases}$$

*has exactly one solution  $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$ , namely  $(h(1), \dots, h(n))$ .*

For a positive integer  $n$ , let  $\xi_n$  denote the following statement: *if a system of equations  $S \subseteq H_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq h(n)$* . The statement  $\xi_n$  says that for subsystems of  $H_n$  the largest known solution is indeed the largest possible.

**Hypothesis 4.** *The statements  $\xi_1, \dots, \xi_{13}$  are true.*

**Proposition 4.** *Every statement  $\xi_n$  is true with an unknown integer bound that depends on  $n$ .*


*Proof.* For every positive integer  $n$ , the system  $H_n$  has a finite number of subsystems. □

**Theorem 17.** *The statement  $\xi_{13}$  proves the following implication: if  $z \in \mathbb{N} \setminus \{0\}$  and  $2^{2^z} + 1$  is composite and greater than  $h(12)$ , then  $2^{2^z} + 1$  is composite for infinitely many positive integers  $z$ .*

*Proof.* Let us consider the equation

$$(x + 1)(y + 1) = 2^{2^z} + 1 \tag{2}$$

in positive integers. By Lemma 3, we can transform equation (2) into an equivalent system of equations  $\mathcal{G}$  which has 13 variables ( $x, y, z$ , and 10 other variables) and which consists of equations of the forms  $\alpha \cdot \beta = \gamma$  and  $2^{2^\alpha} = \gamma$ , see the diagram in Figure 9.



**Fig. 9** Construction of the system  $\mathcal{G}$

Since  $2^{2^z} + 1 > h(12)$ , we obtain that  $2^{2^{2^z} + 1} > h(13)$ . By this, the statement  $\xi_{13}$  implies that the system  $\mathcal{G}$  has infinitely many solutions in positive integers. It means that there are infinitely many composite Fermat numbers. □

**Corollary 6.** Let  $\mathcal{W}_{13}$  denote the set of composite Fermat numbers. The statement  $\xi_{13}$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{W}_{13}$ , and this number equals  $\max(\mathcal{W}_{13})$ , if  $\mathcal{W}_{13}$  is finite. Assuming the statement  $\xi_{13}$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{W}_{13}$ . Assuming the statement  $\xi_{13}$ , the infinity of  $\mathcal{W}_{13}$  is decidable in the limit.

*Proof.* We consider an algorithm which computes  $\max(\mathcal{W}_{13} \cap [1, h(12)])$ . □

## References

- [1] C. H. Bennett, *Chaitin's Omega*, in: *Fractal music, hypercards, and more ...* (M. Gardner, ed.), W. H. Freeman, New York, 1992, 307–319.
- [2] D. Berend and J. E. Harmse, *On polynomial-factorial Diophantine equations*, *Trans. Amer. Math. Soc.* 358 (2006), no. 4, 1741–1779.
- [3] C. K. Caldwell and Y. Gallot, *On the primality of  $n! \pm 1$  and  $2 \times 3 \times 5 \times \cdots \times p \pm 1$* , *Math. Comp.* 71 (2002), no. 237, 441–448, <http://doi.org/10.1090/S0025-5718-01-01315-1>.
- [4] C. S. Calude, H. Jürgensen, S. Legg, *Solving problems with finite test sets*, in: *Finite versus Infinite: Contributions to an Eternal Dilemma* (C. Calude and G. Păun, eds.), 39–52, Springer, London, 2000.
- [5] N. C. A. da Costa and F. A. Doria, *On the foundations of science (LIVRO): essays, first series*, E-papers Serviços Editoriais Ltda, Rio de Janeiro, 2013.
- [6] F. G. Dorais, *Can the twin prime problem be solved with a single use of a halting oracle?* July 23, 2011, <http://mathoverflow.net/questions/71050>.
- [7] M. Erickson, A. Vazzana, D. Garth, *Introduction to number theory*, 2nd ed., CRC Press, Boca Raton, FL, 2016.
- [8] J.-M. De Koninck and F. Luca, *Analytic number theory: Exploring the anatomy of integers*, American Mathematical Society, Providence, RI, 2012.
- [9] M. Křížek, F. Luca, L. Somer, *17 lectures on Fermat numbers: from number theory to geometry*, Springer, New York, 2001.
- [10] F. Luca, *The Diophantine equation  $P(x) = n!$  and a result of M. Overholt*, *Glas. Mat. Ser. III* 37 (57) (2002), no. 2, 269–273
- [11] M. Mignotte and A. Pethő, *On the Diophantine equation  $x^p - x = y^q - y$* , *Publ. Mat.* 43 (1999), no. 1, 207–216.
- [12] W. Narkiewicz, *Rational number theory in the 20th century: From PNT to FLT*, Springer, London, 2012.
- [13] M. Overholt, *The Diophantine equation  $n! + 1 = m^2$* , *Bull. London Math. Soc.* 25 (1993), no. 2, 104.
- [14] PARI/GP online documentation, [http://pari.math.u-bordeaux.fr/dohtml/html/Arithmetic\\_functions.html](http://pari.math.u-bordeaux.fr/dohtml/html/Arithmetic_functions.html).
- [15] P. Ribenboim, *The new book of prime number records*, Springer, New York, 1996, <http://doi.org/10.1007/978-1-4612-0759-7>.
- [16] S. Siksek, *Chabauty and the Mordell–Weil Sieve*, in: *Advances on Superelliptic Curves and Their Applications* (eds. L. Beshaj, T. Shaska, E. Zhupa), 194–224, IOS Press, Amsterdam, 2015, <http://dx.doi.org/10.3233/978-1-61499-520-3-194>.



- [17] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, *Smallest prime factor of*  $A020549(n) = (n!)^2 + 1$ , <http://oeis.org/A282706>.
- [18] E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics*, 2nd ed., Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [19] Wolfram MathWorld, *Perfect Cuboid*, <http://mathworld.wolfram.com/PerfectCuboid.html>.
- [20] Wolfram MathWorld, *Sophie Germain prime*, <http://mathworld.wolfram.com/SophieGermainPrime.html>.
- [21] S. Y. Yan, *Number theory for computing*, 2nd ed., Springer, Berlin, 2002.
- [22] A. A. Zenkin, *Super-induction method: logical acupuncture of mathematical infinity*, Twentieth World Congress of Philosophy, Boston, MA, August 10–15, 1998, <http://www.bu.edu/wcp/Papers/Logi/LogiZenk.htm>.
- [23] A. A. Zenkin, *Superinduction: new logical method for mathematical proofs with a computer*, in: J. Cachro and K. Kijania-Placek (eds.), Volume of Abstracts, 11th International Congress of Logic, Methodology and Philosophy of Science, August 20–26, 1999, Cracow, Poland, p. 94, The Faculty of Philosophy, Jagiellonian University, Cracow, 1999.

Apoloniusz Tyszka  
 University of Agriculture  
 Faculty of Production and Power Engineering  
 Balicka 116B, 30-149 Kraków, Poland  
 E-mail: rttyszka@cyf-kr.edu.pl