

On *ZFC*-formulae  $\varphi(x)$  for which we know a non-negative integer  $n$  such that  $\{x \in \mathbb{N} : \varphi(x)\} \subseteq \{x \in \mathbb{N} : x \leq n - 1\}$  if the set  $\{x \in \mathbb{N} : \varphi(x)\}$  is finite

Apoloniusz Tyszką

### Abstract

Let  $g(3) = 4$ , and let  $g(n + 1) = g(n)!$  for every integer  $n \geq 3$ . For an integer  $n \in \{3, \dots, 16\}$ , let  $\Psi_n$  denote the following statement: *if a system of equations  $S \subseteq \{x_i! = x_k : (i, k \in \{1, \dots, n\}) \wedge (i \neq k)\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq g(n)$* . For every statement  $\Psi_n$ , the bound  $g(n)$  cannot be decreased. The author's hypothesis says that the statements  $\Psi_3, \dots, \Psi_{16}$  hold true. We say that a non-negative integer  $m$  is a threshold number of a set  $X \subseteq \mathbb{N}$ , if  $X$  is infinite if and only if  $X$  contains an element greater than  $m$ . The following problem is open: *define a set  $X \subseteq \mathbb{N}$  that satisfies the following conditions: (1) the relation  $n \in X$  is qualitatively the same for every  $n \in \mathbb{N}$ , (2) a known algorithm for every  $n \in \mathbb{N}$  decides whether or not  $n \in X$ , (3) a known algorithm returns a threshold number of  $X$ , (4) new elements of  $X$  are still discovered, (5) we do not know any algorithm deciding the finiteness of  $X$* . We define a set  $X \subseteq \mathbb{N}$  that satisfies conditions (2)–(5). The statement  $\Psi_9$  implies that the set of primes of the form  $n^2 + 1$  solves the problem and the set of primes of the form  $n! + 1$  solves the problem. The statement  $\Psi_{16}$  implies that the set of twin primes solves the problem.

**Key words and phrases:** Brocard-Ramanujan equation  $x! + 1 = y^2$ , composite Fermat numbers, decidability in the limit, finiteness of a set, incompleteness of *ZFC*, infiniteness of a set, prime numbers of the form  $n^2 + 1$ , prime numbers of the form  $n! + 1$ , single query to an oracle for the halting problem, twin primes.

**2010 Mathematics Subject Classification:** 03D20, 11A41.

## 1 Introduction

The phrase "we know a non-negative integer  $n$ " in the title means that we know an algorithm which returns  $n$ . The title cannot be formalised in *ZFC* because the phrase "we know a non-negative integer  $n$ " refers to currently known non-negative integers  $n$  with some property. A formally stated title may look like this: On *ZFC*-formulae  $\varphi(x)$  for which there exists a non-negative integer  $n$  such that *ZFC* proves that

$$\text{card}(\{x \in \mathbb{N} : \varphi(x)\}) < \infty \implies \{x \in \mathbb{N} : \varphi(x)\} \subseteq \{x \in \mathbb{N} : x \leq n - 1\}$$

Unfortunately, this formulation admits formulae  $\varphi(x)$  without any known non-negative integer  $n$  such that *ZFC* proves the above implication.

**Lemma 1.** *For every non-negative integer  $n$ ,  $\text{card}(\{x \in \mathbb{N} : x \leq n - 1\}) = n$ .*

**Corollary 1.** *The title altered to "On *ZFC*-formulae  $\varphi(x)$  for which we know a non-negative integer  $n$  such that  $\text{card}(\{x \in \mathbb{N} : \varphi(x)\}) \leq n$  if the set  $\{x \in \mathbb{N} : \varphi(x)\}$  is finite" involves a weaker assumption on  $\varphi(x)$ .*

## 2 Subsets of $\mathbb{N}$ and their threshold numbers

We say that a non-negative integer  $m$  is a threshold number of a set  $X \subseteq \mathbb{N}$ , if  $X$  is infinite if and only if  $X$  contains an element greater than  $m$ , cf. [27] and [28]. If a set  $X \subseteq \mathbb{N}$  is empty or infinite, then any non-negative integer  $m$  is a threshold number of  $X$ . If a set  $X \subseteq \mathbb{N}$  is non-empty and finite, then the all threshold numbers of  $X$  form the set  $\{\max(X), \max(X) + 1, \max(X) + 2, \dots\}$ .

It is conjectured that the set of prime numbers of the form  $n^2 + 1$  is infinite, see [16, pp. 37–38]. It is conjectured that the set of prime numbers of the form  $n! + 1$  is infinite, see [3, p. 443]. A twin prime is a prime number that differs from another prime number by 2. The twin prime conjecture states that the set of twin primes is infinite, see [16, p. 39]. It is conjectured that the set of composite numbers of the form  $2^{2^n} + 1$  is infinite, see [11, p. 23] and [12, pp. 158–159]. A prime  $p$  is said to be a Sophie Germain prime if both  $p$  and  $2p + 1$  are prime, see [25]. It is conjectured that the set of Sophie Germain primes is infinite, see [19, p. 330]. For each of these sets, we do not know any threshold number.

The following statement: *for every non-negative integer  $n$  there exist*

$$\text{prime numbers } p \text{ and } q \text{ such that } p + 2 = q \text{ and } p \in \left[10^n, 10^{n+1}\right] \quad (\text{T})$$

is a  $\Pi_1$  statement which strengthens the twin prime conjecture, see [4, p. 43]. C. H. Bennett claims that most mathematical conjectures can be settled indirectly by proving stronger  $\Pi_1$  statements, see [1]. The statement (T) is equivalent to the non-halting of a Turing machine. If a set  $X \subseteq \mathbb{N}$  is computable and we know a threshold number of  $X$ , then the infinity of  $X$  is equivalent to the halting of a Turing machine.

The height of a rational number  $\frac{p}{q}$  is denoted by  $H\left(\frac{p}{q}\right)$  and equals  $\max(|p|, |q|)$  provided  $\frac{p}{q}$  is written in lowest terms. The height of a rational tuple  $(x_1, \dots, x_n)$  is denoted by  $H(x_1, \dots, x_n)$  and equals  $\max(H(x_1), \dots, H(x_n))$ .

**Proposition 1.** *The equation  $x^5 - x = y^2 - y$  has only finitely many rational solutions, see [15, p. 212]. The known rational solutions are  $(x, y) = (-1, 0), (-1, 1), (0, 0), (0, 1), (1, 0), (1, 1), (2, -5), (2, 6), (3, -15), (3, 16), (30, -4929), (30, 4930), \left(\frac{1}{4}, \frac{15}{32}\right), \left(\frac{1}{4}, \frac{17}{32}\right), \left(-\frac{15}{16}, -\frac{185}{1024}\right), \left(-\frac{15}{16}, \frac{1209}{1024}\right)$ , and the existence of other solutions is an open question, see [20, pp. 223–224].*

**Corollary 2.** *The set  $\mathcal{T} = \{n \in \mathbb{N} : \text{the equation } x^5 - x = y^2 - y \text{ has a rational solution of height } n\}$  is finite. We know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{T}$ . We do not know any algorithm which returns a threshold number of  $\mathcal{T}$ .*

**Open Problem 1.** (cf. Corollary 3) *Define a set  $X \subseteq \mathbb{N}$  that satisfies the following conditions:*

- (1) *the relation  $n \in X$  is qualitatively the same for every  $n \in \mathbb{N}$ ,*
- (2) *a known algorithm for every  $n \in \mathbb{N}$  decides whether or not  $n \in X$ ,*
- (3) *a known algorithm returns a threshold number of  $X$ ,*
- (4) *new elements of  $X$  are still discovered,*
- (5) *we do not know any algorithm deciding the finiteness of  $X$ .*

Let  $\mathcal{L}$  denote the following system of equations:

$$\begin{cases} x^2 + y^2 = s^2 \\ x^2 + z^2 = t^2 \\ y^2 + z^2 = u^2 \\ x^2 + y^2 + z^2 = v^2 \end{cases}$$

Let  $\mathcal{F}$  denote the set

$$\left\{k \in \mathbb{N} \setminus \{0\} : \left(\text{the system } \mathcal{L} \text{ has no solutions in } \{1, \dots, k\}^7\right) \wedge \left(\text{the system } \mathcal{L} \text{ has a solution in } \{1, \dots, k+1\}^7\right)\right\}$$

Let  $\mathcal{P}$  denote the set of prime numbers, and let  $\mathcal{Z}$  denote the set

$$\{k \in \mathbb{N} \setminus \{0\} : \text{the system } \mathcal{L} \text{ has a solution in } \{1, \dots, k\}^7\}$$

A perfect cuboid is a cuboid having integer side lengths, integer face diagonals, and an integer space diagonal.

**Proposition 2.** ([24]) *No perfect cuboids are known.*

**Corollary 3.** (cf. Open Problem 1) *The set  $\mathcal{Z} \cup \left( \left[ 2, 9^{99999} \right] \cap \mathcal{P} \right)$  satisfies conditions (2)–(5).*

**Corollary 4.** *We know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{F}$ . ZFC proves that  $\text{card}(\mathcal{F}) \in \{0, 1\}$ . We do not know any algorithm which returns  $\text{card}(\mathcal{F})$ . We do not know any algorithm which returns a threshold number of  $\mathcal{F}$ .*

Let

$$\mathcal{H} = \begin{cases} \mathbb{N}, & \text{if } \sin\left(9^{99999}\right) < 0 \\ \mathbb{N} \cap \left[ 0, \sin\left(9^{99999}\right) \cdot 9^{99999} \right) & \text{otherwise} \end{cases}$$

We do not know whether or not the set  $\mathcal{H}$  is finite.

**Proposition 3.** *The number  $9^{99999}$  is a threshold number of  $\mathcal{H}$ . We know an algorithm which decides the equality  $\mathcal{H} = \mathbb{N}$ . If  $\mathcal{H} \neq \mathbb{N}$ , then the set  $\mathcal{H}$  consists of all integers from 0 to a non-negative integer which can be computed by a known algorithm. We know an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{H}$ .*

Let

$$\mathcal{K} = \begin{cases} \{n\}, & \text{if } (n \in \mathbb{N}) \wedge (2^{\aleph_0} = \aleph_{n+1}) \\ \{0\}, & \text{if } 2^{\aleph_0} \geq \aleph_\omega \end{cases}$$

**Theorem 1.** *ZFC proves that  $\text{card}(\mathcal{K}) = 1$ . If ZFC is consistent, then for every  $n \in \mathbb{N}$  the sentences " $n$  is a threshold number of  $\mathcal{K}$ " and " $n$  is not a threshold number of  $\mathcal{K}$ " are not provable in ZFC. If ZFC is consistent, then for every  $n \in \mathbb{N}$  the sentences " $n \in \mathcal{K}$ " and " $n \notin \mathcal{K}$ " are not provable in ZFC.*

*Proof.* It suffices to observe that  $2^{\aleph_0}$  can attain every value from the set  $\{\aleph_1, \aleph_2, \aleph_3, \dots\}$ , see [7] and [10, p. 232].  $\square$

### 3 A Diophantine equation whose non-solvability expresses the consistency of ZFC

Gödel's second incompleteness theorem and the Davis-Putnam-Robinson- Matiyasevich theorem imply the following theorem.

**Theorem 2.** ([5, p. 35]) *There exists a polynomial  $D(x_1, \dots, x_m)$  with integer coefficients such that if ZFC is arithmetically consistent, then the sentences "The equation  $D(x_1, \dots, x_m) = 0$  is solvable in non-negative integers" and "The equation  $D(x_1, \dots, x_m) = 0$  is not solvable in non-negative integers" are not provable in ZFC.*

**Remark 1.** ([9, p. 53]) *The polynomial  $D(x_1, \dots, x_m)$  is not effectively known.*

Let  $\mathcal{Y}$  denote the set of all non-negative integers  $k$  such that the equation  $D(x_1, \dots, x_m) = 0$  has no solutions in  $\{0, \dots, k\}^m$ . Since the set  $\{0, \dots, k\}^m$  is finite, there exists an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{Y}$ . Theorem 2 implies the next theorem.

**Theorem 3.** *For every  $n \in \mathbb{N}$ , ZFC proves that  $n \in \mathcal{Y}$ . If ZFC is arithmetically consistent, then the sentences " $\mathcal{Y}$  is finite" and " $\mathcal{Y}$  is infinite" are not provable in ZFC. If ZFC is arithmetically consistent, then for every  $n \in \mathbb{N}$  the sentences " $n$  is a threshold number of  $\mathcal{Y}$ " and " $n$  is not a threshold number of  $\mathcal{Y}$ " are not provable in ZFC.*

Let  $\mathcal{E}$  denote the set of all non-negative integers  $k$  such that the equation  $D(x_1, \dots, x_m) = 0$  has a solution in  $\{0, \dots, k\}^m$ . Since the set  $\{0, \dots, k\}^m$  is finite, there exists an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{E}$ . Theorem 2 implies the next theorem.

**Theorem 4.** *The set  $\mathcal{E}$  is empty or infinite. In both cases, every non-negative integer  $n$  is a threshold number of  $\mathcal{E}$ . If ZFC is arithmetically consistent, then the sentences " $\mathcal{E}$  is empty", " $\mathcal{E}$  is not empty", " $\mathcal{E}$  is finite", and " $\mathcal{E}$  is infinite" are not provable in ZFC.*

Let  $\mathcal{V}$  denote the set

$$\{k \in \mathbb{N} : (\text{the polynomial } D(x_1, \dots, x_m) \text{ has no solutions in } \{0, \dots, k\}^m) \wedge \\ (\text{the polynomial } D(x_1, \dots, x_m) \text{ has a solution in } \{0, \dots, k+1\}^m)\}.$$

Since the sets  $\{0, \dots, k\}^m$  and  $\{0, \dots, k+1\}^m$  are finite, there exists an algorithm which for every  $n \in \mathbb{N}$  decides whether or not  $n \in \mathcal{V}$ . According to Remark 1, at present we are not able to write a computer program that realizes such an algorithm. Theorem 2 implies the next theorem.

**Theorem 5.** (6) *ZFC proves that  $\text{card}(\mathcal{V}) \in \{0, 1\}$ .* (7) *For every  $n \in \mathbb{N}$ , ZFC proves that  $n \notin \mathcal{V}$ .* (8) *ZFC does not prove the emptiness of  $\mathcal{V}$ , if ZFC is arithmetically consistent.* (9) *For every  $n \in \mathbb{N}$ , the sentence " $n$  is a threshold number of  $\mathcal{V}$ " is not provable in ZFC, if ZFC is arithmetically consistent.* (10) *For every  $n \in \mathbb{N}$ , the sentence " $n$  is not a threshold number of  $\mathcal{V}$ " is not provable in ZFC, if ZFC is arithmetically consistent.*

**Open Problem 2.** *Define a simple algorithm  $A$  such that  $A$  returns 0 or 1 on every input  $k \in \mathbb{N}$  and the set*

$$\mathcal{V} = \{k \in \mathbb{N} : \text{the program } A \text{ returns 1 on input } k\}$$

*satisfies conditions (6)–(10).*

## 4 Basic lemmas

**Lemma 2.** *For every positive integers  $x$  and  $y$ ,  $x! \cdot y = y!$  if and only if*

$$(x + 1 = y) \vee (x = y = 1)$$

Let  $\Gamma(k)$  denote  $(k - 1)!$ .

**Lemma 3.** *For every positive integers  $x$  and  $y$ ,  $x \cdot \Gamma(x) = \Gamma(y)$  if and only if*

$$(x + 1 = y) \vee (x = y = 1)$$

**Lemma 4.** *For every non-negative integers  $b$  and  $c$ ,  $b + 1 = c$  if and only if*

$$2^{2^b} \cdot 2^{2^b} = 2^{2^c}$$


**Lemma 5.** *(Wilson's theorem, [8, p. 89]). For every positive integer  $x$ ,  $x$  divides  $(x - 1)! + 1$  if and only if  $x = 1$  or  $x$  is prime.*

## 5 Hypothetical statements $\Psi_3, \dots, \Psi_{16}$

For an integer  $n \geq 3$ , let  $\mathcal{U}_n$  denote the following system of equations:

$$\begin{cases} \forall i \in \{1, \dots, n-1\} \setminus \{2\} & x_i! = x_{i+1} \\ & x_1 \cdot x_2 = x_3 \\ & x_2 \cdot x_2 = x_3 \end{cases}$$

The diagram in Figure 1 illustrates the construction of the system  $\mathcal{U}_n$ .



**Fig. 1** Construction of the system  $\mathcal{U}_n$

Let  $g(3) = 4$ , and let  $g(n+1) = g(n)!$  for every integer  $n \geq 3$ .

**Lemma 6.** *For every integer  $n \geq 3$ , the system  $\mathcal{U}_n$  has exactly two solutions in positive integers, namely  $(1, \dots, 1)$  and  $(2, 2, g(3), \dots, g(n))$ .*

Let

$$B_n = \{x_i! = x_k : (i, k \in \{1, \dots, n\}) \wedge (i \neq k)\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$$

For an integer  $n \geq 3$ , let  $\Psi_n$  denote the following statement: *if a system of equations  $\mathcal{S} \subseteq B_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq g(n)$ .* The statement  $\Psi_n$  says that for subsystems of  $B_n$  the largest known solution is indeed the largest possible.

**Hypothesis 1.** *The statements  $\Psi_3, \dots, \Psi_{16}$  are true.*

**Remark 2.** *By Lemma 2 and algebraic lemmas in [22, p. 110], the statement  $\forall n \in \mathbb{N} \setminus \{0, 1, 2\} \Psi_n$  implies that there is an algorithm which takes as input a factorial Diophantine equation, and returns an integer such that this integer is greater than the solutions in positive integers, if these solutions form a finite set. This conclusion is a bit strange because a computable upper bound on non-negative integer solutions does not exist for exponential Diophantine equations with a finite number of solutions, see [14, p. 300]. Therefore, the statement  $\forall n \in \mathbb{N} \setminus \{0, 1, 2\} \Psi_n$  seems to be false.*

**Lemma 7.** *Every statement  $\Psi_n$  is true with an unknown integer bound that depends on  $n$ .*

*Proof.* For every positive integer  $n$ , the system  $B_n$  has a finite number of subsystems. □

**Lemma 8.** *For every statement  $\Psi_n$ , the bound  $g(n)$  cannot be decreased.*


*Proof.* It follows from Lemma 6 because  $\mathcal{U}_n \subseteq B_n$ . □

## 6 The Brocard-Ramanujan equation $x! + 1 = y^2$

Let  $\mathcal{A}$  denote the following system of equations:

$$\begin{cases} x_1! = x_2 \\ x_2! = x_3 \\ x_5! = x_6 \\ x_4 \cdot x_4 = x_5 \\ x_3 \cdot x_5 = x_6 \end{cases}$$

Lemma 2 and the diagram in Figure 2 explain the construction of the system  $\mathcal{A}$ .



**Fig. 2** Construction of the system  $\mathcal{A}$

**Lemma 9.** For every  $x_1, x_4 \in \mathbb{N} \setminus \{0, 1\}$ , the system  $\mathcal{A}$  is solvable in positive integers  $x_2, x_3, x_5, x_6$  if and only if  $x_1! + 1 = x_4^2$ . In this case, the integers  $x_2, x_3, x_5, x_6$  are uniquely determined by the following equalities:

$$\begin{aligned} x_2 &= x_1! \\ x_3 &= (x_1!)! \\ x_5 &= x_1! + 1 \\ x_6 &= (x_1! + 1)! \end{aligned}$$

*Proof.* It follows from Lemma 2. □

It is conjectured that  $x! + 1$  is a perfect square only for  $x \in \{4, 5, 7\}$ , see [23, p. 297]. A weak form of Szpiro's conjecture implies that there are only finitely many solutions to the equation  $x! + 1 = y^2$ , see [17].

**Theorem 6.** If the equation  $x_1! + 1 = x_4^2$  has only finitely many solutions in positive integers, then the statement  $\Psi_6$  guarantees that each such solution  $(x_1, x_4)$  belongs to the set  $\{(4, 5), (5, 11), (7, 71)\}$ .


*Proof.* Suppose that the antecedent holds. Let positive integers  $x_1$  and  $x_4$  satisfy  $x_1! + 1 = x_4^2$ . Then,  $x_1, x_4 \in \mathbb{N} \setminus \{0, 1\}$ . By Lemma 9, the system  $\mathcal{A}$  is solvable in positive integers  $x_2, x_3, x_5, x_6$ . Since  $\mathcal{A} \subseteq \mathcal{B}_6$ , the statement  $\Psi_6$  implies that  $x_6 = (x_1! + 1)! \leq g(6) = g(5)!$ . Hence,  $x_1! + 1 \leq g(5) = g(4)!$ . Consequently,  $x_1 < g(4) = 24$ . If  $x_1 \in \{1, \dots, 23\}$ , then  $x_1! + 1$  is a perfect square only for  $x_1 \in \{4, 5, 7\}$ . □

## 7 Are there infinitely many prime numbers of the form $n^2 + 1$ ?

Edmund Landau's conjecture states that there are infinitely many primes of the form  $n^2 + 1$ , see [16, pp. 37–38]. Let  $\mathcal{B}$  denote the following system of equations:

$$\begin{cases} x_2! = x_3 & x_1 \cdot x_1 = x_2 \\ x_3! = x_4 & x_3 \cdot x_5 = x_6 \\ x_5! = x_6 & x_4 \cdot x_8 = x_9 \\ x_8! = x_9 & x_5 \cdot x_7 = x_8 \end{cases}$$

Lemma 2 and the diagram in Figure 3 explain the construction of the system  $\mathcal{B}$ .



**Fig. 3** Construction of the system  $\mathcal{B}$

**Lemma 10.** For every integer  $x_1 \geq 2$ , the system  $\mathcal{B}$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1^2 + 1$  is prime. In this case, the integers  $x_2, \dots, x_9$  are uniquely determined by the following equalities:

$$\begin{aligned}
 x_2 &= x_1^2 & x_7 &= \frac{(x_1^2)! + 1}{x_1^2 + 1} \\
 x_3 &= (x_1^2)! & x_8 &= (x_1^2)! + 1 \\
 x_4 &= ((x_1^2)!)! & x_9 &= ((x_1^2)! + 1)! \\
 x_5 &= x_1^2 + 1 & & \\
 x_6 &= (x_1^2 + 1)! & & 
 \end{aligned}$$

*Proof.* By Lemma 2, for every integer  $x_1 \geq 2$ , the system  $\mathcal{B}$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1^2 + 1$  divides  $(x_1^2)! + 1$ . Hence, the claim of Lemma 10 follows from Lemma 5.  $\square$

**Lemma 11.** There are only finitely many tuples  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  which solve the system  $\mathcal{B}$  and satisfy  $x_1 = 1$ .

*Proof.* If a tuple  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  solves the system  $\mathcal{B}$  and  $x_1 = 1$ , then  $x_1, \dots, x_9 \leq 2$ . Indeed,  $x_1 = 1$  implies that  $x_2 = x_1^2 = 1$ . Hence, for example,  $x_3 = x_2! = 1$ . Therefore,  $x_8 = x_3 + 1 = 2$  or  $x_8 = 1$ . Consequently,  $x_9 = x_8! \leq 2$ .  $\square$

**Theorem 7.** (cf. Theorem 11) The statement  $\Psi_9$  proves the following implication: if there exists an integer  $x_1 \geq 2$  such that  $x_1^2 + 1$  is prime and greater than  $g(7)$ , then there are infinitely many primes of the form  $n^2 + 1$ .

*Proof.* Suppose that the antecedent holds. By Lemma 10, there exists a unique tuple  $(x_2, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^8$  such that the tuple  $(x_1, x_2, \dots, x_9)$  solves the system  $\mathcal{B}$ . Since  $x_1^2 + 1 > g(7)$ , we obtain that  $x_1^2 \geq g(7)$ . Hence,  $(x_1^2)! \geq g(7)! = g(8)$ . Consequently,

$$x_9 = ((x_1^2)! + 1)! \geq (g(8) + 1)! > g(8)! = g(9)$$

Since  $\mathcal{B} \subseteq \mathcal{B}_9$ , the statement  $\Psi_9$  and the inequality  $x_9 > g(9)$  imply that the system  $\mathcal{B}$  has infinitely many solutions  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$ . According to Lemmas 10 and 11, there are infinitely many primes of the form  $n^2 + 1$ .  $\square$

**Corollary 5.** Let  $\mathcal{X}_9$  denote the set of primes of the form  $n^2 + 1$ . The statement  $\Psi_9$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{X}_9$ , and this number equals  $\max(\mathcal{X}_9)$ , if  $\mathcal{X}_9$  is finite. Assuming the statement  $\Psi_9$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{X}_9$ . Assuming the statement  $\Psi_9$ , the infinity of  $\mathcal{X}_9$  is decidable in the limit.

*Proof.* We consider an algorithm which computes  $\max(\mathcal{X}_9 \cap [1, g(7)])$ .  $\square$

## 8 Are there infinitely many prime numbers of the form $n! + 1$ ?

It is conjectured that there are infinitely many primes of the form  $n! + 1$ , see [3, p. 443].

**Theorem 8.** (cf. Theorem 12) *The statement  $\Psi_9$  proves the following implication: if there exists an integer  $x_1 \geq g(6)$  such that  $x_1! + 1$  is prime, then there are infinitely many primes of the form  $n! + 1$ .*


*Proof.* We leave the analogous proof to the reader. □

## 9 The twin prime conjecture

A twin prime is a prime number that differs from another prime number by 2. The twin prime conjecture states that there are infinitely many twin primes, see [16, p. 39]. Let  $C$  denote the following system of equations:

$$\left\{ \begin{array}{ll} x_1! = x_2 & x_2 \cdot x_4 = x_5 \\ x_2! = x_3 & x_5 \cdot x_6 = x_7 \\ x_4! = x_5 & x_7 \cdot x_9 = x_{10} \\ x_6! = x_7 & x_4 \cdot x_{11} = x_{12} \\ x_7! = x_8 & x_3 \cdot x_{12} = x_{13} \\ x_9! = x_{10} & x_9 \cdot x_{14} = x_{15} \\ x_{12}! = x_{13} & x_8 \cdot x_{15} = x_{16} \\ x_{15}! = x_{16} & \end{array} \right.$$

Lemma 2 and the diagram in Figure 4 explain the construction of the system  $C$ .



**Fig. 4** Construction of the system  $C$

**Lemma 12.** *For every  $x_4, x_9 \in \mathbb{N} \setminus \{0, 1, 2\}$ , the system  $C$  is solvable in positive integers  $x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}$  if and only if  $x_4$  and  $x_9$  are prime and  $x_4 + 2 = x_9$ . In this case, the integers  $x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}$  are uniquely determined by the following equalities:*



$$\begin{array}{ll}
x_1 = x_4 - 1 & x_{11} = \frac{(x_4 - 1)! + 1}{x_4} \\
x_2 = (x_4 - 1)! & x_{12} = (x_4 - 1)! + 1 \\
x_3 = ((x_4 - 1)!)! & x_{13} = ((x_4 - 1)! + 1)! \\
x_5 = x_4! & x_{14} = \frac{(x_9 - 1)! + 1}{x_9} \\
x_6 = x_9 - 1 & x_{15} = (x_9 - 1)! + 1 \\
x_7 = (x_9 - 1)! & x_{16} = ((x_9 - 1)! + 1)! \\
x_8 = ((x_9 - 1)!)! & \\
x_{10} = x_9! & 
\end{array}$$

*Proof.* By Lemma 2, for every  $x_4, x_9 \in \mathbb{N} \setminus \{0, 1, 2\}$ , the system  $C$  is solvable in positive integers  $x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}$  if and only if

$$(x_4 + 2 = x_9) \wedge (x_4 | ((x_4 - 1)! + 1)) \wedge (x_9 | ((x_9 - 1)! + 1))$$

Hence, the claim of Lemma 12 follows from Lemma 5.  $\square$

**Lemma 13.** *There are only finitely many tuples  $(x_1, \dots, x_{16}) \in (\mathbb{N} \setminus \{0\})^{16}$  which solve the system  $C$  and satisfy  $(x_4 \in \{1, 2\}) \vee (x_9 \in \{1, 2\})$ .*

*Proof.* If a tuple  $(x_1, \dots, x_{16}) \in (\mathbb{N} \setminus \{0\})^{16}$  solves the system  $C$  and  $(x_4 \in \{1, 2\}) \vee (x_9 \in \{1, 2\})$ , then  $x_1, \dots, x_{16} \leq 7!$ . Indeed, for example, if  $x_4 = 2$  then  $x_6 = x_4 + 1 = 3$ . Hence,  $x_7 = x_6! = 6$ . Therefore,  $x_{15} = x_7 + 1 = 7$ . Consequently,  $x_{16} = x_{15}! = 7!$ .  $\square$

**Theorem 9.** (cf. Theorem 13) *The statement  $\Psi_{16}$  proves the following implication: if there exists a twin prime greater than  $g(14)$ , then there are infinitely many twin primes.*

*Proof.* Suppose that the antecedent holds. Then, there exist prime numbers  $x_4$  and  $x_9$  such that  $x_9 = x_4 + 2 > g(14)$ . Hence,  $x_4, x_9 \in \mathbb{N} \setminus \{0, 1, 2\}$ . By Lemma 12, there exists a unique tuple

$$(x_1, x_2, x_3, x_5, x_6, x_7, x_8, x_{10}, x_{11}, x_{12}, x_{13}, x_{14}, x_{15}, x_{16}) \in (\mathbb{N} \setminus \{0\})^{14}$$

such that the tuple  $(x_1, \dots, x_{16})$  solves the system  $C$ . Since  $x_9 > g(14)$ , we obtain that  $x_9 - 1 \geq g(14)$ . Therefore,  $(x_9 - 1)! \geq g(14)! = g(15)$ . Hence,  $(x_9 - 1)! + 1 > g(15)$ . Consequently,

$$x_{16} = ((x_9 - 1)! + 1)! > g(15)! = g(16)$$

Since  $C \subseteq B_{16}$ , the statement  $\Psi_{16}$  and the inequality  $x_{16} > g(16)$  imply that the system  $C$  has infinitely many solutions in positive integers  $x_1, \dots, x_{16}$ . According to Lemmas 12 and 13, there are infinitely many twin primes.  $\square$

**Corollary 6.** (cf. [6]) *Let  $\mathcal{X}_{16}$  denote the set of twin primes. The statement  $\Psi_{16}$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{X}_{16}$ , and this number equals  $\max(\mathcal{X}_{16})$ , if  $\mathcal{X}_{16}$  is finite. Assuming the statement  $\Psi_{16}$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{X}_{16}$ . Assuming the statement  $\Psi_{16}$ , the infinity of  $\mathcal{X}_{16}$  is decidable in the limit.*


*Proof.* We consider an algorithm which computes  $\max(\mathcal{X}_{16} \cap [1, g(14)])$ .  $\square$

## 10 Hypothetical statements $\Delta_5, \dots, \Delta_{14}$ and their consequences

Let  $\lambda(5) = \Gamma(25)$ , and let  $\lambda(n + 1) = \Gamma(\lambda(n))$  for every integer  $n \geq 5$ . For an integer  $n \geq 5$ , let  $\mathcal{J}_n$  denote the following system of equations:

$$\left\{ \begin{array}{l} \forall i \in \{1, \dots, n-1\} \setminus \{3\} \Gamma(x_i) = x_{i+1} \\ x_1 \cdot x_1 = x_4 \\ x_2 \cdot x_3 = x_5 \end{array} \right.$$

Lemma 3 and the diagram in Figure 5 explain the construction of the system  $\mathcal{J}_n$ .



**Fig. 5** Construction of the system  $\mathcal{J}_n$

For every integer  $n \geq 5$ , the system  $\mathcal{J}_n$  has exactly two solutions in positive integers, namely  $(1, \dots, 1)$  and  $(5, 24, 23!, 25, \lambda(5), \dots, \lambda(n))$ . For an integer  $n \geq 5$ , let  $\Delta_n$  denote the following statement: if a system of equations  $\mathcal{S} \subseteq \{\Gamma(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq \lambda(n)$ .

**Hypothesis 2.** The statements  $\Delta_5, \dots, \Delta_{14}$  are true.

**Proposition 4.** Lemmas 3 and 5 imply that the statements  $\Delta_n$  have similar consequences as the statements  $\Psi_n$ .

**Remark 3.** By Lemma 3 and algebraic lemmas in [22, p. 110], the statement  $\forall n \in \mathbb{N} \setminus \{0, 1, 2, 3, 4\} \Delta_n$  implies that there is an algorithm which takes as input a factorial Diophantine equation, and returns an integer such that this integer is greater than the solutions in positive integers, if these solutions form a finite set. This conclusion is a bit strange because a computable upper bound on non-negative integer solutions does not exist for exponential Diophantine equations with a finite number of solutions, see [14, p. 300]. Therefore, the statement  $\forall n \in \mathbb{N} \setminus \{0, 1, 2, 3, 4\} \Delta_n$  seems to be false.

**Theorem 10.** The statement  $\Delta_6$  implies that any prime number  $p \geq 25$  proves the infinitude of primes.

*Proof.* It follows from Lemmas 3 and 5. We leave the details to the reader.  $\square$

## 11 Hypothetical statements $\Sigma_3, \dots, \Sigma_{16}$ and their consequences

Let  $\Gamma_n(k)$  denote  $(k-1)!$ , where  $n \in \{3, \dots, 16\}$  and  $k \in \{2\} \cup [2^{2^{n-3}} + 1, \infty) \cap \mathbb{N}$ . For an integer  $n \in \{3, \dots, 16\}$ , let

$$Q_n = \{\Gamma_n(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$$

For an integer  $n \in \{3, \dots, 16\}$ , let  $P_n$  denote the following system of equations:

$$\begin{cases} x_1 \cdot x_1 = x_1 \\ \Gamma_n(x_2) = x_1 \\ \forall i \in \{2, \dots, n-1\} x_i \cdot x_i = x_{i+1} \end{cases}$$


**Lemma 14.** For every integer  $n \in \{3, \dots, 16\}$ ,  $P_n \subseteq Q_n$  and the system  $P_n$  with  $\Gamma$  instead of  $\Gamma_n$  has exactly one solution in positive integers  $x_1, \dots, x_n$ , namely  $(1, 2^{2^0}, 2^{2^1}, 2^{2^2}, \dots, 2^{2^{n-2}})$ .

For an integer  $n \in \{3, \dots, 16\}$ , let  $\Sigma_n$  denote the following statement: if a system of equations  $\mathcal{S} \subseteq Q_n$  with  $\Gamma$  instead of  $\Gamma_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then every tuple  $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$  that solves the original system  $\mathcal{S}$  satisfies  $x_1, \dots, x_n \leq 2^{2^{n-2}}$ .

**Hypothesis 3.** The statements  $\Sigma_3, \dots, \Sigma_{16}$  are true.

**Lemma 15.** (cf. Lemma 3) For every integer  $n \in \{4, \dots, 16\}$  and for every positive integers  $x$  and  $y$ ,  $x \cdot \Gamma_n(x) = \Gamma_n(y)$  if and only if  $(x + 1 = y) \wedge (x \geq 2^{2^{n-3}} + 1)$ .

Let  $\mathcal{Z}_9 \subseteq Q_9$  be the system of equations in Figure 6.



**Fig. 6** Construction of the system  $\mathcal{Z}_9$

**Lemma 16.** For every positive integer  $x_1$ , the system  $\mathcal{Z}_9$  is solvable in positive integers  $x_2, \dots, x_9$  if and only if  $x_1 > 2^{2^{9-4}}$  and  $x_1^2 + 1$  is prime. In this case, positive integers  $x_2, \dots, x_9$  are uniquely determined by  $x_1$ . For every positive integer  $n$ , at most finitely many tuples  $(x_1, \dots, x_9) \in (\mathbb{N} \setminus \{0\})^9$  begin with  $n$  and solve the system  $\mathcal{Z}_9$  with  $\Gamma$  instead of  $\Gamma_9$ .

*Proof.* It follows from Lemmas 3, 5, and 15. □

**Lemma 17.** ([21]). The number  $(13!)^2 + 1 = 38775788043632640001$  is prime.

**Lemma 18.**  $\left( (13!)^2 \geq 2^{2^{9-3}} + 1 = 18446744073709551617 \right) \wedge \left( \Gamma_9((13!)^2) > 2^{2^{9-2}} \right)$ .

**Theorem 11.** (cf. Theorem 7) The statement  $\Sigma_9$  implies the infinitude of primes of the form  $n^2 + 1$ .

*Proof.* It follows from Lemmas 16–18. □


**Theorem 12.** (cf. Theorem 8) The statement  $\Sigma_9$  implies that any prime of the form  $n! + 1$  with  $n \geq 2^{2^{9-3}}$  proves the infinitude of primes of the form  $n! + 1$ .

*Proof.* We leave the proof to the reader. □

**Corollary 7.** Let  $\mathcal{Y}_9$  denote the set of primes of the form  $n! + 1$ . The statement  $\Sigma_9$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{Y}_9$ , and this number equals  $\max(\mathcal{Y}_9)$ , if  $\mathcal{Y}_9$  is finite. Assuming the statement  $\Sigma_9$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{Y}_9$ . Assuming the statement  $\Sigma_9$ , the infinity of  $\mathcal{Y}_9$  is decidable in the limit.

*Proof.* We consider an algorithm which computes  $\max(\mathcal{Y}_9 \cap [1, (2^{2^{9-3}} - 1)! + 1])$ . □

Let  $\mathcal{Z}_{14} \subseteq \mathcal{Q}_{14}$  be the system of equations in Figure 7.



**Fig. 7** Construction of the system  $\mathcal{Z}_{14}$

**Lemma 19.** *For every positive integer  $x_1$ , the system  $\mathcal{Z}_{14}$  is solvable in positive integers  $x_2, \dots, x_{14}$  if and only if  $x_1$  and  $x_1 + 2$  are prime and  $x_1 \geq 2^{2^{14-3}} + 1$ . In this case, positive integers  $x_2, \dots, x_{14}$  are uniquely determined by  $x_1$ . For every positive integer  $n$ , at most finitely many tuples  $(x_1, \dots, x_{14}) \in (\mathbb{N} \setminus \{0\})^{14}$  begin with  $n$  and solve the system  $\mathcal{Z}_{14}$  with  $\Gamma$  instead of  $\Gamma_{14}$ .*

*Proof.* It follows from Lemmas 3, 5, and 15. □


**Lemma 20.** ([26, p. 87]) *The numbers  $459 \cdot 2^{8529} - 1$  and  $459 \cdot 2^{8529} + 1$  are prime (Harvey Dubner).*

**Lemma 21.**  $459 \cdot 2^{8529} - 1 > 2^{2^{14-2}} = 2^{4096}$ .

**Theorem 13.** (cf. Theorem 9) *The statement  $\Sigma_{14}$  implies the infinitude of twin primes.*

*Proof.* It follows from Lemmas 19–21. □

A prime  $p$  is said to be a Sophie Germain prime if both  $p$  and  $2p + 1$  are prime, see [25]. It is conjectured that there are infinitely many Sophie Germain primes, see [19, p. 330]. Let  $\mathcal{Z}_{16} \subseteq \mathcal{Q}_{16}$  be the system of equations in Figure 8.



**Fig. 8** Construction of the system  $\mathcal{Z}_{16}$

**Lemma 22.** For every positive integer  $x_1$ , the system  $\mathcal{Z}_{16}$  is solvable in positive integers  $x_2, \dots, x_{16}$  if and only if  $x_1$  is a Sophie Germain prime and  $x_1 \geq 2^{2^{16-3}} + 1$ . In this case, positive integers  $x_2, \dots, x_{16}$  are uniquely determined by  $x_1$ . For every positive integer  $n$ , at most finitely many tuples  $(x_1, \dots, x_{16}) \in (\mathbb{N} \setminus \{0\})^{16}$  begin with  $n$  and solve the system  $\mathcal{Z}_{16}$  with  $\Gamma$  instead of  $\Gamma_{16}$ .

*Proof.* It follows from Lemmas 3, 5, and 15. □

**Lemma 23.** ([19, p. 330])  $8069496435 \cdot 10^{5072} - 1$  is a Sophie Germain prime (Harvey Dubner).

**Lemma 24.**  $8069496435 \cdot 10^{5072} - 1 > 2^{2^{16-2}}$ .

**Theorem 14.** The statement  $\Sigma_{16}$  implies the infinitude of Sophie Germain primes.

*Proof.* It follows from Lemmas 22–24. □

**Theorem 15.** The statement  $\Sigma_6$  proves the following implication: if the equation  $x(x+1) = y!$  has only finitely many solutions in positive integers  $x$  and  $y$ , then each such solution  $(x, y)$  belongs to the set  $\{(1, 2), (2, 3)\}$ .

*Proof.* We leave the proof to the reader. □

The question of solving the equation  $x(x+1) = y!$  was posed by P. Erdős, see [2]. F. Luca proved that the *abc* conjecture implies that the equation  $x(x+1) = y!$  has only finitely many solutions in positive integers, see [13].

**Theorem 16.** The statement  $\Sigma_6$  proves the following implication: if the equation  $x! + 1 = y^2$  has only finitely many solutions in positive integers  $x$  and  $y$ , then each such solution  $(x, y)$  belongs to the set  $\{(4, 5), (5, 11), (7, 71)\}$ .

*Proof.* We leave the proof to the reader. □

## 12 Hypothetical statements $\Omega_3, \dots, \Omega_{16}$ and their consequences

For an integer  $n \in \{3, \dots, 16\}$ , let  $\Omega_n$  denote the following statement: *if a system of equations  $\mathcal{S} \subseteq \{\Gamma(x_i) = x_k : i, k \in \{1, \dots, n\}\} \cup \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\}$  has a solution in integers  $x_1, \dots, x_n$  greater than  $2^{2^{n-2}}$ , then  $\mathcal{S}$  has infinitely many solutions in positive integers  $x_1, \dots, x_n$ .* For every  $n \in \{3, \dots, 16\}$ , the statement  $\Sigma_n$  implies the statement  $\Omega_n$ .

**Lemma 25.** *The number  $(65!)^2 + 1$  is prime and  $65! > 2^{2^{9-2}}$ .*

*Proof.* The following PARI/GP ([18]) command

```
(04:04) gp > isprime((65!)^2+1, {flag=2})
%1 = 1
```

is shown together with its output. This command performs the APRCL primality test, the best deterministic primality test algorithm ([26, p. 226]). It rigorously shows that the number  $(65!)^2 + 1$  is prime.  $\square$

**Lemma 26.** *If positive integers  $x_1, \dots, x_9$  solve the system  $\mathcal{Z}_9$  and  $x_1 > 2^{2^{9-2}}$ , then  $x_1 = \min(x_1, \dots, x_9)$ .*

**Theorem 17.** *The statement  $\Omega_9$  implies the infinitude of primes of the form  $n^2 + 1$ .*

*Proof.* It follows from Lemmas 16 and 25–26.  $\square$

**Lemma 27.** *If positive integers  $x_1, \dots, x_{14}$  solve the system  $\mathcal{Z}_{14}$  and  $x_1 > 2^{2^{14-2}}$ , then  $x_1 = \min(x_1, \dots, x_{14})$ .*

**Theorem 18.** *The statement  $\Omega_{14}$  implies the infinitude of twin primes.*

*Proof.* It follows from Lemmas 19–21 and 27.  $\square$

## 13 Are there infinitely many composite Fermat numbers?

Integers of the form  $2^{2^n} + 1$  are called Fermat numbers. Primes of the form  $2^{2^n} + 1$  are called Fermat primes, as Fermat conjectured that every integer of the form  $2^{2^n} + 1$  is prime, see [12, p. 1]. Fermat correctly remarked that  $2^{2^0} + 1 = 3$ ,  $2^{2^1} + 1 = 5$ ,  $2^{2^2} + 1 = 17$ ,  $2^{2^3} + 1 = 257$ , and  $2^{2^4} + 1 = 65537$  are all prime, see [12, p. 1].

**Open Problem 3.** ([12, p. 159]) *Are there infinitely many composite numbers of the form  $2^{2^n} + 1$ ?*

Most mathematicians believe that  $2^{2^n} + 1$  is composite for every integer  $n \geq 5$ , see [11, p. 23]. Let

$$H_n = \{x_i \cdot x_j = x_k : i, j, k \in \{1, \dots, n\}\} \cup \{2^{2^{x_i}} = x_k : i, k \in \{1, \dots, n\}\}$$

Let  $h(1) = 1$ , and let  $h(n+1) = 2^{2^{h(n)}}$  for every positive integer  $n$ .

**Lemma 28.** *The following subsystem of  $H_n$*

$$\begin{cases} x_1 \cdot x_1 = x_1 \\ \forall i \in \{1, \dots, n-1\} 2^{2^{x_i}} = x_{i+1} \end{cases}$$

*has exactly one solution  $(x_1, \dots, x_n) \in (\mathbb{N} \setminus \{0\})^n$ , namely  $(h(1), \dots, h(n))$ .*

For a positive integer  $n$ , let  $\xi_n$  denote the following statement: *if a system of equations  $S \subseteq H_n$  has only finitely many solutions in positive integers  $x_1, \dots, x_n$ , then each such solution  $(x_1, \dots, x_n)$  satisfies  $x_1, \dots, x_n \leq h(n)$* . The statement  $\xi_n$  says that for subsystems of  $H_n$  the largest known solution is indeed the largest possible.

**Hypothesis 4.** *The statements  $\xi_1, \dots, \xi_{13}$  are true.*

**Lemma 29.** *Every statement  $\xi_n$  is true with an unknown integer bound that depends on  $n$ .*


*Proof.* For every positive integer  $n$ , the system  $H_n$  has a finite number of subsystems. □

**Theorem 19.** *The statement  $\xi_{13}$  proves the following implication: if  $z \in \mathbb{N} \setminus \{0\}$  and  $2^{2^z} + 1$  is composite and greater than  $h(12)$ , then  $2^{2^z} + 1$  is composite for infinitely many positive integers  $z$ .*

*Proof.* Let us consider the equation

$$(x + 1)(y + 1) = 2^{2^z} + 1 \tag{E}$$

in positive integers. By Lemma 4, we can transform the equation (E) into an equivalent system of equations  $\mathcal{G}$  which has 13 variables ( $x, y, z$ , and 10 other variables) and which consists of equations of the forms  $\alpha \cdot \beta = \gamma$  and  $2^{2^\alpha} = \gamma$ , see the diagram in Figure 9.



**Fig. 9** Construction of the system  $\mathcal{G}$

Since  $2^{2^z} + 1 > h(12)$ , we obtain that  $2^{2^{2^{2^z} + 1}} > h(13)$ . By this, the statement  $\xi_{13}$  implies that the system  $\mathcal{G}$  has infinitely many solutions in positive integers. It means that there are infinitely many composite Fermat numbers. □

**Corollary 8.** *Let  $\mathcal{W}_{13}$  denote the set of composite Fermat numbers. The statement  $\xi_{13}$  implies that we know an algorithm such that it returns a threshold number of  $\mathcal{W}_{13}$ , and this number equals  $\max(\mathcal{W}_{13})$ , if  $\mathcal{W}_{13}$  is finite. Assuming the statement  $\xi_{13}$ , a single query to an oracle for the halting problem decides the infinity of  $\mathcal{W}_{13}$ . Assuming the statement  $\xi_{13}$ , the infinity of  $\mathcal{W}_{13}$  is decidable in the limit.*

*Proof.* We consider an algorithm which computes  $\max(\mathcal{W}_{13} \cap [1, h(12)])$ . □

## References

- [1] C. H. Bennett, *Chaitin's Omega*, in: *Fractal music, hypercards, and more ...* (M. Gardner, ed.), W. H. Freeman, New York, 1992, 307–319.
- [2] D. Berend and J. E. Harmse, *On polynomial-factorial Diophantine equations*, *Trans. Amer. Math. Soc.* 358 (2006), no. 4, 1741–1779.
- [3] C. K. Caldwell and Y. Gallot, *On the primality of  $n! \pm 1$  and  $2 \times 3 \times 5 \times \cdots \times p \pm 1$* , *Math. Comp.* 71 (2002), no. 237, 441–448, <http://doi.org/10.1090/S0025-5718-01-01315-1>.
- [4] C. S. Calude, H. Jürgensen, S. Legg, *Solving problems with finite test sets*, in: *Finite versus Infinite: Contributions to an Eternal Dilemma* (C. Calude and G. Păun, eds.), 39–52, Springer, London, 2000.
- [5] N. C. A. da Costa and F. A. Doria, *On the foundations of science (LIVRO): essays, first series*, E-papers Serviços Editoriais Ltda, Rio de Janeiro, 2013.
- [6] F. G. Dorais, *Can the twin prime problem be solved with a single use of a halting oracle?* July 23, 2011, <http://mathoverflow.net/questions/71050>.
- [7] W. B. Easton, *Powers of regular cardinals*, *Ann. Math. Logic* 1 (1970), 139–178.
- [8] M. Erickson, A. Vazzana, D. Garth, *Introduction to number theory*, 2nd ed., CRC Press, Boca Raton, FL, 2016.
- [9] H. Friedman, *The incompleteness phenomena*, in: *Proceedings of the AMS Centennial Symposium 1988*, 49–84, Amer. Math. Soc., Providence, RI, 1992.
- [10] T. Jech, *Set theory*, Springer, Berlin, 2003.
- [11] J.-M. De Koninck and F. Luca, *Analytic number theory: Exploring the anatomy of integers*, American Mathematical Society, Providence, RI, 2012.
- [12] M. Křížek, F. Luca, L. Somer, *17 lectures on Fermat numbers: from number theory to geometry*, Springer, New York, 2001.
- [13] F. Luca, *The Diophantine equation  $P(x) = n!$  and a result of M. Overholt*, *Glas. Mat. Ser. III* 37 (57) (2002), no. 2, 269–273
- [14] Yu. Matiyasevich, *Existence of noneffectivizable estimates in the theory of exponential Diophantine equations*, *J. Sov. Math.* vol. 8, no. 3, 1977, 299–311, <http://dx.doi.org/10.1007/bf01091549>.
- [15] M. Mignotte and A. Pethő, *On the Diophantine equation  $x^p - x = y^q - y$* , *Publ. Mat.* 43 (1999), no. 1, 207–216.
- [16] W. Narkiewicz, *Rational number theory in the 20th century: From PNT to FLT*, Springer, London, 2012.
- [17] M. Overholt, *The Diophantine equation  $n! + 1 = m^2$* , *Bull. London Math. Soc.* 25 (1993), no. 2, 104.
- [18] PARI/GP *online documentation*, [http://pari.math.u-bordeaux.fr/dochtml/html/Arithmetic\\_functions.html](http://pari.math.u-bordeaux.fr/dochtml/html/Arithmetic_functions.html)
- [19] P. Ribenboim, *The new book of prime number records*, Springer, New York, 1996, <http://doi.org/10.1007/978-1-4612-0759-7>.



- [20] S. Siksek, *Chabauty and the Mordell–Weil Sieve*, in: *Advances on Superelliptic Curves and Their Applications* (eds. L. Beshaj, T. Shaska, E. Zhupa), 194–224, IOS Press, Amsterdam, 2015, <http://dx.doi.org/10.3233/978-1-61499-520-3-194>.
- [21] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*, *Smallest prime factor of  $A020549(n) = (n!)^2 + 1$* , <http://oeis.org/A282706>.
- [22] A. Tyszka, *A hypothetical upper bound on the heights of the solutions of a Diophantine equation with a finite number of solutions*, *Open Comput. Sci.* 8 (2018), no. 1, 109–114, <http://doi.org/10.1515/comp-2018-0012>.
- [23] E. W. Weisstein, *CRC Concise Encyclopedia of Mathematics*, 2nd ed., Chapman & Hall/CRC, Boca Raton, FL, 2002.
- [24] Wolfram MathWorld, *Perfect Cuboid*, <http://mathworld.wolfram.com/PerfectCuboid.html>.
- [25] Wolfram MathWorld, *Sophie Germain prime*, <http://mathworld.wolfram.com/SophieGermainPrime.html>.
- [26] S. Y. Yan, *Number theory for computing*, 2nd ed., Springer, Berlin, 2002.
- [27] A. A. Zenkin, *Super-induction method: logical acupuncture of mathematical infinity*, Twentieth World Congress of Philosophy, Boston, MA, August 10–15, 1998, <http://www.bu.edu/wcp/Papers/Logi/LogiZenk.htm>.
- [28] A. A. Zenkin, *Superinduction: new logical method for mathematical proofs with a computer*, in: J. Cachro and K. Kijania-Placek (eds.), *Volume of Abstracts, 11th International Congress of Logic, Methodology and Philosophy of Science*, August 20–26, 1999, Cracow, Poland, p. 94, The Faculty of Philosophy, Jagiellonian University, Cracow, 1999.

Apoloniusz Tyszka  
 University of Agriculture  
 Faculty of Production and Power Engineering  
 Balicka 116B, 30-149 Kraków, Poland  
 E-mail: [rtyyszka@cyf-kr.edu.pl](mailto:rtyyszka@cyf-kr.edu.pl)